

Risk of cancer in small and diminutive colorectal polyps

Prasanna L. Ponugoti, M.D.¹

Oscar W. Cummings, M.D.²

Douglas K. Rex, M.D.¹

Indiana University School of Medicine,

¹Division of Gastroenterology and Hepatology, Indianapolis, IN

²Division of Surgical Pathology, Indianapolis, IN

Address correspondence and reprint requests to:

Douglas K. Rex

550 N. University Blvd, Suite 4100

Indianapolis, IN 46202

Phone: 317-948-8741

Fax: 917-944-5449

Email: drex@iupui.edu

This work was supported by a gift to the Indiana University Foundation in the name of Douglas K. Rex by Scott and Kay Schurz of Bloomington, Indiana, and their children.

This is the author's manuscript of the article published in final edited form as:

Ponugoti, P. L., Cummings, O. W., & Rex, D. K. (2017). Risk of cancer in small and diminutive colorectal polyps. *Digestive and Liver Disease*, 49(1), 34–37. <https://doi.org/10.1016/j.dld.2016.06.025>

Abstract

The prevalence of cancer in small and diminutive polyps is relevant to “resect and discard” and CT colonography reporting recommendations.

We evaluated a prospectively collected colonoscopy polyp database to identify polyps < 10 mm and those with cancer or advanced histology (high-grade dysplasia or villous elements)

Of 32,790 colonoscopies, 15,558 colonoscopies detected 42,630 polyps < 10 mm in size. A total of 4,790 lesions were excluded as they were not conventional adenomas or serrated class lesions.

There were 23,524 conventional adenomas < 10 mm of which 22,952 were tubular adenomas.

There were 14,316 serrated class lesions of which 13,589 were hyperplastic polyps and the remainder were sessile serrated polyps. Of all conventional adenomas, 96 had high-grade dysplasia including 0.3% of adenomas \leq 5 mm in size and 0.8% of adenomas 6-9 mm in size. Of all conventional adenomas, 2.1% of those \leq 5 mm in size and 5.6% of those 6-9 mm in size were advanced. Among 36,107 polyps \leq 5 mm in size and 6,523 polyps 6-9 mm in size, there were no cancers.

These results support the safety of resect and discard as well as current CT colonography reporting recommendations for small and diminutive polyps.

Key words: Diminutive polyps, small polyps, colorectal cancer, adenomas, serrated lesions, resect and discard, CT colonography.

INTRODUCTION

The risk of cancer in small and diminutive colorectal polyps remains an important factor in several polyp management paradigms. For example, the risk of cancer in the resect and discard paradigm is important, since resect and discard ¹ could potentially result in a small polyp with cancer being discarded after resection, and the cancer thus going unrecognized. Such an event could result in an adverse outcome for a patient if a cancer recurrence developed. Similarly, the prevalence of cancer in diminutive and small polyps is important in computed tomography (CT) colonography management paradigms, where diminutive polyps are not reported ², and small polyps may not be recommended for immediate resection ².

Early studies of cancer prevalence in small and diminutive polyps found a substantial cancer risk, which exceeded 0.4% in polyps 6 to 9 mm in size in some studies ³⁻⁷. However, more recent colonoscopic studies found a much lower prevalence ⁸⁻¹⁵. These differences may reflect in part the improved potential for polyp identification resulting from progressive improvements in colonoscope imaging. Thus, increasing emphasis on maximizing adenoma detection could result in a broader range of lesions removed, including many with a more subtle and flat morphology than was detectable in early studies. Contrary to some suggestions, flat morphology without depression is not associated with an increased risk of advanced pathology and may be associated with a lower risk of advanced pathology ¹⁶⁻¹⁸.

The prevalence of cancer in diminutive polyps is of great interest to some patients who are considering whether to participate in the resect and discard paradigm of small polyp management ¹⁹. Determining this risk of cancer within narrow confidence limits is an important goal. We now report the largest single experience on the prevalence of cancer in small and diminutive polyps. This study exceeds the size of all prior studies on this topic combined.

MATERIALS AND METHODS

We prospectively maintained a database of colonoscopies and polyp findings continuously since the year 2000 in our endoscopy units. The current report describes colonoscopies performed beginning in February 2004 and extending to September 2015. Permission to review the database was granted by the Institutional Review Board at Indiana University School of Medicine on September 29, 2015.

The database includes the endoscopist performing the examination, polyp size (as measured by endoscopist estimate), polyp location in the colon (by endoscopist estimate), and pathology as reported by our university pathologists.

Conventional adenomas were those interpreted by the pathologists as tubular, tubulovillous, or villous. Serrated class lesions were those interpreted as hyperplastic polyp, sessile serrated polyp, sessile serrated adenoma, or serrated adenoma, or traditional serrated adenoma.

Statistical analysis was descriptive.

RESULTS

A total of 32,790 colonoscopies were performed during the study period by 53 different endoscopists. There were 15,558 procedures that identified 42,630 polyps < 10 mm in size for which the polyp was resected and a pathology report was created. The mean age was 60.1 years (59.7 years in females, 60.2 years in males). There were 1,890 procedures in persons < 50 years of age, and 649 in persons 80 years and older.

Polyps (n=4,790) that were not considered conventional adenomas or serrated class lesions were excluded from further analysis. These included normal tissue/mucosa (n = 3,833),

inflammatory polyps (n = 574), lymphoid follicles (n = 293), granulation tissue (n = 60), hamartoma (n = 14), carcinoid tumor (n=13), metastatic malignant melanoma (n = 2), and granular cell tumor (n = 1).

There were 23,524 conventional adenomas < 10 mm in size, of which 22,952 (97.6%) were tubular adenomas. There were 14,316 serrated class lesions < 10 mm in size, of which 13,589 (95%) were interpreted as hyperplastic polyps, and the remainder were sessile serrated polyps or "serrated adenoma" (Table 1). Polyps < 10 mm in size comprised 88.4% of all resected polyps.

Among all conventional adenomas, 96 had high-grade dysplasia, including 0.3% of adenomas \leq 5 mm in size and 0.8% of conventional adenomas 6 to 9 mm in size. The fraction of conventional adenomas \leq 5 mm in size that were advanced (had either villous elements or high-grade dysplasia) was 2.1%. The fraction of 6-9 mm conventional adenomas that were advanced was 5.6%.

There were no cancers among 36,107 polyps \leq 5 mm in size and 6,523 polyps 6 to 9 mm in size.

DISCUSSION

In this report, we describe the largest reported experience with identification of cancer in colorectal polyps < 10 mm in size. This study exceeds the size of all previous studies on this topic combined (Table 2). We found no cancers in nearly 36,000 polyps \leq 5 mm in size, and none in 6,523 polyps 6 to 9 mm in size. Our results indicate that the risk of cancer in diminutive and small polyps is very low. These results substantially improve the projected safety of resect and discard paradigm¹ and of current reporting recommendations for CT colonography².

Our results are consistent with much of the previously published literature^{3-15,20-27}, particularly reports from the last 10 years (Table 2). As noted earlier, the trends toward decreasing prevalence rates of cancer in small and diminutive polyps may reflect the improved imaging capabilities of colonoscopes, and increasing emphasis on identification of flat lesions which do not increase the risk of cancer compared to polypoid lesions¹⁶⁻¹⁸.

The overall distribution of polyp sizes in this study suggests that the tendency of endoscopists in our unit is to underestimate polyp size, particularly since the percentage of all polyps < 1 cm in size was comparable to other studies^{3-15,21-27}, despite our center receiving many referrals for large and complex polypectomies²⁸. Further the fraction of polyps \leq 5 mm and 6-9 mm in size was comparable to or higher than prior studies^{3-17,22-29}. Any underestimation of polyp size would tend to cause the apparent prevalence of cancer in polyps of different size ranges to increase. Despite this apparent tendency to underestimate polyp size, no cancers were identified in small and diminutive polyps.

The prevalence of villous elements and high-grade dysplasia in small and diminutive polyps in the study was very low, and lower than noted in a number of other reports (Table 2). However, we have demonstrated that interpretation of villous elements and dysplasia grade in small polyps is subject to extreme interobserver variation and poor agreement even among experts²⁹. These features have caused the British Society of Gastroenterology to ignore villous elements and dysplasia grade in their post polypectomy surveillance guideline³⁰. We previously reported that experts in gastrointestinal pathology do not use the same definitions of high-grade dysplasia²⁹. Specifically, pathologists who utilize cytologic criteria for high-grade dysplasia report much higher prevalence rates of high-grade dysplasia compared to those who use only morphologic criteria²⁹. At the direction of the gastrointestinal pathologists in our unit, our

pathologists utilize only morphologic criteria for high-grade dysplasia, which undoubtedly contributes to the low prevalence of high-grade dysplasia in our study. Similarly, there is marked interobserver variation in the interpretation of sessile serrated polyp vs. hyperplastic polyp^{31,32}. We had previously demonstrated that experts identify a higher percentage of sessile serrated polyps than are identified by our own pathologists³². However, experts also exhibit significant interobserver variation³¹. Therefore, our data on the prevalence of sessile serrated polyp reflect a conservative approach to interpretation of this lesion, as well as evolving awareness of sessile serrated polyp by pathologists during the study period. Our focus in this paper is on the prevalence of cancer in small and diminutive polyps. Interpretation of cancer by community pathologists in colorectal polyps appears to be consistent and reliable³³, as does the assignment of colorectal polyps to the conventional adenoma vs. serrated class^{1,33}. Thus, the pathology reports with regards to the prevalence of cancer in diminutive and small polyps should be reliable.

Strengths of our study include its large size, which far exceeds the size of any previous study on this topic, and is slightly larger than the size of all previous studies on this topic combined (Table 2). There are few databases available that include pathology results on individual polyps < 10 mm in size.

Limitations include the retrospective nature of the study, though the database was accumulated prospectively. Additionally, polyp size was estimated endoscopically, though as we noted above the distribution of polyp sizes does not suggest any tendency to overestimate lesion size.

In summary, we did not identify any cancers in a large sample of diminutive and small colorectal polyps. This result favors the feasibility of strategies such as resect and discard¹, and

favors the appropriateness of current CT colonography reporting recommendations for diminutive and small polyps ².

References

1. Rex DK, Kahi C, O'Brien M, et al. The American Society for Gastrointestinal Endoscopy PIVI (Preservation and Incorporation of Valuable Endoscopic Innovations) on real-time endoscopic assessment of the histology of diminutive colorectal polyps. *Gastrointest Endosc* 2011;73:419-22.
2. Zalis ME, Barish MA, Choi JR, et al. CT colonography reporting and data system: a consensus proposal. *Radiology* 2005;236:3-9.
3. Shinya H, Wolff WI. Morphology, anatomic distribution and cancer potential of colonic polyps. *Ann Surg* 1979;190:679-83.
4. Gschwantler M, Kriwanek S, Langner E, et al. High-grade dysplasia and invasive carcinoma in colorectal adenomas: a multivariate analysis of the impact of adenoma and patient characteristics. *Eur J Gastroenterol Hepatol* 2002;14:183-8.
5. Odom SR, Duffy SD, Barone JE, Ghevariya V, McClane SJ. The rate of adenocarcinoma in endoscopically removed colorectal polyps. *Am Surg* 2005;71:1024-6.
6. Yoo TW, Park DI, Kim YH, et al. Clinical significance of small colorectal adenoma less than 10 mm: the KASID study. *Hepatogastroenterology* 2007;54:418-21.
7. Butterly LF, Chase MP, Pohl H, Fiarman GS. Prevalence of clinically important histology in small adenomas. *Clinical gastroenterology and hepatology : the official clinical practice journal of the American Gastroenterological Association* 2006;4:343-8.
8. Kim DH, Pickhardt PJ, Taylor AJ. Characteristics of advanced adenomas detected at CT colonographic screening: implications for appropriate polyp size thresholds for polypectomy versus surveillance. *AJR Am J Roentgenol* 2007;188:940-4.
9. Lieberman D, Moravec M, Holub J, Michaels L, Eisen G. Polyp size and advanced histology in patients undergoing colonoscopy screening: implications for CT colonography. *Gastroenterology* 2008;135:1100-5.
10. Graser A, Stieber P, Nagel D, et al. Comparison of CT colonography, colonoscopy, sigmoidoscopy and faecal occult blood tests for the detection of advanced adenoma in an average risk population. *Gut* 2009;58:241-8.
11. Bretagne JF, Manfredi S, Piette C, Hamonic S, Durand G, Riou F. Yield of high-grade dysplasia based on polyp size detected at colonoscopy: a series of 2295 examinations following a positive fecal occult blood test in a population-based study. *Dis Colon Rectum* 2010;53:339-45.
12. Chaput U, Alberto SF, Terris B, et al. Risk factors for advanced adenomas amongst small and diminutive colorectal polyps: a prospective monocenter study. *Dig Liver Dis* 2011;43:609-12.
13. Tsai FC, Strum WB. Prevalence of advanced adenomas in small and diminutive colon polyps using direct measurement of size. *Dig Dis Sci* 2011;56:2384-8.
14. Denis B, Bottlaender J, Weiss AM, Peter A, Breysacher G, Chiappa P, Gendre I, Perrin P. Some diminutive colorectal polyps can be removed and discarded without pathologic examination. *Endoscopy* 2011;43:81-6.
15. Gupta N, Bansal A, Rao D, et al. Prevalence of advanced histological features in diminutive and small colon polyps. *Gastrointest Endosc* 2012;75:1022-30.
16. Kudo S, Lambert R, Allen JI, et al. Nonpolypoid neoplastic lesions of the colorectal mucosa. *Gastrointest Endosc* 2008;68(4 Suppl):S3-47.
17. Bianco MA, Cipolletta L, Rotondano G, et al. Prevalence of nonpolypoid colorectal neoplasia: an Italian multicenter observational study. *Endoscopy* 2010;42:279-85.
18. Tsuda S, Veress B, Toth E, Fork F. Flat and depressed colorectal tumours in southern Swedish population: a prospective chromoendoscopic and histopathological study. *Gut* 2002;51:550-5.
19. Rex DK, Patel NJ, Vemulapalli KC. A survey of patient acceptance of resect and discard for diminutive polyps. *Gastrointest Endosc* 2015;82:376-80 e1.

20. O'Brien MJ, Winawer SJ, Zauber AG, et al. The National Polyp Study. Patient and polyp characteristics associated with high-grade dysplasia in colorectal adenomas. *Gastroenterology* 1990;98:371-9.
21. Weston AP, Campbell DR. Diminutive colonic polyps: histopathology, spatial distribution, concomitant significant lesions, and treatment complications. *Am J Gastroenterol* 1995;90:24-8.
22. Aldridge AJ, Simson JN. Histological assessment of colorectal adenomas by size. Are polyps less than 10 mm in size clinically important? *Eur J Surg* 2001;167:777-81.
23. Kapsoritakis AN, Potamianos SP, Koukourakis MI, et al. Diminutive polyps of large bowel should be an early target for endoscopic treatment. *Dig Liver Dis* 2002;34:137-40.
24. Pickhardt PJ, Choi JR, Hwang I, et al. Computed tomographic virtual colonoscopy to screen for colorectal neoplasia in asymptomatic adults. *N Engl J Med* 2003;349:2191-200.
25. Church JM. Clinical significance of small colorectal polyps. *Dis Colon Rectum* 2004;47:481-5.
26. Sprung DJ. Prevalence of adenocarcinoma in small adenomas. *American Journal of Gastroenterology* 2006;101:S199-S.
27. Kaltenbach T, Rastogi A, Rouse RV, et al. Real-time optical diagnosis for diminutive colorectal polyps using narrow-band imaging: the VALID randomised clinical trial. *Gut* 2015;64:1569-77.
28. Liaquat H, Rohn E, Rex DK. Prophylactic clip closure reduced the risk of delayed postpolypectomy hemorrhage: experience in 277 clipped large sessile or flat colorectal lesions and 247 control lesions. *Gastrointest Endosc* 2013;77:401-7.
29. Lasisi F, Mouchli A, Riddell R, et al. Agreement in interpreting villous elements and dysplasia in adenomas less than one centimetre in size. *Dig Liver Dis* 2013;45:1049-55.
30. Cairns SR, Scholefield JH, Steele RJ, et al. Guidelines for colorectal cancer screening and surveillance in moderate and high risk groups (update from 2002). *Gut* 2010;59:666-89.
31. Khalid O, Radaideh S, Cummings OW, O'Brien MJ, Goldblum JR, Rex DK. Reinterpretation of histology of proximal colon polyps called hyperplastic in 2001. *World J Gastroenterol* 2009;15:3767-70.
32. Abdeljawad K, Vemulapalli KC, Kahi CJ, Cummings OW, Snover DC, Rex DK. Sessile serrated polyp prevalence determined by a colonoscopist with a high lesion detection rate and an experienced pathologist. *Gastrointest Endosc* 2015;81:517-24.
33. Rex DK, Alikhan M, Cummings O, Ulbright TM. Accuracy of pathologic interpretation of colorectal polyps by general pathologists in community practice. *Gastrointest Endosc* 1999;50:468-74.

Table 1. Prevalence of cancer in diminutive and small colorectal polyps

Polyp Pathology	Size	Total number of polyps	Tubular adenoma n (%)	Tubulovillous adenoma n (%)	Villous adenoma n (%)	High-grade Dysplasia n (%)	Cancer n (%)	Hyperplastic polyp n (%)	Sessile serrated polyp n (%)
	≤ 5 mm	19,559	19,191 (98.1%)	357 (1.8%)	11 (0.1%)	66 (0.3%)	0	-	-
	6-9 mm	3,965	3,761 (94.9%)	202 (5.1%)	2 (0.1%)	30 (0.8%)	0	-	-
	≤ 5 mm	12,214	-	-	-	-	0	11,772 (96.4%)	442 (3.6%)
	6-9 mm	2,102	-	-	-	-	0	1,817 (86.4%)	285 (13.6%)

Table 2: Previously published data on the prevalence of advanced adenomas and cancer in small and diminutive polyps

Study	Number of Diminutive Adenomas (Size Range)	Number of Diminutive Adenomas with Advanced Histology (%)	Number of Diminutive Carcinomas (%)	Number of Small Adenomas (Size Range)	Number of Small Adenomas with Advanced Histology (%)	Number of Small Carcinomas (%)
Shinya et al. 1979 ³				1661 (5-9 mm)	249 (15%)**	8 (0.5%)
National Polyp Study, 1990 ²⁰	1270 (<6 mm)	25 (2%)**		1230 (6-10 mm)	155 (12.6%)**	-
Weston et al. 1995 ²¹	1964 (<= 5 mm)	5 (0.3%)	0			
Aldridge et al. 2001 ²²	194 (<= 5 mm)	71 (36.5%)**	0	163 (6-10 mm)	99 (60.7%)**	2 (0.01%)
Gshwantler et al. 2002 ⁴	3016 (<5 mm)	561 (18.6%)**	0	2789 (5-10 mm)	1080 (38.7%)**	26 (0.9%)
Kapsoritakis et al. 2002 ²³	293 (<= 5 mm)	6 (2%)*	0			
Pickhardt et al. 2003 ²⁴	966 (<=5 mm)	1 (0.1%)	0	262 (6-9 mm)		0
Church et al. 2004 ²⁵	4381 (<6 mm)	91 (2.1%)	2 (0.05%)	666 (6-10 mm)	65 (9.8%)	1 (0.15%)
Odom et al. 2005 ⁵	2851 (<= 5 mm)		1 (0.03%)	152 (6-10 mm)		1 (0.65%)
Butterly et al. 2006 ⁷	1305 (<6 mm)	34 (2.6%)**	1 (0.08%)	487 (6-9 mm)	38 (7.8%)**	2 (0.4%)
Sprung et al. 2006 ²⁶				6694 (5-10 mm)		2 (0.03%)
Yoo et al. 2007 ⁶	3303 (<= 5 mm)	5 (0.15%)	1 (<0.01%)	1432 (6-9 mm)	31 (2.2%)	7 (0.49%)
Kim et al. 2007 ⁸	2006 (<=5 mm)	4 (0.2%)	0			
Lieberman et al. 2008 ⁹	3744 (<=5 mm)	62 (1.7%)***	1 (0.02%)	1198 (6-9 mm)	77 (6.4%)***	2 (0.2%)
Graser, 2009 ¹⁰	418 (<= 5 mm)	7 (1.7%)**	0	56 (6-9 mm)	6 (10.7%)**	0
Bretagne et al. 2010 ¹¹	535 (<=5 mm)	15 (2.8%)*	0	219 (6-9 mm)	34 (15.5%)*	0
Chaput et al. 2011 ¹²	342 (<=5 mm)	16 (4.7%)	0	72 (6-9 mm)	25 (35.2%)	0
Tsai et al. 2011 ¹³	1025 (<= 5 mm)	105 (10%)	0	247 (6-9 mm)	67 (27%)	0
Denis et al. 2011 ¹⁴	180 (<= 5 mm)	22 (12.2%)		29 (6-9 mm)	13 (44%)	0
Gupta et al. 2012 ¹⁵	1620 (<= 5 mm)	9 (0.6%)**	0	455 (6-9 mm)	7 (1.5%)**	0
Kaltenbach et al. 2014 ²⁷	975 (<= 5 mm)	1 (0.10%)	0			
Pooled data	30,388	1,040 (3.42%)	6 (0.02%)	17,812	1,946 (10.9%)	51 (0.28%)

*: High grade dysplasia only

** : High grade dysplasia, villous and tubulovillous category

*** : High grade dysplasia, villous and tubulovillous, serrated category