



JMIR Res Protoc. 2015 Oct-Dec; 4(4): e143. Published online 2015 Dec 30.

doi: 10.2196/resprot.4416: 10.2196/resprot.4416

PMCID: PMC4712346 | PMID: [26717906](https://pubmed.ncbi.nlm.nih.gov/26717906/)

Development and Pilot Evaluation of a Tablet-Based Application to Improve Quality of Care in Child Mental Health Treatment

Monitoring Editor: Gunther Eysenbach

Reviewed by Josef Ruzek and Aviv Shachak

[Kenneth J Ruggiero](#), PhD,^{1,2} [Brian E Bunnell](#), MS,¹ [Arthur R Andrews III](#), PhD,³ [Tatiana M Davidson](#), PhD,^{1,3} [Rochelle F Hanson](#), PhD,³ [Carla Kmett Danielson](#), PhD,³ [Benjamin E Saunders](#), PhD,³ [Kathryn Soltis](#), BS,^{1,3} [Caleb Yarian](#),⁴ [Brian Chu](#), PhD,⁵ and [Zachary W Adams](#), PhD³

¹ Technology Applications Center for Healthful Lifestyles, College of Nursing, Medical University of South Carolina, Charleston, SC, United States

² Ralph H Johnson VA Medical Center, Charleston, SC, United States

³ National Crime Victims Research and Treatment Center, Department of Psychiatry and Behavioral Sciences, Medical University of South Carolina, Charleston, SC, United States

⁴ Fuzzco, Charleston, SC, United States

⁵ Rutgers, The State University of New Jersey, Piscataway, NJ, United States

Kenneth J Ruggiero, Technology Applications Center for Healthful Lifestyles, College of Nursing, Medical University of South Carolina, 99 Jonathan Lucas St, Charleston, SC, 29425, United States, Phone: 1 843 792 3687, Fax: 1 843 792 2099, Email: ruggierk@musc.edu.

✉Corresponding author.

Corresponding Author: Kenneth J Ruggiero ruggierk@musc.edu

Received 2015 Mar 11; Revisions requested 2015 Jun 24; Revised 2015 Jun 29; Accepted 2015 Jun 29.

Copyright ©Kenneth J Ruggiero, Brian E Bunnell, Arthur R Andrews III, Tatiana M Davidson, Rochelle F Hanson, Carla Kmett Danielson, Benjamin E Saunders, Kathryn Soltis, Caleb Yarian, Brian Chu, Zachary W Adams. Originally published in JMIR Research Protocols (<http://www.researchprotocols.org>), 30.12.2015.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Research Protocols, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.researchprotocols.org>, as well as this copyright and license information must be included.

Abstract

Background

Children need access to high quality mental health care. Effective treatments now exist for a wide range of mental health conditions. However, these interventions are delivered with variable effectiveness in traditional mental health service settings. Innovative solutions are needed to improve treatment delivery quality and effectiveness.

Objective

The aim of this study was to develop a scalable, sustainable technology-based approach to improve the quality of care in child mental health treatment.

Methods

A tablet-based resource was developed with input from mental health training experts, mental health providers, and patients. A series of qualitative data collection phases (ie, expert interviews, patient and provider focus groups, usability testing) guided the initial concept and design of the resource, and then its refinement. The result was an iPad-based “e-workbook” designed to improve child engagement and provider fidelity in implementation of a best-practice treatment. We are currently conducting a small scale randomized controlled trial to evaluate the feasibility of e-workbook facilitated child mental health treatment with 10 providers and 20 families recruited from 4 local community-based mental health clinics.

Results

Usability and focus group testing yielded a number of strong, favorable reactions

from providers and families. Recommendations for refining the e-workbook also were provided, and these guided several improvements to the resource prior to initiating the feasibility trial, which is currently underway.

Conclusions

This study aimed to develop and preliminarily evaluate a tablet-based application to improve provider fidelity and child engagement in child mental health treatment. If successful, this approach may serve as a key step toward making best-practice treatment more accessible to children and families. As various technologies continue to increase in popularity worldwide and within the health care field more specifically, it is essential to rigorously test the usability, feasibility, acceptability, and effectiveness of novel health technology solutions. It is also essential to ensure that patients and providers drive decision making that supports the development of these resources to ensure that they can be seamlessly integrated into practice.

Trial Registration

Clinicaltrials.gov NCT01915160; <https://clinicaltrials.gov/ct2/show/NCT01915160>
(Archived by WebCite at <http://www.webcitation.org/6cPIiQDpu>)

Keywords: technology, mobile health, child mental health treatment, feasibility test, fidelity, patient engagement, traumatic stress

Introduction

Background

One in four US children experiences a mental health disorder with severe impairment or distress during their childhood [1-3]. Ensuring that these children have access to the highest quality mental health care is a top public health priority. Efficacious child and adolescent treatments exist for a wide range of mental health disorders [4]. However, these treatments are delivered with variable fidelity in mental health service settings, even among well-trained providers [5-10]. Provider fidelity generally refers to the degree to which a clinician adheres to a treatment protocol and delivers the treatment competently [11]. Drifting or deviating from empirically

supported treatment protocols can diminish an intervention's potency and effectiveness, leading to a major quality shortfall [10,12-14]. Statewide and national dissemination and implementation initiatives are underway to narrow these gaps [15-17]. However, the problem of weak and inconsistent provider fidelity persists and must be addressed to improve the quality of care.

Technology-Based Resources and Access to High Quality Care

Recent technological advances offer an opportunity to support the effective delivery of best-practice interventions. This can be achieved with portable mobile apps that assist providers as they implement treatment activities that are challenging to deliver with high fidelity and child engagement. Technology-based decision support tools have been developed in the broader health care field, and initial data suggest that this approach improves clinical decision making and adherence to best practices and treatment protocols [18-19]. Novel technology-based therapy tools can offer a standardized guiding framework for providers to follow as they progress through a treatment and incorporate several design features to promote provider fidelity to a treatment model, such as (1) inclusion of a diverse yet finite set of in-session and homework activities that are all consistent with the goals of the treatment (ie, rather than off-topic or off-task activities that may encourage drift); (2) presentation of key intervention-related concepts in a consistent manner across providers and clients; and (3) assessment and tracking of progress through a treatment model for each client. Moreover, studies in child education suggest that the integration of interactive games, touch screen learning, video demonstrations, and other engaging features enhance child engagement in learning activities [20-21]. Together, these data increase confidence that technology-based approaches may have value toward improving child engagement and provider fidelity in child mental health treatment. Increased child engagement is particularly important during mental health treatment sessions, because engagement has been shown to reduce risk for dropout, which is another pervasive problem in mental health treatment that limits its impact [22].

Efforts to develop technology-based solutions for mental health care must account for limited resources available in community mental health settings, including the cost and providers' time [23-25]. Contemporary mobile devices such as tablets and mobile phones are low cost and increasingly ubiquitous [26]. Integration of these devices into practice is therefore likely to be feasible from a cost perspective. With regard to time and effort, it is important that novel solutions are user-friendly and

able to be integrated readily into practice with minimal provider training and preparation. This is most likely to be achieved when providers and patients are the key drivers of the development process, working closely with the research team at each phase of design, development, and evaluation. A patient- and provider-centered approach is critical for successful implementation and dissemination.

Research is needed that directs the process of developing novel health care solutions and that measures their potential to improve the quality of mental health care. Research reviews suggest that technology-based tools, broadly considered, effectively enhance mental health care; however, the majority of this research focuses on self-help tools for adults [27-29] or other resources used by patients outside the context of formal treatment sessions [30-32]. Our protocol differs from prior approaches by examining the benefits of mobile device apps used *in session* with an emphasis on interventions designed for children and their caregivers.

Selection of a Treatment Model With Which to Test a Novel Technology-Based Solution

Selecting an appropriate treatment model is an important step in the process of evaluating a new health technology solution. Ideally, the treatment model should have a strong evidence base and high potential for cross-application with other treatment approaches to enhance generalizability of the data. We selected Trauma-Focused Cognitive Behavioral Therapy (TF-CBT) [33-34]; because it is a well-established treatment for children in the mental health field [33-38], has ample evidence supporting its effectiveness [33-38], is widely used, and has been disseminated internationally. Moreover, TF-CBT uniquely addresses multiple symptom domains commonly encountered in mental health treatment settings, including posttraumatic stress, depressed mood, and disruptive behavior. Development of a tablet-based application, or “e-workbook,” for TF-CBT therefore would appear to have high potential to enhance the relevance of our data to a range of established treatments for youth. TF-CBT also requires caregiver involvement, which will allow us to explore the use of in-session resources with adults as well as children. In our pilot work, we conducted structured 30-min interviews with 21 certified national TF-CBT trainers, which revealed significant provider interest in and likely acceptance of tablet-based aids for delivering TF-CBT. These interviews also provided strong direction around key challenges that tablet-based resources can overcome [39].

Purpose and Aims

The purpose of this project was to develop and pilot a tablet-based e-workbook that is designed to increase quality of care in child mental health treatment via improvements in child engagement and provider fidelity. The potential impact of an e-workbook approach, if successful, is extraordinarily high in light of its scalability and sustainability. In the current protocol, we describe a strategy to develop an e-workbook to augment delivery of TF-CBT, a well-established treatment for children and their caregivers. If this approach is found to have utility in practice, it can be applied to a wide range of treatment approaches. The general outline of the protocol addresses three aims: (1) to develop an e-workbook to support delivery of treatment with high fidelity; (2) to conduct usability tests of the e-workbook with families and providers; and (3) to conduct a feasibility trial comparing TF-CBT vs iPad-facilitated TF-CBT.

Methods

Study Design

The aims of this investigation were accomplished in three phases. Phase I included the initial development of the TF-CBT e-workbook. Phase II included focus groups and individual interviews with 21 providers and 24 children (aged 8-16 years) to inform the refinement of the e-workbook (ie, alpha testing). It also included internal beta testing to guide the final editing and debugging process. Finally, Phase III, which is currently underway, features a feasibility trial with 10 providers and 20 families to examine the feasibility of the methodology that we propose to use in a future randomized controlled trial (RCT) as well as the feasibility and acceptability of implementing the TF-CBT e-workbook in community mental health and child welfare agencies.

Phase I: Development of the TF-CBT e-Workbook

Technical Approach to Development The e-workbook was developed as a Web-based, rather than native (ie, device-specific) application. Although this approach requires the user to have Internet access, an advantage is that Web-based tools are

accessible to providers on any of the wide range of web-connected devices. A native application would ultimately require reprogramming and updating for each operating system (Android, iOS, Windows). Although no technology can be completely “future proof,” it was determined that Web-based tools would have the highest potential to remain useful in meeting the aims of this project as technology progresses. Native versions of the app will likely represent the final step toward disseminating this resource once it is fully evaluated and refined.

The application is responsive; that is, it detects the type of device that is in use and adjusts to it for optimal look and feel. For example, the application can detect an iPad and use JavaScript to convert desktop mouse actions to accelerometer actions. This functionality enables the use of features such as shaking the device to trigger a response, inertial finger drawing, and the built-in microphone. Conversely, those actions would naturally degrade to normal mouse actions if a user were to access the resource via desktop computer. As the application detects different devices, the responsive layout naturally adapts to the devices’ native display parameters. This achieves a fully integrated and device-agnostic application and increases potential for adoption across practice settings.

Content Development for the TF-CBT e-Workbook Qualitative data, collected regularly as part of consultation calls conducted in TF-CBT training programs, provided valuable direction in the technical and content development phases. Specifically, these data were used to determine which components of TF-CBT are most vulnerable to drift and what activities could be developed to overcome challenges to child engagement and provider fidelity in each TF-CBT component. For example, providers who were trained in TF-CBT reported that they used the psychoeducation, anxiety management, and coping components of TF-CBT at least 50% of the time with their child trauma cases; but that the exposure components were reported least frequently (26-50%) [40]. Therefore, greater attention was paid to emphasizing exposure-specific elements of TF-CBT rather than general psychotherapy skills. The research team also carefully reviewed data collected from expert clinicians [39] to identify areas in which technology-based resources would have high potential to enhance (1) provider fidelity; (2) child engagement in treatment activities; (3) child or caregiver understanding of key treatment concepts; (4) likelihood of skill acquisition; or (5) patient adherence to homework exercises. Taken together, these data and observations were used to create the content and format of the e-workbook activities and resources listed in [Table 1](#).

We developed numerous resources ([Table 1](#)) or “chapters” for use by providers during individual sessions with each child. Introduction videos were created for children and are available on the first screen of each chapter. Each video depicts a teenager who explains the rationale for the chapter and presents brief examples that illustrate completion of the activity. Our decision to feature older youth (ages 15-16) in most of the videos was based on focus group feedback from children aged 8-15 years. Specifically, youth at the younger end of the age range stated that they would be equally pleased with younger or older actors in the videos, whereas youth at the older end of the age range agreed unanimously that actors should be older adolescents, and that they would likely experience very little connection to a younger actor. Some videos are brief (ie, 30-60 second) clips designed for a provider to use after introducing a concept or teaching a skill with the goal of enhancing engagement and reinforcing what the provider taught during the session (eg, videos demonstrating the CBT triangle in the cognitive coping component of TF-CBT). Some chapters feature interactive touch screen games, such as drag-and-drop activities, drawing tools, trivia-style card games, and animated relaxation activities. Each activity was developed to address an element of the TF-CBT protocol that was identified by experts and providers as challenging to implement with high fidelity and engagement.

Some resources included activities for providers to complete with caregivers. These consisted of an extensive collection of video clips with narration that demonstrate a wide range of effective behavior management skills, including common mistakes and how to correct them. Videos were not intended to replace provider instruction and demonstration, but to support providers’ attempts to teach caregivers how to apply skills across a broad range of situations and settings. Additional resources were developed for the provider. These consisted primarily of content that assists with session preparation tasks, such as setting session goals and potential agenda items, tracking, updating assigned homework activities, and guiding their clinical decision making as families progress in TF-CBT. We also created provider tabs for most resources that provide discussion points, tips, and ideas for supplemental exercises.

Phase II: Alpha and Beta Testing

Procedure The primary purpose of alpha testing in software and intervention resource development is to assess reactions and obtain direct input from end users

regarding design, content, and functionality. These data are needed to guide improvements to the resource. Participants were provided with tablets (ie, iPads) in either individual interviews or focus group settings to interact with a select set of resources within the TF-CBT e-workbook. Providers were given access to all activities listed in [Table 1](#), and children were given access to a representative sample of activities. Participants were asked a series of semistructured questions administered by trained interviewers. The questions asked participants to provide reactions to each tool, ranging from reactions to the overall “look and feel” of the interface, ease of navigation, and technical problems they experience. An average of 5 minutes was dedicated to review and discuss each resource. Qualitative data collected during these sessions were audiotaped for transcription, coding, and data analysis.

Participants Alpha testing was completed via focus groups with providers (n=22) and via focus groups or individual interviews with children (n=24). Focus group sizes were generally between 5-8 participants.

Children aged 8-11 years and 12-16 years participated in separate groups; average age was 13.0 years. Children were recruited from clinical sites (local child advocacy and mental health treatment centers) and local schools with on-site mental health services with high rates of trauma exposure. Inclusion/exclusion criteria for clinic-referred children were consistent with those used in prior RCTs with TF-CBT: children were 8-16 years old, had a history of at least one potentially traumatic event, had current clinically elevated symptoms of PTSD, and did not have active suicidal or homicidal ideations or symptoms of psychosis. Participants recruited from the schools were also 8-16 years old but not required to have a reported trauma history or a particular symptom profile. This combined clinical/nonclinical sample was used to maximize input on the look-and-feel of the TF-CBT e-workbook in addition to applicability and clarity of content to youth in the targeted age range. Sample of children was 65% female, 86% Black/African American, 14% white.

Providers were recruited from several child advocacy and mental health centers. Providers were eligible if they were fully trained in TF-CBT and carried active child trauma cases at the time of the study. There was diversity in the provider sample with respect to discipline (64% counseling, 18% social work, 18% psychology), credentials (18% doctorate, 23% MSW/LCSW/LISW/LMSW, 36% LPC, 23% other Master’s degree), and years of experience delivering child trauma treatment (46%

5<years, 27% 3-5 years, 27% 1-3 years). Providers also varied in the settings in which they worked (45% mental health clinic, 55% child advocacy center), with several providers delivering TF-CBT in schools or patients' homes through outreach clinics. The provider sample was 86% female, 73% Caucasian, 18% Black/African American, 4.5% Native American, and 4.5% other (unspecified).

Data Analysis Focus group and interview responses were audiorecorded and transcribed. First, a content analysis of the interview responses was conducted through multiple close readings of the transcriptions by two independent coders (authors on this paper). Each coder generated an independent list of thematic categories and subcategories based on their review of the data (eg, usefulness of a specific tool; age appropriateness). These themes were then further developed and ordered by the primary coder and reviewed and edited by the second coder. The coders then met in a consensus conference to discuss the categories, resolve questions, and refine the thematic categories. Once this was accomplished, the themes were again reviewed. After additional discussion to review and refine categories and resolve questions, the final thematic categories were completed and higher-order categories were developed. We have previously used similar analytic approaches to qualitative research with a range of public sector health care patient and provider populations. [39,41-43] Results of these analyses, reported in [Table 2](#) (providers) and [Table 3](#) (children), were used to refine and fine tune the TF-CBT e-workbook. The vast majority of recommendations given by children and providers were addressed in the context of the current grant prior to feasibility testing. Some recommendations were cost prohibitive, but can be addressed with future funding in preparation for a large-scale RCT.

Beta testing followed the alpha testing phase and is a form of usability testing in which refined or revised resources are evaluated by a new set of providers for acceptability and functionality. This was to ensure that any errors existing within the TF-CBT e-workbook were identified and corrected in preparation for the feasibility trial. Members of the development and investigative team reviewed all the revised components of the resource, and additional edits and refinements to content, appearance, and functionality were made in response. [Figures 1-7](#) include screenshots of several e-workbook activities.

Phase III: Feasibility Trial

Approach The goal of the ongoing feasibility trial is to examine the feasibility of implementing the TF-CBT e-workbook in community mental health service agencies, and to demonstrate the randomized controlled trial (RCT) methodology we propose to use in a future large-scale trial to test the effectiveness of the TF-CBT e-workbook in clinics across the state. Resources and data resulting from this project are designed to position us well for a large-scale RCT to examine the incremental benefit of e-workbooks to best-practice interventions compared to standard care without technology. The strengths and limitations associated with conducting a small-scale feasibility RCT versus an open trial were carefully considered. A major benefit of the latter approach included increased quantity of feasibility data related to the use of the e-workbook. On balance, this option was outweighed by the importance of demonstrating the feasibility of the full RCT methodology. This would allow us to identify and address barriers in recruitment, retention, fidelity assessment, and other procedures essential to successful conduct of a RCT. This approach is consistent with expert recommendations to pilot test the feasibility of methods to be used in large RCTs, use data yielded by such studies to “debug” the methodology, and assess optimal strategies to executing the RCT [44].

Design The feasibility trial will involve 10 providers recruited from four participating community-based mental health and child welfare agencies where TF-CBT is delivered. Providers will have been fully trained in TF-CBT and carry active child trauma cases at the onset of the trial. Providers will be block randomized by site to either the e-workbook facilitated TF-CBT (n=5) or standard TF-CBT (TAU) (n=5) conditions. Each provider will treat at least two children for a total sample of 20 children. We will also pursue an exploratory aim regarding the feasibility of using the e-workbook across multiple treatment delivery settings by including a small subsample of mental health providers who are primarily based in school settings (at least two per study condition).

An independent, trained evaluator who is blind to the study condition will conduct baseline, mid-treatment, and post-treatment clinical assessments. Families will be reimbursed \$30 for each administration of the assessment battery. All sessions will be audiorecorded and coded for fidelity and child engagement by independent coders.

Alternatives to this design decision were considered carefully. For example, the design could have included each provider treating one e-workbook case and one

standard TF-CBT case, with randomized ordering of conditions by provider. An advantage of this approach is that provider factors are less likely to have strong impact on clinical outcomes in this small-scale trial. A weakness of the approach is that use of the e-workbook could affect performance on subsequent standard TF-CBT cases for providers assigned to deliver TF-CBT using the e-workbook first.

Additionally, it was determined that if providers within an agency were assigned to different starting conditions, this could increase risk of contamination across conditions with standard TF-CBT providers hearing about, viewing, and potentially even using elements of the e-workbook with standard TF-CBT cases. Overall, it was concluded that block randomization by clinic to the two study conditions was a preferable design strategy.

Participants

Providers A minimum of 10 providers who have been trained in TF-CBT and maintain active child trauma caseloads will be recruited from four participating clinics in the Charleston tri-county area. The clinics include both outpatient mental health clinics and child advocacy centers. We will make every effort to recruit a racially/ethnically diverse sample of providers to allow integration of feedback from providers with diverse backgrounds. We will also attempt to recruit a balanced sample with respect to sex and gender. Inclusion criteria for providers will be as follows. Providers will be full- or part-time employees of the participating clinic and had to have obtained at least a Master's degree in social work, counseling, clinical psychology, or a related field. Each provider will provide treatment to at least two study cases during the feasibility trial. Providers will be encouraged to enroll up to 5 children to increase the likelihood that at least 2 cases will be completed per provider during the trial.

Patients A minimum of 20 children aged 5-15 years and their caregivers will be recruited into the feasibility trial. These children will be recruited from participating clinics in the local area. We will make every effort to recruit a racially/ethnically diverse sample of youth (eg, >25% African American) to ensure integration of feedback from patients with diverse backgrounds. Inclusion criteria will be as follows. Participating children will be aged 5-15 years, have experienced at least one potentially traumatic event (eg, sexual assault, physical assault, witnessed violence, disaster, serious accident), and have at least one symptom on each PTSD symptom cluster (re-experiencing, avoidance, hyperarousal) based on a diagnostic interview.

Cases will be excluded when the child or caregiver exhibits psychotic symptoms (eg, active hallucinations, delusions, impaired thought processes); significant cognitive disability, developmental delays, or pervasive developmental disorder; or active suicidal or homicidal ideations. Children also will be excluded when there is no consistent caregiver available to participate (eg, short-term foster care placement, restrictions by child protective services). These criteria are consistent with those used in prior TF-CBT clinical trials.

Assessment A trained evaluator both blinded to treatment condition and fully trained in the administration of all measures will administer the assessment battery. All measures are well-validated and widely used instruments in the mental health field and in the treatment outcome research. Assessment will occur on three occasions: at baseline (ie, during the screening interview), at mid-treatment (ie, after the sixth treatment session, approximately two months post-baseline), and at post-treatment (ie, after the twelfth treatment session, approximately four months post-baseline). Treatment fidelity and child engagement will be measured via observational coding of audiotapes.

Treatment Fidelity Fidelity to the TF-CBT and e-workbook-facilitated TF-CBT protocols will be measured via coding of audiotaped treatment sessions by independent, trained raters. Treatment sessions will be audiotaped for both study conditions (n=20 x ~10 sessions average=~200 audiotapes). Ratings will be completed using a behaviorally specific coding system of TF-CBT provider behavior that was modified for the current study to ensure relevance to the e-workbook condition [45]. The coding system will be used to calculate providers' fidelity to each TF-CBT component. An additional eight items focus on general therapy skills, not specific to TF-CBT, including establishing an agenda, providing a treatment rationale, and assigning homework. Additional items were created to identify use of the e-workbook activities to differentiate the two study conditions. Two independent raters will listen to audiorecorded treatment session tapes and complete the modified fidelity measure to code the use and extensiveness of specific treatment techniques depicted on the recordings. Raters will be trained in the coding system and meet biweekly throughout the remainder of the feasibility trial to ensure maintenance of acceptable levels of accuracy and interrater reliability. Discrepant ratings will be reviewed until consensus is achieved. If the two raters cannot reach consensus, the PI and Co-Is will make final decisions.

Child Engagement Child engagement will be measured via coding of audiotaped sessions by independent, trained raters. The Child Involvement Ratings Scale (CIRS) [46-47], a 6-item scale that measures child engagement for each session, will be used for this purpose. Four “positive” involvement items and two “negative” involvement items are rated for each session on a 6-point scale (ie, “not at all” to “a great deal present”). The positive-involvement items emphasize the extent to which children initiate discussions, demonstrate enthusiasm, self-disclose, and demonstrate understanding. Negative-involvement items address withdrawal or avoidance in treatment. Coders will provide ratings based on two 10-min segments of session audiotapes (ie, beginning at min 10 and min 40). Child engagement ratings on the CIRS have been associated with clinical outcomes [47] and provider flexibility in delivery of EBTs [48]. Excellent internal consistency and strong interrater reliability have been reported for this measure [47-48].

Child-Report of Functioning

Clinical Interview The Kiddie Schedule for Affective Disorders and Schizophrenia for School Age Children-Present and Lifetime version (K-SADS-PL PTSD module) [49] is a semistructured interview that will be used to assess PTSD symptom levels and diagnostic status. The K-SADS-PL is well-established and used widely. It also has been used in numerous TF-CBT RCTs [33]. The K-SADS-PL also assesses functional impairment in school, social, and family life.

Self-Report Instruments The Center for Epidemiological Studies Depression Scale for Children (CES-DC) assesses the severity of depressive symptomatology in children. The CES-DC is a 20-item self-report depression inventory with possible scores ranging from 0-60. Scores over 15 are indicative of significant levels of depressive symptoms [50,51]. The UCLA PTSD Index for Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV), Child Version (UCLA-PTSD) [52-54] will be used to assess the severity of PTSD symptomatology in children. The UCLA PTSD assesses exposure to traumatic events and all 17 DSM-IV symptoms of PTSD. Psychometric research has yielded significant support for the reliability, construct validity, and PTSD criterion-related validity [54]. The Therapeutic Alliance Scale for Children (TASC) [55-57] will be used to measure levels of therapeutic alliance. The TASC is an 8-item measure of the child’s alliance with the provider using a 4-point scale. It has good internal consistency and interrater reliability [57]. The Child/Adolescent Satisfaction Questionnaire (CASQ) [58] is a 15-item instrument that

assesses child satisfaction with mental health treatment.

Caregiver Report of Functioning

Clinical Interview The K-SADS-PL [49] also will be administered to caregivers to assess children's PTSD symptoms and functional impairment.

Caregiver Report Instruments The UCLA PTSD Index for DSM-IV, Parent Version [59] will be used to assess the severity of the child's PTSD symptomatology from the perspective of the caregiver. The Child Behavior Checklist-Parent Report (CBCL) [60] is a widely used measure of behavioral and social maladjustment in children, as perceived by the caregiver. The CBCL has strong psychometric properties. The Beck Depression Inventory (BDI) [61-62] will be used to assess severity of caregivers' depressive symptomatology. The BDI is a 21-item self-report scale of depression and is widely researched and has excellent concurrent and discriminant validity [63]. The Caregiver Satisfaction Questionnaire (CSQ) [58] is a 15-item instrument that assesses caregiver satisfaction with mental health treatment. The Working Alliance Inventory (WAI-short form) [64] is a 12-item measure of the parent-therapist alliance using a 7-point scale (ie, "never" to "always").

User Reactions to TF-CBT e-Workbook To obtain direct input on the e-workbook's design, content, and functionality, post-treatment assessments will include semistructured interviews with children, caregivers, and providers after completion of e-workbook-facilitated TF-CBT. Patients and providers will complete semistructured interviews about their experiences with the e-workbook and how it affected treatment. Interviews will be audiotaped, transcribed, and interpreted using the same approach used during alpha testing.

Data Analysis Plan For both conditions, several patient- and provider-level variables, as well as data collection procedures will be assessed and described. Patient-level variables will include the percentage of eligible patients recruited, treatment attrition, study retention, and session attendance. Recruitment will be assessed by the proportion of patients who agree to participate as compared to the total number solicited to enroll. Attrition will be assessed by examining the proportion of patients who prematurely terminate treatment. Qualitative analyses will be conducted to identify themes in termination. Study retention refers to the proportion of patients who complete all assessment points associated with the treatment protocol,

including those who terminate treatment prematurely.

Provider-level variables will include provider recruitment to participate, fidelity to TF-CBT procedures, use of the resources within the e-workbook, and adherence to the session audiotaping protocol. Recruitment will be assessed by the proportion of providers who agreed to participate as compared to the total number approached. Provider fidelity will consist of the proportion of completed treatment components for the intervention according to the fidelity measure described previously. Providers will be interviewed at the end of treatment with e-workbook cases to provide feedback on the usability of the e-workbook activities. Other data to be summarized include kappa coefficients between independent fidelity raters, scheduling and logistical barriers to completing assessment, and communication successes and failures between study staff and clinic site staff around recruitment efforts to demonstrate feasibility of study procedures for our planned RCT.

Results

At the time of manuscript submission, Phases I and II of the study (mobile application development and refinement) were completed, and initial recruitment for Phase III (feasibility trial) is underway. No data have been cleaned or analyzed for the feasibility trial component of the project. All aspects of this federally funded study have been approved by the institutional review board (IRB) at the institution where the research is being conducted. Usability and focus group testing yielded a number of strong, favorable reactions from providers and families. Recommendations for refining the e-workbook also were provided, and these guided several improvements to the resource prior to initiating the feasibility trial, which is currently underway.

Discussion

The current protocol advances methods for developing technologies for use in mental health. First, it moves the field forward by developing and evaluating technology-based tools designed specifically to support treatment delivery and quality of care by targeting provider fidelity and, second, through engaging children. This represents a key step toward making EBTs more accessible to children and families. Existing technology-based tools largely target the (adult) patient directly by assisting them in self-care or homework adherence [25-43,63]. Such resources are not designed to support providers in the delivery of interventions with fidelity, and therefore are

unlikely to have significant direct effect on the quality of care that families receive when they present for treatment in community mental health service agencies. Second, this line of research will provide valuable data about the potential for technology-based resources to enhance children's engagement in treatment. This is an important, but significantly underdeveloped, area of research that may have critical implications for the health care field. Third, we capitalized on recent technological advances by developing Web-based applications that are optimized for mobile devices. This allows us to test the resources on a tablet (eg, iPad) while ensuring that we achieve a fully integrated and device-agnostic application that will operate with the most current technologies without the need to rewrite the application on a device-by-device basis. Fourth, we developed a wide range of tools (eg, videos, interactive games, drawing applications) that providers will use with children and adults (caregivers). This will ensure collection of valuable feasibility data that have high relevance to several design formats and target populations. Alternatively, a narrower focus on a specific patient population (eg, adults with depression) or specific type of resource (eg, videos only) would have had relatively less potential to significantly advance the field.

This study represents a unique opportunity to capitalize on the increasing use of mobile phone, Internet, and tablets by both patients and providers to support providers' efforts to deliver empirically supported treatments with a high level of fidelity. As these technologies continue to increase in popularity worldwide and within the health care field more specifically, it is essential to rigorously test the usability, feasibility, acceptability, and effectiveness of these resources. It is also essential to use input from patients and providers to drive decision-making related to development of these resources to ensure that they can be seamlessly integrated into practice. This study takes an initial step toward evaluating the feasibility and utility of implementing best-practice treatment with the assistance of a patient- and provider-centered tablet-based e-workbook resource. If feasibility is supported by our pilot trial, we will propose a rigorous, statewide efficacy evaluation powered to examine the impact on child mental health outcomes. Data yielded from such an evaluation will have tremendous value for purposes of developing and disseminating highly accessible resources that are designed to enhance the quality of treatment delivered to children and families in a wide range of service settings.

Acknowledgments

We thank Judy Cohen, MD, Esther Deblinger, PhD, and Anthony Mannarino, PhD, who developed TF-CBT and have provided valuable input and feedback throughout every phase of this project. We thank Ben Jacobs of Fuzzco for his significant role in the design and development process of the TF-CBT e-workbook. We thank Josh Nissenboim of Fuzzco for his leadership and partnership in support of this project and related research. We thank Joel Sherrill of NIMH for his thoughtful feedback on early conceptualizations of this project. We also thank the families and providers who participated in this study for their time and insights. This research and the authors' time and effort were supported by National Institute of Mental Health grants R34-MH096907, R01-MH81056, R34-MH104470, and T32-MH18869; National Institute on Drug Abuse grants K12-DA031794, K23-DA038257, and R01-DA031285; and Substance Abuse and Mental Health Services Administration grant U79-SM061269. None of these funding agencies were involved in the design of the study; the collection, analysis, or interpretation of data; the writing of the manuscript, or the decision to submit the manuscript for publication.

Abbreviations

BDI	Beck Depression Inventory
CASQ	Child/Adolescent Satisfaction Questionnaire
CBLC	Child Behavior Checklist
CBT	cognitive behavioral therapy
CES-DC	Center for Epidemiological Studies Depression Scale for Children
CIRS	Child Involvement Ratings Scale
CSQ	Caregiver Satisfaction Questionnaire
DSM-IV	Diagnostic and Statistical Manual of Mental Disorders, Fourth

Edition

EBT	evidence-based treatment
e-workbook	tablet-facilitated Trauma-Focused CBT (mobile application)
IRB	institutional review board
KSADS-PL	Kiddie Schedule for Affective Disorders and Schizophrenia for School Age Children-Present and Lifetime version
PTSD	posttraumatic stress disorder
RCT	randomized controlled trial
TAC-R	Treatment Adherence Checklist-Revised
TASC	Therapeutic Alliance Scale for Children
TAU	treatment-as-usual
TF-CBT	Trauma-Focused Cognitive Behavioral Therapy
UCLA-PTSD	UCLA PTSD Index for DSM-IV
WAI	Working Alliance Inventory

Footnotes

Conflicts of Interest: None declared.

References

1. Costello EJ, Egger H, Angold A. 10-year research update review: the epidemiology of child and adolescent psychiatric disorders: I. Methods and public health burden. *J Am Acad Child Adolesc Psychiatry*. 2005 Oct;44(10):972–86. doi: 10.1097/01.chi.0000172552.41596.6f.S0890-8567(09)61759-7 [PubMed: 16175102] [CrossRef: 10.1097/01.chi.0000172552.41596.6f]
2. Kessler RC, Avenevoli S, Costello EJ, Georgiades K, Green JG, Gruber MJ, He J, Koretz D, McLaughlin KA, Petukhova M, Sampson NA, Zaslavsky AM, Merikangas KR. Prevalence, persistence, and sociodemographic correlates of DSM-IV disorders in the National Comorbidity Survey Replication Adolescent Supplement. *Arch Gen Psychiatry*. 2012 Apr;69(4):372–80. doi: 10.1001/archgenpsychiatry.2011.160. <http://europepmc.org/abstract/MED/22147808> .archgenpsychiatry.2011.160 [PMCID: PMC3445020] [PubMed: 22147808] [CrossRef: 10.1001/archgenpsychiatry.2011.160]
3. Merikangas KR, He J, Burstein M, Swanson SA, Avenevoli S, Cui L, Benjet C, Georgiades K, Swendsen J. Lifetime prevalence of mental disorders in U.S. adolescents: results from the National Comorbidity Survey Replication--Adolescent Supplement (NCS-A) *J Am Acad Child Adolesc Psychiatry*. 2010 Oct;49(10):980–9. doi: 10.1016/j.jaac.2010.05.017. <http://europepmc.org/abstract/MED/20855043> .S0890-8567(10)00476-4 [PMCID: PMC2946114] [PubMed: 20855043] [CrossRef: 10.1016/j.jaac.2010.05.017]
4. Chorpita B, Daleiden E, Ebesutani C. Evidence-based treatments for children and adolescents: An updated review of indicators of efficacy and effectiveness. *Clin Psychol Sci Prac*. 2011;18:154–72. [http://www.childfirst.ucla.edu/Chorpita%20et%20al%20\(2011\)-Evidence-Based%20Treatment%20for%20Children.pdf](http://www.childfirst.ucla.edu/Chorpita%20et%20al%20(2011)-Evidence-Based%20Treatment%20for%20Children.pdf) .
5. Drake RE, Torrey WC, McHugo GJ. Strategies for implementing evidence-based practices in routine mental health settings. *Evid Based Ment Health*. 2003 Feb;6(1):6–7. <http://ebmh.bmj.com/cgi/pmidlookup?view=long&pmid=12588814> . [PubMed: 12588814]
6. Garland AF, Brookman-Frazee L, Hurlburt MS, Accurso EC, Zoffness RJ, Haine-Schlagel R, Ganger W. Mental health care for children with disruptive behavior problems: a view inside therapists' offices. *Psychiatr Serv*. 2010 Aug;61(8):788–95. doi: 10.1176/appi.ps.61.8.788. <http://europepmc.org/abstract/MED/20675837> .61/8/788 [PMCID: PMC3019612] [PubMed: 20675837] [CrossRef: 10.1176/appi.ps.61.8.788]
7. Kazak AE, Hoagwood K, Weisz JR, Hood K, Kratochwill TR, Vargas LA, Banez GA. A meta-systems approach to evidence-based practice for children and adolescents. *Am Psychol*. 2010;65(2):85–97. doi: 10.1037/a0017784.2010-02208-011 [PubMed: 20141264] [CrossRef: 10.1037/a0017784]

8. McHugo GJ, Drake RE, Whitley R, Bond GR, Campbell K, Rapp CA, Goldman HH, Lutz WJ, Finnerty MT. Fidelity outcomes in the National Implementing Evidence-Based Practices Project. *Psychiatr Serv*. 2007 Oct;58(10):1279–84. doi: 10.1176/appi.ps.58.10.1279.58/10/1279 [PubMed: 17914003] [CrossRef: 10.1176/appi.ps.58.10.1279]
9. Raghavan R, Inoue M, Ettner SL, Hamilton BH, Landsverk J. A preliminary analysis of the receipt of mental health services consistent with national standards among children in the child welfare system. *Am J Public Health*. 2010 Apr;100(4):742–9. doi: 10.2105/AJPH.2008.151472.AJPH.2008.151472 [PMCID: PMC2836330] [PubMed: 19608950] [CrossRef: 10.2105/AJPH.2008.151472]
10. McCabe O. Crossing the Quality Chasm in Behavioral Health Care: The Role of Evidence-Based Practice. *Professional Psychology: Research and Practice*. 2004;35(6):571–579. doi: 10.1037/0735-7028.35.6.571. [CrossRef: 10.1037/0735-7028.35.6.571]
11. Cross WF, West JC. Examining implementer fidelity: Conceptualizing and measuring adherence and competence. *J Child Serv*. 2011;6(1):18–33. doi: 10.5042/jcs.2011.0123. <http://europepmc.org/abstract/MED/21922026> . [PMCID: PMC3171488] [PubMed: 21922026] [CrossRef: 10.5042/jcs.2011.0123]
12. Bond G, Becker D, Drake R. Measurement of fidelity of implementation of evidence based-practices: Case examples of the IPS fidelity scale. *Clin Psychol Sci Prac*. 2011;18:126–41. doi: 10.1111/j.1468-2850.2011.01244.x. [CrossRef: 10.1111/j.1468-2850.2011.01244.x]
13. McHugo GJ, Drake RE, Teague GB, Xie H. Fidelity to assertive community treatment and client outcomes in the New Hampshire dual disorders study. *Psychiatr Serv*. 1999 Jun;50(6):818–24. [PubMed: 10375153]
14. Schoenwald SK, Sheidow AJ, Letourneau EJ. Toward effective quality assurance in evidence-based practice: links between expert consultation, therapist fidelity, and child outcomes. *J Clin Child Adolesc Psychol*. 2004 Mar;33(1):94–104. doi: 10.1207/S15374424JCCP3301_10. [PubMed: 15028545] [CrossRef: 10.1207/S15374424JCCP3301_10]
15. Karlin BE, Ruzek JI, Chard KM, Eftekhari A, Monson CM, Hembree EA, Resick PA, Foa EB. Dissemination of evidence-based psychological treatments for posttraumatic stress disorder in the Veterans Health Administration. *J Trauma Stress*. 2010 Dec;23(6):663–73. doi: 10.1002/jts.20588. [PubMed: 21171126] [CrossRef: 10.1002/jts.20588]
16. Saunders B. Bringing in evidence-based practice: How do we make EBP TAU?. Leadership Symposium on Evidence-Based Practice in Human Services; January 30, 2009; San Diego, CA. 2009.
17. Markiewicz J, Ebert L, Ling D, Amaya-Jackson L, Kisiel C. *Learning Collaborative Toolkit. Raising*

the standard of care for traumatized children and their families. Los Angeles, CA and Durham, NC: National Center for Child Traumatic Stress; 2006. [2015-07-23].

http://www.nctsn.org/nctsn_assets/pdfs/lc/Module_all.pdf .

18. Kubben PL, van SH, Cornips E M J, Vaccaro AR, Dvorak MF, van Rhijn L W, Scherpbier A J J A, Hoogland H. An evidence-based mobile decision support system for subaxial cervical spine injury treatment. *Surg Neurol Int*. 2011;2:32. doi: 10.4103/2152-7806.78238.

<http://www.surgicalneurologyint.com/article.asp?issn=2152-7806;year=2011;volume=2;issue=1;spage=32;epage=32;aulast=Kubben> . [PMCID: PMC3086168] [PubMed: 21541200] [CrossRef: 10.4103/2152-7806.78238]

19. Roy P, Durieux P, Gillaizeau F, Legall C, Armand-Perroux A, Martino L, Hachelaf M, Dubart A, Schmidt J, Cristiano M, Chretien J, Perrier A, Meyer G. A computerized handheld decision-support system to improve pulmonary embolism diagnosis: a randomized trial. *Ann Intern Med*. 2009 Nov 17;151(10):677–86. doi: 10.7326/0003-4819-151-10-200911170-00003.151/10/677 [PubMed: 19920268] [CrossRef: 10.7326/0003-4819-151-10-200911170-00003]

20. Kang B, Tan S. Impact of digital games on intrinsic/extrinsic motivation, achievement, satisfaction. McFerrin K. eds. Proceedings of Society for Information Technology & Teacher Education International Conference; Society for Information Technology & Teacher Education International Conference; 03/03/2008; Las Vegas, NV. Chesapeake, VA: Association for the Advancement of Computing in Education; 2008. Mar 03, pp. 1825–32.

21. Papastergiou M. Digital Game-Based Learning in high school Computer Science education: Impact on educational effectiveness and student motivation. *Computers & Education*. 2009 Jan;52(1):1–12. doi: 10.1016/j.compedu.2008.06.004. [CrossRef: 10.1016/j.compedu.2008.06.004]

22. McKay M, Hibbert R, Hoagwood K. Integrating evidence based engagement interventions into "real world" child mental health settings. *Brief Treatment and Crisis Intervention*. 2004;4:177–86. doi: 10.1093/brief-treatment/mhh014.

<http://familyinvolvementcenter.org/phocadownloadpap/Integrating%20Evidence-Based%20Engagement%20Interventions.pdf> . [CrossRef: 10.1093/brief-treatment/mhh014]

23. Kazdin AE, Blase SL. Rebooting Psychotherapy Research and Practice to Reduce the Burden of Mental Illness. *Perspect Psychol Sci*. 2011 Jan;6(1):21–37. doi: 10.1177/1745691610393527.6/1/21 [PubMed: 26162113] [CrossRef: 10.1177/1745691610393527]

24. Luxton D, McCann R, Bush N, Mishkind M, Reger G. mHealth for mental health: Integrating smartphone technology in behavioral healthcare. *Prof Psychol Res Pr*. 2011;42:505. doi: 10.1037/a0024485. [CrossRef: 10.1037/a0024485]

25. Amstadter AB, Broman-Fulks J, Zinzow H, Ruggiero KJ, Cercone J. Internet-based interventions for traumatic stress-related mental health problems: a review and suggestion for future research. *Clin Psychol Rev*. 2009 Jul;29(5):410–20. doi: 10.1016/j.cpr.2009.04.001. [http://europepmc.org/abstract/MED/19403215.S0272-7358\(09\)00046-4](http://europepmc.org/abstract/MED/19403215.S0272-7358(09)00046-4) [PMCID: PMC2704915] [PubMed: 19403215] [CrossRef: 10.1016/j.cpr.2009.04.001]
26. Smith A. *Pew Research Center*. 2015. Apr 01, [2015-10-27]. U.S. Smartphone Use in 2015 http://www.pewinternet.org/files/2015/03/PI_Smartphones_0401151.pdf.
27. Carlbring P, Andersson Internet and psychological treatment: How well can they be combined? *Comput Hum Behav*. 2006;22:545–53. doi: 10.1016/j.chb.2004.10.009. [CrossRef: 10.1016/j.chb.2004.10.009]
28. Cuijpers P, Marks IM, van SA, Cavanagh K, Gega L, Andersson G. Computer-aided psychotherapy for anxiety disorders: a meta-analytic review. *Cogn Behav Ther*. 2009;38(2):66–82. doi: 10.1080/16506070802694776.912391320 [PubMed: 20183688] [CrossRef: 10.1080/16506070802694776]
29. Clough BA, Casey LM. Technological adjuncts to enhance current psychotherapy practices: a review. *Clin Psychol Rev*. 2011 Apr;31(3):279–92. doi: 10.1016/j.cpr.2010.12.008.S0272-7358(11)00003-1 [PubMed: 21382535] [CrossRef: 10.1016/j.cpr.2010.12.008]
30. Luxton DD, June JD, Kinn JT. Technology-based suicide prevention: current applications and future directions. *Telemed J E Health*. 2011;17(1):50–4. doi: 10.1089/tmj.2010.0091. [PubMed: 21214382] [CrossRef: 10.1089/tmj.2010.0091]
31. Morris ME, Kathawala Q, Leen TK, Gorenstein EE, Guilak F, Labhard M, Deleeuw W. Mobile therapy: case study evaluations of a cell phone application for emotional self-awareness. *J Med Internet Res*. 2010;12(2):e10. doi: 10.2196/jmir.1371. <http://www.jmir.org/2010/2/e10/> v12i2e10 [PMCID: PMC2885784] [PubMed: 20439251] [CrossRef: 10.2196/jmir.1371]
32. Sandler E. *Psychol Today*. 2010. [2015-10-27]. New mobile apps provide 24-7 check-ins <https://www.psychologytoday.com/blog/promoting-hope-preventing-suicide/201006/new-mobile-apps-provide-24-7-check-ins>.
33. Deblinger E, Mannarino AP, Cohen JA, Steer RA. A follow-up study of a multisite, randomized, controlled trial for children with sexual abuse-related PTSD symptoms. *J Am Acad Child Adolesc Psychiatry*. 2006 Dec;45(12):1474–84. doi: 10.1097/01.chi.0000240839.56114.bb.S0890-8567(09)61805-0 [PubMed: 17135993] [CrossRef: 10.1097/01.chi.0000240839.56114.bb]
34. Cohen JA, Deblinger E, Mannarino AP, Steer RA. A multisite, randomized controlled trial for children with sexual abuse-related PTSD symptoms. *J Am Acad Child Adolesc Psychiatry*. 2004

Apr;43(4):393–402. doi: 10.1097/00004583-200404000-00005.

<http://europepmc.org/abstract/MED/15187799> .S0890-8567(09)61245-4 [PMCID: PMC1201422] [PubMed: 15187799] [CrossRef: 10.1097/00004583-200404000-00005]

35. Cohen JA, Mannarino AP, Iyengar S. Community treatment of posttraumatic stress disorder for children exposed to intimate partner violence: a randomized controlled trial. *Arch Pediatr Adolesc Med*. 2011 Jan;165(1):16–21. doi: 10.1001/archpediatrics.2010.247.165/1/16 [PubMed: 21199975] [CrossRef: 10.1001/archpediatrics.2010.247]

36. Deblinger E, Mannarino AP, Cohen JA, Runyon MK, Steer RA. Trauma-focused cognitive behavioral therapy for children: impact of the trauma narrative and treatment length. *Depress Anxiety*. 2011 Jan;28(1):67–75. doi: 10.1002/da.20744. [PMCID: PMC6675414] [PubMed: 20830695] [CrossRef: 10.1002/da.20744]

37. Silverman WK, Ortiz CD, Viswesvaran C, Burns BJ, Kolko DJ, Putnam FW, Amaya-Jackson L. Evidence-based psychosocial treatments for children and adolescents exposed to traumatic events. *J Clin Child Adolesc Psychol*. 2008 Jan;37(1):156–83. doi: 10.1080/15374410701818293.792182672 [PubMed: 18444057] [CrossRef: 10.1080/15374410701818293]

38. Wethington HR, Hahn RA, Fuqua-Whitley DS, Sipe TA, Crosby AE, Johnson RL, Liberman AM, Mościcki E, Price LN, Tuma FK, Kalra G, Chattopadhyay SK, Task Force on Community Preventive Services The effectiveness of interventions to reduce psychological harm from traumatic events among children and adolescents: a systematic review. *Am J Prev Med*. 2008 Sep;35(3):287–313. doi: 10.1016/j.amepre.2008.06.024.S0749-3797(08)00528-X [PubMed: 18692745] [CrossRef: 10.1016/j.amepre.2008.06.024]

39. Hanson RF, Gros KS, Davidson TM, Barr S, Cohen J, Deblinger E, Mannarino AP, Ruggiero KJ. National trainers' perspectives on challenges to implementation of an empirically-supported mental health treatment. *Adm Policy Ment Health*. 2014 Jul;41(4):522–34. doi: 10.1007/s10488-013-0492-6. <http://europepmc.org/abstract/MED/23605292> . [PMCID: PMC3758397] [PubMed: 23605292] [CrossRef: 10.1007/s10488-013-0492-6]

40. Hanson R, Self-Brown S, Begle A. Factors related to therapist enrollment and retention in Implementation Research: *Lessons learned from the BRidGE project*. 2011

41. Davidson TM, Soltis K, Albia CM, de AM, Ruggiero KJ. Providers' perspectives regarding the development of a web-based depression intervention for Latina/o youth. *Psychol Serv*. 2015 Feb;12(1):37–48. doi: 10.1037/a0037686.2014-34421-001 [PMCID: PMC4333062] [PubMed: 25133417] [CrossRef: 10.1037/a0037686]

42. Davidson TM, Lopez CM, Saulson R, Borkman AL, Soltis K, Ruggiero KJ, de AM, Wingood GM,

- Diclemente RJ, Danielson CK. Development and preliminary evaluation of a behavioural HIV-prevention programme for teenage girls of Latino descent in the USA. *Cult Health Sex*. 2014;16(5):533–46. doi: 10.1080/13691058.2014.891049. <http://europepmc.org/abstract/MED/24697607> . [PMCID: PMC4020958] [PubMed: 24697607] [CrossRef: 10.1080/13691058.2014.891049]
43. Grubaugh A, Gros K, Davidson T, Frueh Bc, Ruggiero K. Providers' perspectives regarding the feasibility and utility of an Internet-based mental health intervention for veterans. *Psychological Trauma: Theory, Research, Practice, and Policy*. 2014;6(6):624–631. doi: 10.1037/a0035772. [CrossRef: 10.1037/a0035772]
44. Leon A, Davis L, Kraemer H. The role and interpretation of pilot studies in clinical research. *Journal of Psychiatric Research*. 2011 May;45(5):626–629. doi: 10.1016/j.jpsychires.2010.10.008. [PMCID: PMC3081994] [PubMed: 21035130] [CrossRef: 10.1016/j.jpsychires.2010.10.008]
45. Hanson R, Deblinger E, Cohen J, Ruggiero K, Suffoletta-Maierle S. *Unpublished manuscript*. Charleston, SC: Unpublished manuscript; 2004. The Treatment Adherence Checklist-Revised.
46. Chu B, Kendall P. *Child Involvement Rating Scale (CIRS): Scoring manual*. Piscataway, NJ Department of Clinical Psychology, GSAPP, Rutgers University: Unpublished scoring manual; 1999.
47. Chu BC, Kendall PC. Positive association of child involvement and treatment outcome within a manual-based cognitive-behavioral treatment for children with anxiety. *J Consult Clin Psychol*. 2004 Oct;72(5):821–9. doi: 10.1037/0022-006X.72.5.821.2004-19094-010 [PubMed: 15482040] [CrossRef: 10.1037/0022-006X.72.5.821]
48. Chu BC, Kendall PC. Therapist responsiveness to child engagement: flexibility within manual-based CBT for anxious youth. *J Clin Psychol*. 2009 Jul;65(7):736–54. doi: 10.1002/jclp.20582. [PubMed: 19391153] [CrossRef: 10.1002/jclp.20582]
49. Kaufman J, Birmaher B, Brent D, Rao U, Flynn C, Moreci P, Williamson D, Ryan N. Schedule for Affective Disorders and Schizophrenia for School-Age Children-Present and Lifetime Version (K-SADS-PL): initial reliability and validity data. *J Am Acad Child Adolesc Psychiatry*. 1997 Jul;36(7):980–8. doi: 10.1097/00004583-199707000-00021.S0890-8567(09)62555-7 [PubMed: 9204677] [CrossRef: 10.1097/00004583-199707000-00021]
50. Weissman M, Orvaschel H, Padian N. Children's symptom and social functioning self-report scales. Comparison of mothers' and children's reports. *J Nerv Ment Dis*. 1980 Dec;168(12):736–40. [PubMed: 7452212]
51. Faulstich M, Carey M, Ruggiero L, Enyart P, Gresham F. Assessment of depression in childhood and adolescence: an evaluation of the Center for Epidemiological Studies Depression Scale for

- Children (CES-DC) *Am J Psychiatry*. 1986 Aug;143(8):1024–7. [PubMed: 3728717]
52. Pynoos R, Rodriguez N, Steinberg A, Stuber M, Frederick C. *UCLA PTSD Index for DSM IV (Adolescent version)* Los Angeles, CA: UCLA Trauma Psychiatry Service; 1998.
53. Pynoos R, Rodriguez N, Steinberg A, Stuber M, Frederick C. *UCLA PTSD Index for DSM IV (Child version)* Los Angeles, CA: UCLA Trauma Psychiatry Service; 1998.
54. Rodriguez N, Steinberg A, Pynoos R. PTSD Index: Preliminary psychometric analysis of the child and parent versions. The Annual Meeting of the International Society for Traumatic Stress Studies; 2001; New Orleans, Louisiana. 2001.
55. Shirk S, Russell R. *Change Processes in Child Psychotherapy: Revitalizing Treatment and Research*. NY: The Guilford Press; 1996.
56. Shirk S, Saiz Cc. Clinical, empirical, and developmental perspectives on the therapeutic relationship in child psychotherapy. *Develop. Psychopathol*. 2008 Oct 31;4(04):713–28. doi: 10.1017/S0954579400004946. [CrossRef: 10.1017/S0954579400004946]
57. Elvins R, Green J. The conceptualization and measurement of therapeutic alliance: an empirical review. *Clin Psychol Rev*. 2008 Oct;28(7):1167–87. doi: 10.1016/j.cpr.2008.04.002.S0272-7358(08)00085-8 [PubMed: 18538907] [CrossRef: 10.1016/j.cpr.2008.04.002]
58. Lang J, Franks R. *Child/Adolescent Satisfaction Questionnaire*. Farmington, CT: Connecticut Center for Effective Practice; 2007.
59. Pynoos R, Rodriguez N, Steinberg A, Stuber M, Frederick C. *UCLA PTSD Index for DSM IV (Parent version)* Los Angeles, CA: UCLA Trauma Psychiatry Service; 1998.
60. Achenbach T. *Manual for the child behavior checklist/4-18 and 1991 profile*. Burlington, VT: Dept. of Psychiatry, University of Vermont; 1991.
61. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry*. 1961 Jun;4:561–71. [PubMed: 13688369]
62. Beck A, Steer R, Carbin Mg. Psychometric properties of the Beck Depression Inventory: Twenty-five years of evaluation. *Clinical Psychology Review*. 1988 Jan;8(1):77–100. doi: 10.1016/0272-7358(88)90050-5. [CrossRef: 10.1016/0272-7358(88)90050-5]
63. Rizvi SL, Dimeff LA, Skutch J, Carroll D, Linehan MM. A pilot study of the DBT coach: an interactive mobile phone application for individuals with borderline personality disorder and substance use disorder. *Behav Ther*. 2011 Dec;42(4):589–600. doi: 10.1016/j.beth.2011.01.003.S0005-7894(11)00050-5 [PubMed: 22035988] [CrossRef:

10.1016/j.beth.2011.01.003]

64. Horvath AO, Greenberg LS. Development and validation of the Working Alliance Inventory. *J Couns Psychol.* 1989;36:223–33.

Figures and Tables

Table 1

Patient-targeted components of the TF-CBT e-workbook by session.

Treatment concept	TF-CBT e-workbook resource	Modality
One for each component of TF-CBT ^a	Introductory videos that provide an overview for the caregiver and child about why this component of treatment is important.	Video clips featuring adolescent-aged subjects
Each TF-CBT component ^a	Interactive homework assignment checklists with activity suggestions.	Interactive application
Psychoeducation ^a	“What Do You Know?” question and answer quiz game, with “card decks” designed to facilitate child-provider education around trauma, domestic violence, sexual abuse, physical abuse, personal safety, disasters, serious accidents, and bullying/peer victimization; these decks can be personalized to each patient and provider.	Interactive touch screen activity with scorekeeping
Psychoeducation	“You are not alone” interactive graphical display that provides accurate statistics about traumatic events and emotional recovery. The provider selects a question to review with the child, and the child then estimates via touch screen interaction how many of children drawn on the screen have had experiences similar to him/her. Correct answers are given with light up figures.	Interactive touch screen activity
Psychoeducation	“Your Body” cartoon that is designed to	Interactive drag-and-

facilitate accurate labeling of body parts via drop touch screen
 drag-and-drop touch screen activity. Both activity
 genders are represented in this activity

^aThis resource was identified by TF-CBT trainers as a necessary component to the toolkit (Hanson et al, 2014).

Table 2

Summary of qualitative feedback from mental health providers (n=22) during alpha testing.

e-workbook component (activity)	Positive feedback (mental health providers)	Recommendations ^a and observations (mental health providers)
Psychoeducation (What Do You Know? card game)	Liked provider tab	Most providers wanted to organize their own decks
	Activity was clear	Most providers wanted to create their own unique cards
	Liked the game component	One provider asked us to label “explicit content” on cards that are particularly sensitive in nature
	Good for full age range; will increase engagement	Ability to “block” certain cards per patient
	Liked interactivity	Some providers asked for more decks of cards—like cyberbullying and Internet safety questions
Psychoeducation (You Are Not Alone activity)	Difference in race/ethnicity	Include percentages and graphical presentations for older kids
	Would use with kids 8-15 years;	Option to view statistics for

very helpful in teaching
psychoeducation

other gender

^aMany items in the *recommendations* column already have been addressed by the development team. Recommendations made by several providers, such as the ability for providers to organize decks and create new cards in the “*What Do You Know?*” game and the recommendation to add decks addressing Internet safety and cyberbullying, were addressed by the developers in the revised e-workbook prior to the feasibility trial. Other recommendations were only voiced by one or two participants, and were considered on a case-by-case basis. Some observations made by providers related to child age; however, these observations often contradicted the perceptions of children. For example, some providers felt that two or three of the chapters were best for children under 12 years of age; but in each of these cases older adolescents (aged 13-16 years) responded very favorably to the chapters. We have informed providers of this feedback from adolescents and have asked providers to be open-minded in the context of the feasibility trial about the value of each resource on the basis of age.

Table 3

Summary of qualitative feedback from children (n=24) during alpha testing.


e-workbook component (activity)	Positive feedback (children)	Recommendations ^a and observations (children)
Psychoeducation (What Do You Know? card game)	Using technology would help kids be more comfortable	Audio narration that reads card to you
	Most youths said that they would like the iPad-based version over the real cards	Too much white background—include more colors and enhance flexibility Add more colors or brighter colors
		Have themes or animation that relates to the question OR that the kid can choose to “make it their own”
Psychoeducation (You Are Not Alone activity)	N/A—Children were not asked about this activity	
Psychoeducation	N/A—Children were not asked about this activity	

(Your Body activity)

Relaxation (breathing/PMR Coaching activity)	Activity makes sense	Balloon is fine, maybe integrate other graphics, like lungs, or a kid breathing in and out
	Would be helpful to learn to calm down	Change white screens to something else—have options for different background colors (preferred neon)
	App version is very straightforward	Make activity available to kids/teens at home


^aMany items in the *recommendations* column already have been addressed by the development team. Recommendations made by several providers, such as the ability for providers to organize decks and create new cards in the “*What Do You Know?*” game and the recommendation to add decks addressing Internet safety and cyberbullying, were addressed by the developers in the revised e-workbook prior to the feasibility trial. Other recommendations were only voiced by one or two participants, and were considered on a case-by-case basis. Some observations made by providers related to child age; however, these observations often contradicted the perceptions of children. For example, some providers felt that two or three of the chapters were best for children under 12 years of age; but in each of these cases older adolescents (aged 13-16 years) responded very favorably to the chapters. We have informed providers of this feedback from adolescents and have asked providers to be open-minded in the context of the feasibility trial about the value of each resource on the basis of age.

Figure 1



Screenshot of the eight “decks” of virtual cards in the “What Do You Know?” psychoeducation activity. Clinicians and families select the most pertinent decks for each child. The cards within each deck are customizable by the clinician to enhance relevance to each family’s needs.


Figure 2



An example card from the Domestic Violence deck. Users can swipe left or right through the deck to view


additional cards. A scoreboard is presented at the bottom to increase engagement.

Figure 3




A Provider tab is located at the top of the screen and contains specific notes for each card. These notes may be used as discussion points during session.

Figure 4



At the end of each activity, progress is tracked graphically via color-coding (ie, green=complete). On this page, module-specific homework assignments may be assigned. The Menu button at the top of each screen allows users to access the Home screen or end a session and assign homework.

Figure 5




One activity in the Affective Regulation module involves directing the child to list as many different feeling words in 90 seconds as he or she is able to. This activity uses a free-write function. The Provider tab at the bottom of the screen provides tips for clinicians on strategies for using this activity effectively in session.

Figure 6



Another Affective Regulation activity is Feelings Charades. Users “spin” a virtual wheel by dragging down on the feeling words. They are then instructed to act out the feeling on which the wheel lands. A scoreboard is available to increase engagement with the activity.

Figure 7



An activity in the Cognitive Coping module has children practice identifying and categorizing thoughts, feelings, and behaviors by dragging the words and phrases on the left of the screen (shown in gray) to the appropriate icon on the right of the screen. Feedback is provided in response to each user interaction.