



Virilization and abdominal mass in a newborn female: A case report



Natalie A. Drucker, MD^{a, b, *}, Brian W. Gray, MD^a, Johanna Askegard-Giesmann, MD^{a, 1}

^a Division of Pediatric Surgery, Department of Surgery, Indiana University School of Medicine, Riley Hospital for Children at IU Health, 705 Riley Hospital Drive, Indianapolis, IN, 46202, USA

^b Division of General Surgery, Department of Surgery, Indiana University School of Medicine, 545 Barnhill Drive, Emerson Hall 125, Indianapolis, IN, 46202, USA

ARTICLE INFO

Article history:

Received 29 March 2017

Received in revised form

7 May 2017

Accepted 8 May 2017

Available online 10 May 2017

Keywords:

Congenital ovarian tumor

Ambiguous genitalia

Virilization

Juvenile granulosa cell tumor

ABSTRACT

We describe virilization in a newborn female secondary to bilateral congenital juvenile granulosa cell tumor (JGCT). The patient presented with abdominal mass and ambiguous genitalia at birth, and bilateral ovarian masses were discovered on further imaging. The patient underwent bilateral salpingo-oophorectomy in staged procedures, as it became apparent that we could not spare the ovaries. Diagnosis of JGCT was confirmed by surgical pathology. She required no adjuvant therapy and has no signs of recurrence at two-year follow-up.

© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Virilization and ambiguous genitalia in a newborn female is an uncommon and complex problem with many possible causes. The most common cause for virilization in genetically female infants is congenital adrenal hyperplasia (CAH). Other conditions in the differential diagnosis include exposure to maternal androgens or exogenous hormones. We present a case of virilization secondary to bilateral congenital juvenile granulosa cell tumor (JGCT) with no other associated abnormalities.

1. Case report

A 5.2-kilogram 37-week gestation female was born to a 42-year-old G5P2202 woman by Cesarean section for failure to progress. Pregnancy was complicated by polyhydramnios and pregnancy-induced hypertension. Ultrasound at 20 weeks was consistent with a normal female fetus. APGARs at birth were 8 and 9. On physical exam, she was noted to have a palpable abdominal mass and ambiguous genitalia. The mid-abdominal mass was firm, mobile, nontender, and 8–10 cm in diameter. She had clitoromegaly with normal labia majora without evidence of posterior labial

fusion. Her urethral and vaginal orifices appeared normal, and she had a patent anus. Basic metabolic panel was normal. Ultrasound demonstrated an abdominal mass with a cystic component. MRI revealed a normal uterus and bilateral cystic masses of the ovaries. On MRI, the larger mass measured 7.4 cm × 5.2 cm × 4.8 cm, and the smaller one measured 4.0 cm × 2.5 cm × 2.2 cm (Fig. 1).

Diagnostic laparoscopy demonstrated a large mass in the midline originating from the left ovary, as well as a smaller mass originating from the right ovary that had undergone torsion. Examination of the left ovary demonstrated multiple small cysts with no discernible ovarian tissue distinct from the mass. The right ovary was not ischemic and was detorsed. Surgical exploration was expedited before return of any laboratory values because of concern for torsion. With the evidently bilateral nature of the pathology, biopsies were obtained of the larger left mass in hopes that any normal ovarian tissue would be able to be preserved.

Pathology demonstrated juvenile granulosa cell tumor (JGCT) with no residual normal ovary tissue. Hormonal studies are listed in Table 1. On day ten of life, she underwent left salpingo-oophorectomy and right ovarian biopsy with peritoneal washings. It was impossible to dissect the left Fallopian tube free from the mass. Right ovary pathology demonstrated JGCT as well, but peritoneal fluid was negative for neoplastic cells. After consultation with oncology, we elected to continue observation with surgery at a later date, to allow the possibility of ovary-sparing surgery.

She was discharged from the hospital at 5 weeks of age. At

* Corresponding author. Indiana University School of Medicine, Department of Surgery, 545 Barnhill Drive, Emerson Hall 125, Indianapolis, IN, 46202, USA.

E-mail address: ndrucker@iupui.edu (N.A. Drucker).

¹ Present address: Department of Surgery, University of North Dakota, Sanford Children's Hospital – 801 Broadway N, Fargo, North Dakota 58122, USA.




Fig. 1. MRI demonstrating bilateral ovarian masses. Arrowhead: left ovary. Arrow: right ovary.

follow-up, her clitoromegaly had slightly improved, and her right ovary was smaller at $4.2 \times 1.6 \times 1.7$ cm on ultrasound. Her follow-up labs are shown in [Table 1](#). She was readmitted at 4.5 months with small bowel obstruction. She required adhesiolysis, and it was evident that her JGCT had not regressed. Right salpingo-oophorectomy was performed at this time. Pathology demonstrated very small fragments of ovary with multiple nodules and cystic areas of JGCT.

Peritoneal washings were again negative. Therefore, the tumor had been limited to the bilateral ovaries, corresponding to FIGO stage Ib, for which no chemotherapy is indicated. By Children's Oncology Group (COG) staging, it would be stage II.

She was lost to follow-up after discharge for 2 years and seen again at the age of 2.5 years, fortunately without evidence of recurrence. Tumor markers were normal with Inhibin B less than 10 pg/mL and AMH less than 0.1 ng/mL, and ultrasound demonstrated no evidence of recurrent or residual tumor. She will continue long-term surveillance and will require hormone replacement therapy.

2. Discussion

Juvenile granulosa cell tumors (JGCT) are the most common sex cord-stromal tumor (SCST) in the pediatric population. Gynecologic tumors make up less than 5% of pediatric neoplasms, and SCSTs are rare within this group (10%), with germ cell tumors being far more common (60–70%). SCST most commonly presents between four and nine years of age and JGCT are exceedingly rare in infancy [1–4].

In a 20-year series from Pakistan, the median age of presentation of JGCT was fourteen years, with only two occurring before one year of age [5]. Bouffet et al. had similar findings in a comprehensive review in 1997 [1]. Only 25 cases of JGCT in infants less than one year old have been documented [6,7]. Bilateral ovarian JGCT occurs in about three percent of cases, but in infants, this tends to

Table 1
Laboratory Results.

| Lab Test | Initial Result | At Follow-up | Reference Range |
|-------------------------------|----------------|----------------|-------------------|
| Karyotype | 46, XX | – | normal |
| 17- α -OH progesterone | 816 ng/dL | – | <78 ng/dL |
| DHEA | 756 μ g/dL | 35 μ g/dL | 88–356 μ g/dL |
| Testosterone | 588 μ g/dL | <10 μ g/dL | 20–64 μ g/dL |
| Estradiol | 32 pg/mL | <20 pg/mL | <20–53 pg/mL |
| LH | 0.9 mU/mL | – | 0.1–3.3 mU/mL |
| FSH | <0.2 mU/mL | – | 2–15 mU/mL |
| Inhibin A | 77 pg/mL | – | N/A |
| Inhibin B | 345 pg/mL | 195 pg/mL | <111 pg/mL |
| Anti-Mullerian hormone | 18 ng/mL | 32 ng/mL | <4.7 ng/mL |

be associated with other congenital abnormalities (Ollier's disease, Maffucci's syndrome, leprechaunism) [2,8].

Typical presentation of JGCT is abdominal pain and distension, frequently associated with endocrine manifestations such as isosexual pseudoprecocity or virilization in the pre-menarchal population. Occasionally, JGCT may present with peritonitis from torsion or hemorrhagic shock secondary to tumor rupture. Work-up for endocrine disturbances and associated genetic abnormalities is indicated in these patients. Hormone levels should be obtained at baseline as tumor markers.

Once the diagnosis has been established, resection is the appropriate management, as these tumors are hormonally active and do have malignant potential. Ovarian preservation should be emphasized if possible. If normal ovary remains, the tumor should be resected preserving ovarian tissue, especially in the case of bilateral pathology. In this case, there was essentially no normal ovary remaining. Oophorectomy is sufficient in FIGO stage Ia or Ib, but in stage IIa–b, hysterectomy may be considered as well. In infants, surgery may be limited to resection of the cyst or tumor, if it is possible to spare the ovary, as the disease tends to be less aggressive in this age group. Peritoneal washings and biopsy of the contralateral ovary should always be performed [1,2,9].

Chemotherapy is recommended only if the tumor has spread beyond the ovaries. It is uncommon for patients to present with advanced disease. In one study from 2002, seven patients were identified with tumors stage II or higher. Adjuvant cisplatin-based chemotherapy was used with all patients achieving complete clinical remission. Radiotherapy was used in one patient with diffuse peritoneal metastases [10]. Schneider et al., in 2005 described long-term survival and remission in four of six patients with stages II and III JGCT following surgery and adjuvant chemotherapy with cisplatin-based therapy. The two patients that died showed high proliferative activity of the malignant cells, while survivors had lower mitotic counts. Age at diagnosis also appeared to be prognostic, as patients younger than ten years had a more favorable outcome [11].

Treatment of stage Ic tumors, where the tumor is confined to the ovaries but has ruptured or the capsule has been violated at the time of surgery, remains rather controversial. In a study of 62 patients, intraoperative rupture or violation of the tumor capsule did not negatively impact long-term outcome or recurrence, whereas if the tumor had ruptured preoperatively or malignant ascites was present, these patient would likely benefit from chemotherapy [10]. If the rupture occurred intraoperatively and there were ≥ 20 mitoses/10HPF, then adjuvant chemotherapy should be considered [9].

After initial surgical treatment, patients require long-term follow-up and persistent surveillance with imaging and tumor markers. Recurrence has been seen with stage I JGCT up to eight years post-operatively [2].

Interestingly, a case report published in 2009 does provide

evidence of spontaneous regression of JGCT after an infant's parents refused bilateral oophorectomy. Two months after diagnosis of JGCT, the patient had normal hormone levels and this persisted over two years of follow-up [12].

Ovarian sex-cord stromal tumors (SCST), which includes JGCT as a subtype, have also been associated with familial pleuropulmonary blastoma (PPB) and the *DICER1* gene mutation. The ovarian SCST may be the initial clinical presentation of *DICER1* mutations within some families, so genetic testing should be considered in these patients, especially if they have any signs of leprechaunism or if they develop a PPB [13,14]. These families should also be counseled about registering with the PPB Registry at <http://www.ppbregistry.org>.

This is a rare case of congenital bilateral JGCT in a patient with no other genetic abnormalities. Initial surgical management was left salpingo-oophorectomy with delay of right salpingo-oophorectomy in hopes that the tumor would regress and allow ovary-sparing surgery in the future. Especially with bilateral pathology, an ovarian sparing operation should be performed when possible. At her second operation for small bowel obstruction, the tumor remained concerning with no residual normal ovary and right salpingo-oophorectomy was performed. She will continue to follow-up with routine imaging and tumor markers for surveillance of late recurrence.

3. Conclusion

Congenital bilateral JGCT is exceedingly rare. Ovary-sparing surgery should be the primary goal, particularly in infants with bilateral disease, but often this cannot be achieved. With early diagnosis and treatment, outcomes are favorable, but long-term follow-up is imperative to detect any late recurrences.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Financial disclosures

There are no financial relationships relevant to this article to disclose.

Conflicts of interest

There are no conflicts of interest to disclose.

Acknowledgements

Deborah Billmire, M.D. reviewed manuscript and provided expert advice regarding surgical treatment of pediatric ovarian tumors.

James Croop, M.D. reviewed manuscript and provided expert advice regarding oncologic treatment of pediatric ovarian tumors and genetic testing.

References

- [1] Bouffet E, Basset T, Chetail N, Dijoud F, Mollard P, Brunat-Mentigny M, et al. Juvenile granulosa cell tumor of the ovary in infants: a clinicopathologic study of three cases and review of the literature. *J Pediatr Surg* 1997;32(5):762–5.
- [2] Leyva-Carmona M, Vazquez-Lopez MA, Lendinez-Molinos F. Ovarian juvenile granulosa cell tumors in infants. *J Pediatr Hematol Oncol* 2009;31(4):304–6.
- [3] Scully RE. Classification of human ovarian tumors. *Environ Health Perspect* 1987;73:15–25.
- [4] Schultz KA, Ness KK, Nagarajan R, Steiner ME. Adnexal masses in infancy and childhood. *Clin Obstet Gynecol* 2006;49(3):464–79.
- [5] Haroon S, Idrees R, Zia A, Memon A, Fatima S, Kayani N. Ovarian sex cord stromal tumours in children and young girls - a more than two decade clinicopathological experience in a developing country, Pakistan. *Asian Pac J Cancer Prev* 2014;15(3):1351–5.
- [6] Pysker TJ, Hitch DC, Krous HF. Bilateral juvenile granulosa cell tumors in a 4-month-old dysmorphic infant. *A Clin Histol, Ultrastruct Study Am J Surg Pathol* 1981;5(8):789–94.
- [7] Schultz KA, Sencer SF, Messinger Y, Neglia JP, Steiner ME. Pediatric ovarian tumors: a review of 67 cases. *Pediatr Blood Cancer* 2005;44(2):167–73.
- [8] Young RH. Ovarian tumors and tumor-like lesions in the first three decades. *Semin Diagn Pathol* 2014;31(5):382–426.
- [9] Schultz KA, Schneider DT, Pashankar F, Ross J, Frazier L. Management of ovarian and testicular sex cord-stromal tumors in children and adolescents. *J Pediatr Hematol Oncol* 2012;34(Suppl 2):S55–63.
- [10] Schneider DT, Calaminus G, Wessalowski R, Pathmanathan R, Harms D, Gobel U. Therapy of advanced ovarian juvenile granulosa cell tumors. *Klin Padiatr* 2002;214(4):173–8.
- [11] Schneider DT, Calaminus G, Harms D, Gobel U. Ovarian sex cord-stromal tumors in children and adolescents. *J Reprod Med* 2005;50(6):439–46.
- [12] Capito C, Flechtner I, Thibaud E, Emond S, Kalfa N, Jaubert F, et al. Neonatal bilateral ovarian sex cord stromal tumors. *Pediatr Blood Cancer*. 2009;52(3):401–3.
- [13] Schultz KA, Flechtner I, Thibaud E, Emond S, Kalfa N, Jaubert F, et al. Ovarian sex cord-stromal tumors, pleuropulmonary blastoma and *DICER1* mutations: a report from the International Pleuropulmonary Blastoma Registry. *Gynecol Oncol* 2011;122(2):246–50.
- [14] Doros L, Schultz KA, Stewart DR, Bauer AJ, Williams G, Rossi CT, et al. *DICER1*-Related disorders. In: Pagon RA, et al., editors. *GeneReviews*(R); 1993. Seattle (WA).