

Addressing People and Place Microenvironments in Weight Loss Disparities (APP-Me): Design of a randomized controlled trial testing timely messages for weight loss behavior in low income black and white women

Daniel O. Clark^{1,2,3} Preethi Srinivas,² Kunal Bodke,² NiCole Keith,^{1,2} Sula Hood,⁴ Wanzhu Tu^{1,2,4}

Affiliations

¹ Indiana University Center for Aging Research, Indianapolis, Indiana

² Regenstrief Institute, Inc., Indianapolis, Indiana

³ Department of Medicine, Division of General Internal Medicine and Geriatrics, Indiana University School of Medicine, Indianapolis, Indiana

⁴ Indiana University Richard M. Fairbanks School of Public Health, Indianapolis, Indiana

Corresponding Author:

Daniel O. Clark, PhD.

1101 West 10th Street

Indianapolis, IN 46202, United States

daniclar@iupui.edu

This is the author's manuscript of the article published in final edited form as:

Clark, D. O., Srinivas, P., Bodke, K., Keith, N., Hood, S., & Tu, W. (2018). Addressing people and place microenvironments in weight loss disparities (APP-Me): Design of a randomized controlled trial testing timely messages for weight loss behavior in low income Black and White Women. *Contemporary Clinical Trials*. <https://doi.org/10.1016/j.cct.2018.01.006>

Abstract

Background

Behavioral interventions for weight loss have been less effective in lower income and black women. These poorer outcomes may in part be related to these women having more frequent exposures to social and physical situations that are obesogenic, i.e., eating and sedentary cues or situations.

Objectives

Working with obese, lower income black and white women, Addressing People and Place Microenvironments (APP-Me) was designed to create awareness of self-behavior at times and places of frequent eating and sedentary behavior.

Design

APP-Me is being evaluated in a randomized controlled trial with 240 participants recruited from federally qualified health centers located in a single Midwestern city. All participants complete four weeks of ecological momentary assessments (EMA) of situations and behavior. At the end of the four weeks, participants are randomized to enhanced usual care (UC) or UC plus APP-Me.

Methods

APP-Me is an automated short messaging system (SMS). Messages are text, image, audio, or a combination, and are delivered to participants' mobile devices with the intent of creating awareness at the times and places of frequent eating or sedentary behavior. The EMA data inform the timing of message deliveries.

Summary

This project aims to create and test timely awareness messages in a subpopulation that has not responded well to traditional behavioral interventions for weight loss. Novel aspects of the study include the involvement of a low income population, the use of data on time and place of obesogenic behavior, and message delivery time tailored to an individual's behavioral patterns.

Keywords: Randomized controlled trial; weight loss; mobile health; health disparities; user-centered design

Introduction

Obesity rates are high among middle-aged adults,[Ogden] which we here define as 35-64. This is true among poorer and minority adults in particular. Adults living in poverty, for example, have rates that are 50% higher than adults not living in poverty. (1) Racial disparities are also large, and largest in middle-age where 59% of black women have a body-mass index (BMI) of 30 or greater (indicating obesity) while only 36% of white middle-aged women have an obese BMI.(2)

Notably, the National Academy of Medicine (formerly the IOM) identified obesity in the urban poor as high priority research. (3) For middle-aged black women, four of five leading causes of death are obesity-related.(4) The concluding article in an October 2014 *Obesity Reviews* Special Issue on Achieving Healthy Weight in Black American Communities reports that studies of obesity prevention and treatment in black adults are few and insufficient. The authors recommend exploration of electronic health approaches.(5)

Weight loss trials have generally had low representation of adults living in poverty.(6) And while black adults have been represented in large weight loss trials such as the Diabetes Prevention Program (DPP) and Look Ahead,(7) black women in these trials have consistently lost less weight compared to white women.(8-13) Reasons for this are unclear but qualitative studies point to frequent exposure to obesogenic situations for black women.(5, 14) Social support for healthy weight in particular has been hypothesized to be low among black women relative to white.(15-17) [] Regardless, limited success in weight loss among black and poor women translates into significant racial and socioeconomic disparities in health and longevity—a relatively modest weight loss of 2kg has been shown to meaningfully reduce one's risk of hypertension and diabetes.(18)(19)

Several recent studies have tested short message service (SMS) “text” messaging as an approach to monitoring and supporting behaviors conducive to weight loss. (20-24) The theoretical frameworks guiding these studies are social cognitive and self-regulation theories, and transtheoretical model. These studies sent messages from the study staff or machine. The frequency of text messaging ranged from two per week to six times a day. Only one study attempted to send text messages that were coordinated with a participant's reported daily schedule. (23)

Weight loss success was variable across the studies. However, consistently across all of the studies, participants who interacted most with messaging had greater weight loss. Furthermore, only one study enrolled low-income or black women. (23) Attrition was well over 50% at 6 months in this trial, but among those who did not attrit, weight loss was 3.7 kilograms for those receiving supportive texts versus those in standard care.

Our project first measures and compares obesogenic behavior and situations to which urban poor black and white middle-aged women are exposed and, second, tests whether timely messaging via a smartphone application can offset some of the influence of these situations. Based upon the theory of automaticity(25, 26)[] and the input of middle-aged, urban poor women who participated in our iterative user-centered design process, both of which are further discussed below, the messages are intended to give timely awareness for physical activity and portion control, as well as general inspiration for health. The black and white women in the proposed study live in the same neighborhoods of a single urban county and seek care in the same safety-net health system. This health system operates a lifestyle weight loss program called Healthy Me, (10, 27) which is part of enhanced usual care. The purpose of this paper is to describe the study design, assessments, and intervention.

Methods

Addressing People and Place Microenvironments (APP-Me) is funded by the National Institute of Heart, Lung, and Blood Institute. The study is approved by the Indiana University Institutional Review Board and registered in clinicaltrials.gov (NCT03083964).

Setting and population

This study will recruit a sample of 150 non-Hispanic black women and 150 non-Hispanic white women ages 35–64 who receive care in one of eight federally qualified health centers (FQHC). Participants will reside in Indianapolis, Indiana, a city with slightly less than one million persons as of 2016—66% non-Hispanic white and 28% black—and a median household income of \$42,168 (28).

The FQHCs from which participants will be recruited serve a patient population with high rates of poverty—one recently completed study indicated that 50% of obese middle-aged adults in these FQHCs report household incomes of less than \$20,000 per year.(29) Further, two thirds of the participants from this study self-identified as black.

Living in the same city, black and white women in the study will have been exposed to largely the same macro physical environments (e.g., number of fast food restaurants and signs) but perhaps not the same micro level physical or social environments (e.g., food in the household, social network eating behavior). Given the influence of automaticity on weight-related behaviors,(30) addressing the influence of micro level exposures is likely critical to furthering research on disparities in weight loss.

User-Centered Design

In the creation of the awareness tool, we adopted a user-centered design process for the development of consumer-facing health information technology.(31-35) This iterative design process was comprised of three phases that (1) sought to understand the users, their tasks and goals, and the different aspects of their surrounding environments (study phase), (2) design abstract representations followed by more traditional artifacts such as wireframes or user interface prototypes, and (3) evaluate the designs against users' needs and goals.

Phase 1: This study phase included conducting 30-60 minute semi-structured individual interviews with FQHC health coaches and black and white women aged 35-64 years. A basic framework of questions served to probe and understand the characteristics of participants, their perceptions toward nutrition and physical activities, and their corresponding motivations, goals, and barriers. Data collected in this phase were transcribed and used to identify pain points, concerns, needs of users, and implications for design.

We identified three key user requirements that influenced our subsequent design process: (1) lack of information, (2) need for social engagement around weight management, and (3) support while making decisions related to eating or physical activity.

Phase 2: Adapting the work discussed by Iacucci and colleagues (36-39), the design phase included a role-playing game that required players to enact and envision potential products or services that will help support their needs specific to information, social engagement, and

support in healthy weight loss behaviors. This game included a board designed to mimic a city and took into account the mobility of users, their various contexts, activities, and group interactions. Each of three games required 4-5 players, with each player enacting a scenario, identifying user needs for the scenario, proposing a solution, and participating in a group-based discussion of the identified needs and solution by all the players. Participants converged on the necessity for increased awareness of obesogenic behavior, which included healthy eating and increased physical activity. From this, our team envisioned a short messaging system (SMS) aimed at creating awareness at the right place and time to offset the influence of obesogenic behaviors.

We then conducted a series of design sessions to identify the product requirements to design a mobile SMS system. These included: (i) push messages in a timely manner (e.g., during times when participants were likely to consume calories or remain sedentary), (ii) push messages depending on location and social context, (iii) push messages users or their close network ties create in addition to those offered by the study, (iv) push messages in multiple modalities (text, audio, and picture), and (v) collect or receive user feedback on each message. Phase 3: A significant need for a timely and context-sensitive messaging system is understanding the user's behavior and schedule. This motivated the need for a mobile ecological momentary assessment (EMA) system, which is capable of performing experience sampling. Hence, working again with potential participants, our team developed a mobile EMA system. EMA assessments occur in the moment and reduce recall bias⁽⁴⁰⁾ and is considered the gold standard of experiential sampling.⁽⁴¹⁾ EMA question formats were co-designed and evaluated by users, and include eating, drinking, social interaction, physical activity, and location questions. To achieve maximum timely and context learning, we designed an enrollment process where users set approximate times appropriate to receive EMA messages. From within the EMA application, users set approximate times when they wake up, eat, sleep, and wish to not be disturbed. This information guides the timing of EMA questions for each participant. APP-Me uses this EMA information to learn personalized user behavior and send awareness messages to users at the right time.

Following the design and evaluation of EMA component, our team worked with potential participants to develop the initial awareness messaging library. This process involved participants co-designing some sample awareness messages, evaluating the usability of APPME in presenting multi-modal awareness messages, creating personal awareness messages, and providing feedback (as "like" or "dislike") every time they received a message.

In short, our team built a native Android application, APPME, with an enrollment step, EMA capability, personalized message creation and delivery, and user feedback regarding messages (i.e., thumbs up/down). Participants' information and responses are stored on a secure server. APPME uses Firebase Cloud Messaging (FCM), a cross-platform messaging solution that supports reliable delivery of information through push notifications.

Study aim

The study Primary Aim was to randomize participants who have EMA response rates of at least 50 percent to enhanced usual care or enhanced usual care plus APP-Me messaging. APP-Me messages were scheduled for up to four delivery times per day that were partially determined by each participant's EMA patterns (e.g., times of frequent eating). It was hypothesized that, compared to UC participants, APP-Me participants would lose more weight, including a greater proportion achieving weight loss of 2kg or more.

Theoretical framework

The theory of automaticity proposes that behavior is largely based on learned habits, with awareness of behavior low or absent.(25) Habits, and therefore behaviors, are triggered by familiar situations and environments. (26)


Both social and physical environments have been shown to have powerful effects on automatic weight-related behaviors.(42, 43) Experiments have shown that caloric consumption increases from 28% of usual intake while eating with one other person to over 70% while eating with six or more.(30) Moreover, research has shown that, compared to normal weight individuals, obese individuals have an attentional bias when exposed to calorically dense, palatable food cues.(44, 45) Prior research has identified that changing cue exposures may change the behavior. A cue can be virtually anything that triggers a desire, thought, or action in an individual. A laboratory-based study exposed obese individuals to diet words (i.e., a cues) and found less attentional bias toward palatable food words.(44) A recent European randomized trial with overweight dieters offered an “all-you-can-eat”, free bite-sized meat snacks from a tray placed in the corner of a meat store. Participants from this trial entered the meat shop with the smell of grilled chicken coming from a large visible grill in the store. This trial included an experimental arm presenting a prominent poster and a control arm with no poster. The poster exposed participants to an advertisement for a low-calorie recipe available in the store. Control arm participants consumed 40% more snacks than those in the experimental arm.(44) The Theory of Automaticity suggests that exposure to the diet word ‘low calorie’ affects short-term awareness—our brains give more attention to certain stimuli in a situation. Thus we suppose that timely messages may affect attention, positively (more attention to the stimuli) or negatively (less attention to the stimuli).(9)

Participant selection, recruitment and randomization protocol

Recruitment involves several steps. First, health system data managers with access to electronic medical records (EMR) identify patients who meet study BMI and demographic criteria and received a Healthy Me referral. Second, study research assistants (RAs) seek permission from FQHC providers to contact potentially eligible participants. Third, potentially eligible participants are approached at a FQHC visit or receive a telephone call from an RA who invite the patient’s participation in the study. Those indicating interest complete a brief telephone screen. Fourth, those who are eligible are invited to schedule a home visit for informed consent, baseline assessment, and enrollment for EMA. At the end of four weeks of EMA, participants were randomly assigned to the intervention and control groups at a one-to-one ratio and stratified by race, according to a randomization list provided by our biostatistical team. Treatment assignments for individual participants were concealed in envelopes and used sequentially by the study RAs. To select participants with a higher probability of study completion, only those who respond to at least 50 percent of the EMA question “eating these past 15 minutes?” are eligible for randomization. Finally, at 6 months, blinded RAs complete the 6-month follow-up assessment and collect any study equipment. Participant’s weight change from baseline to 6 months was used as the primary outcome of the analysis. Proportions of participants achieving 2 kg weight reduction were compared between the two treatment groups.

Figure 1. Study Recruitment, Randomization, and Assessments.

Figure 1. Study Recruitment, Randomization, and Assessments.



Inclusion and Exclusion Criteria

As noted, eligibility requirements include women within the age range of 35–64yrs, with a BMI of 30 or greater, and who received a Healthy Me referral. In addition, participants are required to be English speaking and self-identify as being non-Hispanic black or non-Hispanic white. Recruitment is stratified such that an equal number of black and white women will be recruited. Those with conditions that may affect weight management are excluded (e.g., psychosis, current cancer treatment, pregnancy or nursing within 6 months, prior bariatric surgery), Current cigarette smokers and alcohol or substance abusers as indicated by their electronic medical record, are also excluded.

Interventions

Enhanced Usual Care. All participants have access to enhanced usual care, which includes access to the Healthy Me program.(10, 27) The Healthy Me program, structured around the 5A's of behavior change,(46) operates in Eskenazi FQHCs with a health coach (47) counseling adult obese patients. The Healthy Me patients are typically referred by their primary care providers after they receive a message about the patient's high BMI in the EMR system. A primary care provider may refer a patient with a BMI of 30 or greater to meet with a Healthy Me coach. Healthy Me coaches are certified in behavior change counseling and fitness instruction, and are present two or more days per week in each of 12 FQHC.

The Healthy Me program emphasizes increased physical activity, healthy food choices, and portion control. Referred patients can meet with coaches to have their current weight-related behavior assessed, receive assistance in problem-solving, and set an action plan for weight loss. The action plan is entered into a Healthy Me database and becomes part of the patient's EMR. Health coaches give patients guidance and information on community-based opportunities for physical activity and also provide weekly FQHC based exercise and nutrition

classes. In addition, dietary and physical activity self-monitoring instructions and logs are provided. A “passport to wellness” incentive program gives patients points for participation that earn them small incentive rewards (e.g., t-shirt, produce coupons, gym trial). If desired, patients can also meet with the FQHC dietitian for nutrition guidance. A consistent message of the program is “eat less, move more.”

APP-Me. The APP-Me intervention is in addition to Healthy Me and consists of personalized, automated messages. The messages are intended to bring participants’ attention and awareness back to healthy eating and physical activity. The messages that achieve this are likely very personal. Hence, the system we created allows participants to create and rate messages. Messages are coded by broad themes (e.g., activity, food, neutral) and subthemes (e.g., gardening, inspiration, grocery shopping, eating out, walking, family, religion, celebrities). An example of a food themed message with a religion subtheme is shown in Figure 2. This message was created by the study team. An example of a neutral themed inspirational message created by a participant during design work is shown in Figure 3.




Figure 2. Example of a study team created food themed message with religious subtheme.



“Mom, you can do it. Don’t give up!”

Figure 3. Example of a participant created neutral themed inspirational message.

Both study staff and participants create the messages. Our team uses a web-based system to produce a master library of messages. As for participant-created messages, text is created and saved, images are taken or selected from camera gallery, and audio-recordings are made—all within the app. When a participant creates a message, it is saved to her message library.

To avoid predictability, messages are randomly generated from both the master and participant libraries. As these were received, participants are able to respond by selecting a thumbs up or thumbs down icon. These responses provide data for the prioritizing or deletion of messages from a participant’s library.

Participants receive a minimum of four messages per day. Every two weeks, participants have the opportunity to increase this frequency by one message. Delivery is scheduled for morning, mid-day, early and late evening but the actual times of these four periods are based upon each participant’s onboarding selections (e.g., wake and physical activity times) and EMA-determined times or places of frequent eating. Onboarding data automatically populate the message

scheduler but EMA patterns are reviewed by team members prior to being considered for automatic scheduling. By initially observing patterns and accumulating data, we will be able to create an algorithm of EMA patterns that will then allow automated message scheduling based on EMA data.

Measures

Baseline RA-administered assessments are completed in participants' homes to allow a physical environment assessment of the home. The in-home baseline assessment requires less than 30 minutes. Interviewer-administered assessments and surveys are carried out using Research Electronic Data Capture (REDCap). Table 1 shows the assessments and schedule.

We obtain some covariates to allow us to adjust for potentially important correlates of weight loss or gain and any group differences. Age, gender, race, and smoking are obtained initially from EMR data but confirmed at the time of screening. Literacy and numeracy affects weight management and varies by ethnicity(48) and we use the New Vital Sign as a representation of both. (49) Food security and binge eating are measured as covariates but also as screeners for need of assistance. Our IRB protocol contains a procedure for referral in cases of a positive screen. Eating disorders (i.e., anorexia nervosa), however, are rare in our target population and a formal screen is not practical.

Table 1. Measures

Construct or Object	Measure or Source	Frequency
Age	EMR	Baseline (B)
Household income	Interview Qs	B
Employment hours	Interview Qs	B
Years of education	Interview Qs	B
Health literacy and numeracy	New Vital Sign(49)	B
Current/Desired weight and body shape	Interview Qs and body figure rating scale	B, 6 months
Disease diagnoses	EMR	B, 6
Prescription medications	EMR	B, 6
Smoking status	Interview Qs	B, 6
Perceived health	SF-36 scale	B, 6
Depression and Anxiety	PHQ-2 and GAD-2 scales	B, 6
Food security	USDA 6-item Form USDA Food Security Survey(50)	B, 6
Binge eating	SCOFF screener(51)	B, 6
Physical activity	Accelerometry	4 weeks
Eating frequency	EMA	4 weeks
Social network	Interview Qs	B
Physical home environment	Interview Qs and home assessment	B
Social contacts	EMA	4 weeks
Height	Portable stadiometer	B
Weight (primary outcome)	Portable scale	B, 6 months

The baseline assessment includes a semi-structured individual interview with questions on how many televisions are in the home, whether there is a scale or fitness equipment, how often fast food, chips, sodas, fruit juices, desserts, and fried foods are in the home, and whether these foods are visible or hidden. The RA also records whether there is a television in or visible from the kitchen, whether a television was turned on when entering the home, and whether there is visible food, frying pans, or fitness equipment.

The RAs also measure the participants' weight using the Tanita WB-800S scale, which has a capacity of 660lbs +/- 0.1lbs. Participants stand flat footed, shoes removed, while wearing light clothing. Two separate measurements are taken and averaged; if weight differs by more than 0.2lbs, a third measurement is taken; and the least congruent measurement is dropped.

Immediately following RA-administered assessments, participants completed onboarding questions within the study app, which includes setting a usual schedule of sleep, wake, and do not disturb times to inform the timing of EMA assessments. We have chosen a maximum of three samples in each quadrant of a participant's waking day; or a maximum of 12 samples per day. As our investigation of eating behavior patterns does not require detailed data on food type or energy intake, to minimize burden and maximize response the EMA assessment asks a single eating and single drinking question, "Eating now or in the past 15 minutes?" [Yes, No] and "Drinking now or in the past 15 minutes?" [No, nothing to drink, Water, Coffee/Tea—no milk or sugar, Other].

The APP-Me app captures location (as longitude and latitude) every 10 seconds from the mobile device. Using the Google Places API, we obtained a probability that the captured latitude and longitude represents a particular place. To improve this, when it appears from GPS that a participant's location changed we ask participants via EMA to report their current location (e.g., home, work, restaurant, retail store, etc). Over time, our system constructs a library of

frequented places for a participant and any future latitude and longitude readings with coordinates within 50 meters of a participant's frequent place are assumed to be that place.

We ask participants to give the first names of up to four persons (ties) they feel close to, any additional household members, and how many people live in their household. This facilitates responses to the EMA question "With anyone?"—the first names of close contacts are displayed as response options. To avoid reminding participants of possible loneliness, this question is not asked when at home for participant's who report living alone.

EMA responses, GPS location, and accelerometry data are captured from the mobile devices are stored in a secure Indiana University server, and later merged with interviewer-administered data stored in REDCap. Participants either use their own smartphone, or a study supplied smartphone (Google Pixel, 32 GB).

Sample Size and Power

The sample size of this study was determined to ensure adequate power for assessing the effect of the APP-Me intervention on weight loss. For this aim, we assume that weight loss in the study sample has a standard deviation of 10lbs (4.535 kg). We intend to recruit and prospectively follow 240 women, with 120 women in each treatment arm, to estimate the intervention effect. Assuming an attrition of 20% based on our prior work in this population[Counsell et al, 2006; Clark et al, 2018], the proposed sample size of 300 (n=150 black and n=150 white women) gives us an expected sample of 240 at 6 months, which is 92% power to detect a 2.0 kg difference in weight changes between the two intervention arms using a two-sample t test at the 0.05 significance level. A 2.0 kg weight loss was judged to be clinically meaningful in that it has been shown in randomized trials to result in a 20% reduction in 3yr risk of hypertension[Stevens] and a 32% reduction in 3yr risk of type 2 diabetes.[Hamman] If we dichotomize the outcome as success (and failure) of achieving 2kg weight loss, using a chi-square test, we will have 86% power to detect a 20% difference in the rate of weight loss success between the two treatment arms (e.g., 28% vs 48%).

Analysis

The primary Specific Aim is to assess 6 month weight change between the two study arms. The main hypothesis is that more participants in the APP-Me arm will lose 2kg or more by the end of follow-up. We will test this hypothesis using a chi-square test. We will secondarily re-examine the balance of baseline characteristics of the study subjects by race and by intervention arm, as well as other relevant characteristics, such as the baseline weight. In necessary, effects of unbalanced participant characteristics will be adjusted as covariates in logistic regression analysis. Similarly, we will also assess the effects of obesogenic exposures on weight loss in the enhanced usual care arm using weight change as the outcome in a linear regression analysis. With the real time exposure data generated by the study, we will evaluate the influences of individual exposures on weight loss in the UC subjects. The effects of other contextual variables such as places of eating and activity, or lack thereof, can be accommodated similarly. We hypothesize that less frequent exposure to obesogenic cues is associated with greater weight loss. As a part of the secondary exploration, we will conduct separate analyses for black and white women to quantify the efficacy of the treatment in each race group, and to estimate race-specific magnitudes of exposure influence. We will assess the racial disparity in 6 month weight change. We will explore whether racial disparities in 6 month weight change are less in the APP-Me arm by comparing the amount of weight loss differences

due to intervention between blacks and whites. We will include a race by intervention interaction in the linear regression model analysis to test the potentially differential treatment effects in the two race groups. As in previous studies,(52) the difference in the race coefficients with and without exposure variables will represent the effect size. All analysis will be conducted using SAS software. P-values less than 0.05 will be considered as statistically significant.

Summary

APP-Me is an outcome of more than a decade of collaboration in health promotion with safety-net health care providers and the patients they serve. This project was inspired by the need to support patients outside of usual clinic physical space and time when patients are experiencing the situations that trigger eating or sedentary behavior (e.g., at home alone and at evening times) and to increase the frequency of support for weight management, which may be especially helpful for persons exposed to significant obesogenic environments and behavior in their daily lives. APP-Me is also inspired by recent advances in theoretical models of automatic behaviors and the influence of momentary situations on those behaviors. APP-Me is enabled by very recent advances in mobile technology, our highly diverse project team, and our embedment in this large urban safety-net health system.

Our project is innovative in several ways. First, we have worked for approximately 12 months with urban low-income women with obesity in the design of both EMA and awareness messaging. The app was constructed with their input at every step – few health applications have been designed with and for urban low-income women. Second, our use of an intense 4-week period of EMA data collection as a guide to the timing of awareness message delivery is novel. Third, we are aware of just one study that has incorporated user-created messages into a weight loss intervention such as APP-Me (53, 54). Fourth, APP-Me is being tested on top of a traditional coach-based model of weight loss support that is operated within safety-net clinical sites.

Commercial apps exist that provide something similar to the awareness messages APP-Me is testing. Fitbit, for example, sends messages that are intended to be motivating in relation to goals set by users within the app. We are not aware that the impact of these messages has been tested. We are also not aware of a weight management or fitness app that uses EMA data to guide the timing of messaging. MoodMission is an app that comes closest perhaps in that it has an onboarding process that includes questions about the times of day your mood may need support. Messages are then delivered at those times. With very many commercial apps and new products and existing product updates continuously added, it is likely that something similar to APP-Me exists or will arise. As pragmatic behavioral scientists, our objective is to design weight loss solutions for those most vulnerable to obesity and to establish an evidence-base for potential users or to offer guidance for future improvements or redirections.

Whether a machine-delivered message can carry the emotional support required for momentary behavior change is not known. Certainly, impactful message designs and optimal delivery timing are highly personalized and require continuous learning and tailoring. Sustaining engagement in such messages over time again points to a need for personalization and continuous learning. Many commercial apps enlist social ties to provide accountability and social engagement (e.g., Twinbody, Thirty, PumpUp). Social networks are smaller in lower-income adults (55), so future efforts to include social ties in APP-Me may require working with providers, pastors, or persons who may not be known to the user but who also use the app. APP-Me is a first step in the direction of continuous learning for personalization of scalable awareness messaging around weight-related behaviors in a highly vulnerable and understudied

population. We anticipate that the APP-Me platform will provide guidance for future work by us and others.

Acknowledgments

This project is supported by National Institutes of Health Grant R01 HL128494.

ACCEPTED MANUSCRIPT

References

1. McGovern L, Miller G, Hughes-Cromwick P. The relative contribution of multiple determinants to health outcomes: Project HOPE; 2014.
2. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011-2012. *JAMA : the journal of the American Medical Association*. 2014;311(8):806-14.
3. Institute of Medicine. *Comparative Effectiveness Research*. Washington, DC: 2009.
4. National Center for Health Statistics. *Healthy People 2010 Final Review*. Atlanta, GA: Centers Disease Control and Prevention; 2013.
5. Kumanyika SK, Whitt-Glover MC, Haire-Joshu D. What works for obesity prevention and treatment in black Americans? *Research directions. Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2014;15 Suppl 4:204-12.
6. Samuel-Hodge CD, Gizlice Z, Cai J, Brantley PJ, Ard JD, Svetkey LP. Family functioning and weight loss in a sample of african americans and whites. *Annals of behavioral medicine : a publication of the Society of Behavioral Medicine*. 2010;40(3):294-301.
7. Pi-Sunyer X. The look AHEAD trial: a review and discussion of its outcomes. *Current nutrition reports*. 2014;3(4):387-91.
8. The Diabetes Prevention Program: baseline characteristics of the randomized cohort. The Diabetes Prevention Program Research Group. *Diabetes care*. 2000;23(11):1619-29.
9. Bargh JA, Chen M, Burrows L. Automaticity of social behavior: direct effects of trait construct and stereotype-activation on action. *Journal of personality and social psychology*. 1996;71(2):230-44.
10. Clark D, Chrysler L, Perkins A, Keith NR, Willis DR, Abernathy G, et al. Screening, referral, and participation in a weight management program implemented in five CHCs. *Journal of health care for the poor and underserved*. 2010;21(2):617-28.
11. Clark DO, Keith N, Chrysler L, Perkins AJ, Willis D. Design and reach of a primary care weight management program. *J Health Care Poor Underserved*. 2008b;19(1):171-9.
12. Bronner Y, Boyington JE. Developing weight loss interventions for African-American women: elements of successful models. *J Natl Med Assoc*. 2002;94(4):224-35.
13. Kumanyika SK, Wadden TA, Shults J, Fassbender JE, Brown SD, Bowman MA, et al. Trial of family and friend support for weight loss in African American adults. *Archives of internal medicine*. 2009;169(19):1795-804.
14. Keith NR, Xu H, de Groot M, Hemmerlein K, Clark DO. Identifying Contextual and Emotional Factors to Explore Weight Disparities between Obese Black and White Women. *Clinical medicine insights Women's health*. 2016;9(Suppl 1):85.
15. Ard JD, Greene LE, Malpede CZ, Jefferson WK. Association between body image disparity and culturally specific factors that affect weight in Black and White women. *Ethnicity & disease*. 2007;17(2 Suppl 2):S2-34-9.
16. Kumanyika SK, Whitt-Glover MC, Gary TL, Prewitt TE, Odoms-Young AM, Banks-Wallace J, et al. Expanding the obesity research paradigm to reach African American communities. *Preventing chronic disease*. 2007;4(4):A112.
17. Smith Barnes A, Kimbro RT. Descriptive study of educated African American women successful at weight-loss maintenance through lifestyle changes. *Journal of general internal medicine*. 2012;27(10):1272-9.
18. Stevens VJ, Obarzanek E, Cook NR, Lee IM, Appel LJ, Smith West D, et al. Long-term weight loss and changes in blood pressure: results of the Trials of Hypertension Prevention, phase II. *Annals of internal medicine*. 2001;134(1):1-11.
19. Hamman RF, Wing RR, Edelstein SL, Lachin JM, Bray GA, Delahanty L, et al. Effect of weight loss with lifestyle intervention on risk of diabetes. *Diabetes care*. 2006;29(9):2102-7.

20. Agboola S, Jethwani K, Lopez L, Searl M, O'Keefe S, Kvedar J. Text to move: a randomized controlled trial of a text-messaging program to improve physical activity behaviors in patients with type 2 diabetes mellitus. *Journal of medical Internet research*. 2016;18(11).
21. Kinnafick F-E, Thøgersen-Ntoumani C, Duda J. The effect of need supportive text messages on motivation and physical activity behaviour. *Journal of behavioral medicine*. 2016;39(4):574-86.
22. Fischer HH, Fischer IP, Pereira RI, Furniss AL, Rozwadowski JM, Moore SL, et al. Text message support for weight loss in patients with prediabetes: a randomized clinical trial. *Diabetes care*. 2016;39(8):1364-70.
23. Lin M, Mahmooth Z, Dedhia N, Frutchey R, Mercado CE, Epstein DH, et al. Tailored, interactive text messages for enhancing weight loss among African American adults: the TRIMM randomized controlled trial. *The American journal of medicine*. 2015;128(8):896-904.
24. Shapiro JR, Koro T, Doran N, Thompson S, Sallis JF, Calfas K, et al. Text4Diet: a randomized controlled study using text messaging for weight loss behaviors. *Preventive medicine*. 2012;55(5):412-7.
25. Marteau TM, Hollands GJ, Fletcher PC. Changing human behavior to prevent disease: the importance of targeting automatic processes. *Science*. 2012;337(6101):1492-5.
26. Fogg BJ, editor A behavior model for persuasive design. *Proceedings of the 4th international Conference on Persuasive Technology*; 2009: ACM.
27. Clark DO, Frankel RM, Morgan DL, Ricketts G, Bair MJ, Nyland KA, et al. The meaning and significance of self-management among socioeconomically vulnerable older adults. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*. 2008;63(5):S312-S9.
28. QuickFacts A. US Census Bureau. Internet Website: <http://quickfacts.census.gov> (accessed 22 May 2009). 2010.
29. Keith NRX, H, Weiner M, Clark DO. An RCT of Clinic, Community, and Video-Conference Behavior Change Counseling for Weight Loss in Urban Poor Adults (in preparation). 2017.
30. Wansink B. *Mindless Eating: Why We Eat More Than We Think*. Reprint Edition ed. New York, NY: Bantam Books; 2010.
31. Beyer H, Holtzblatt K. *Contextual Design: A Customer-Centered Approach to Systems Designs* (Morgan Kaufmann Series in Interactive Technologies). 1997.
32. Buxton B. *Sketching user experiences: getting the design right and the right design*: Morgan Kaufmann; 2010.
33. Nielsen J. Iterative user-interface design. *Computer*. 1993;26(11):32-41.
34. Rogers Y, Sharp H, Preece J. *Interaction design: beyond human-computer interaction*: John Wiley & Sons; 2011.
35. Snyder C. *Paper prototyping: The fast and easy way to design and refine user interfaces*: Morgan Kaufmann; 2003.
36. Iacucci G, Mäkelä A, Ranta M, Mäntylä M, editors. Visualizing context, mobility and group interaction: role games to design product concepts for mobile communication. *Proceedings COOP*; 2000.
37. Iacucci G, Kuutti K. Everyday life as a stage in creating and performing scenarios for wireless devices. *Personal and Ubiquitous Computing*. 2002;6(4):299-306.
38. Iacucci G, Iacucci C, Kuutti K, editors. Imagining and experiencing in design, the role of performances. *Proceedings of the second Nordic conference on Human-computer interaction*; 2002: ACM.
39. Kuutti K, Iacucci G, Iacucci C, editors. Acting to know: improving creativity in the design of mobile services by using performances. *Proceedings of the 4th conference on Creativity & cognition*; 2002: ACM.
40. Shiffman S, Stone AA, Hufford MR. Ecological momentary assessment. *Annu Rev Clin Psychol*. 2008;4:1-32.

41. Kahneman D, Krueger AB, Schkade DA, Schwarz N, Stone AA. A survey method for characterizing daily life experience: the day reconstruction method. *Science*. 2004;306(5702):1776-80.
42. Wansink B. *Mindless Eating: Why We Eat More Than We Think*. New York, NY: Bantam Books; 2006.
43. Kremers SP, Eves FF, Andersen RE. Environmental changes to promote physical activity and healthy dietary behavior. *J Environ Public Health*. 2012;2012:470858.
44. Papias EK. Goal Priming in Dieters: Recent Insights and Applications. *Curr Obes Rep*. 2012;1(2):99-105.
45. Hofman W, van Koningsbruggen GM, Stroebe W, Ramanathan S, Aarts H. As pleasure unfolds. *Psychol Sci*. 2010;21(12):1863-70.
46. Whitlock EP, Orleans CT, Pender N, Allan J. Evaluating primary care behavioral counseling interventions: an evidence-based approach. *American journal of preventive medicine*. 2002;22(4):267-84.
47. Bennett HD, Coleman EA, Parry C, Bodenheimer T, Chen EH. Health coaching for patients with chronic illness. *Fam Pract Manag*. 2010;17(5):24-9.
48. Rothman RL, Housam R, Weiss H, Davis D, Gregory R, Gebretsadik T, et al. Patient understanding of food labels: the role of literacy and numeracy. *American journal of preventive medicine*. 2006;31(5):391-8.
49. Weiss BD, Mays MZ, Martz W, Castro KM, DeWalt DA, Pignone MP, et al. Quick assessment of literacy in primary care: the newest vital sign. *Ann Fam Med*. 2005;3(6):514-22.
50. Blumberg SJ, Bialostosky K, Hamilton WL, Briefel RR. The effectiveness of a short form of the Household Food Security Scale. *Am J Public Health*. 1999;89(8):1231-4.
51. Morgan JF, Reid F, Lacey JH. The SCOFF questionnaire: assessment of a new screening tool for eating disorders. *Bmj*. 1999;319(7223):1467-8.
52. Stewart JC, Zielke DJ, Hawkins MA, Williams DR, Carnethon MR, Knox SS, et al. Depressive symptom clusters and 5-year incidence of coronary artery calcification: the coronary artery risk development in young adults study. *Circulation*. 2012;126(4):410-7.
53. de Vries RA, Truong KP, Kwint S, Drossaert CH, Evers V, editors. *Crowd-Designed Motivation: Motivational messages for exercise adherence based on behavior change theory*. Proceedings of the 2016 CHI Conference on Human Factors in Computing Systems; 2016: ACM.
54. de Vries RA, Zaga C, Bayer F, et al. Experts Get Me Started, Peers Keep Me Going: Comparing Crowd versus Expert-Designed Motivational Text Messages for Exercise Behavior Change. *EAI International Conference on Pervasive Computing Technologies for Healthcare: ACM*; 2017.
55. Christakis NA, Fowler JH. The spread of obesity in a large social network over 32 years. *N Engl J Med*. 2007;357(4):370-9.