

1
2
3
4 **Using an integrated conceptual framework to investigate parents' HPV vaccine decision for**
5 **their daughters and sons**
6
7
8

9
10 Gilla K. Shapiro,^{1,2,*} Ovidiu Tatar,² Rhonda Amsel,¹ Gillian Prue,³ Gregory D. Zimet,⁴ Barbel
11 Knauper,¹ Zeev Rosberger.^{1,2,5,6.}
12
13

14
15 ¹ Department of Psychology, McGill University, 1205 Dr. Penfield Avenue, Montreal, Quebec,
16 H3A 1B1
17

18
19 ² Lady Davis Institute for Medical Research, Jewish General Hospital, 4333 Côte St-Catherine
20 Road, Montreal, Quebec, H3T 1E4
21

22
23 ³ School of Nursing and Midwifery, Medical Biology Centre, Queen's University Belfast, 97
24 Lisburn Road, Belfast BT9 7BL, Northern Ireland
25

26
27 ⁴ Section of Adolescent Medicine, Department of Pediatrics, Indiana University School of
28 Medicine, Indianapolis, Indiana, USA
29

30
31 ⁵ Louise Granofsky Psychosocial Oncology Program, Jewish General Hospital, 4333 Côte St-
32 Catherine Road, Montreal, Quebec, H3T 1E4
33

34
35 ⁶ Departments of Psychiatry and Oncology, McGill University, 1205 Dr. Penfield Avenue,
36 Montreal, Quebec, H3A 1B1.
37
38
39
40

41 **Corresponding author:** Gilla K. Shapiro, Department of Psychology, McGill University, 1205
42 Dr. Penfield Avenue, Montreal, Quebec, H3A 1B1; telephone: +1-514-340-8222, ext. 3978;
43 Email: gilla.shapiro@mail.mcgill.ca
44
45
46
47

48 **Abstract Word Count:** 247 words
49
50

51 **Manuscript Word Count:** 3,554
52
53
54
55
56
57
58
59
60

Abstract

Despite being an effective cancer prevention strategy, human papillomavirus (HPV) vaccination in Canada remain suboptimal. This study is the first to concurrently evaluate HPV vaccine knowledge, attitudes, and the decision-making stage of Canadian parents for their school-aged daughters and sons. Data were collected through an online survey from a nationally representative sample of Canadian parents of 9-16 year old children from August to September 2016. Measures included socio-demographics, validated scales to assess HPV vaccine knowledge and attitudes (using the Health Belief Model), and parents' HPV vaccination adoption stage using the Precaution Adoption Process Model (PAPM; six stages: unaware, unengaged, undecided, decided not, decided to, or vaccinated). 3,779 parents' survey responses were analyzed (1,826 parents of sons and 1,953 parents of daughters). There was a significant association between child's gender and PAPM stage of decision-making, with parents of boys more likely to report being in earlier PAPM stages. In multinomial logistic regression analyses parents of daughters (compared to sons), parents of older children, and parents with a health care provider recommendation had decreased odds of being in any earlier PAPM stage as compared to the last PAPM stage (i.e. vaccinated). Parents who were in the 'decided not to vaccinate' stage had significantly greater odds of reporting perceived vaccine harms, lack of confidence, risks, and vaccine conspiracy beliefs. Future research could use these findings to investigate theoretically informed interventions to specifically target subsets of the population with particular attention towards addressing knowledge gaps, perceived barriers, and concerns of parents.

Keywords

Human papillomavirus vaccination; vaccine uptake; vaccine intentions; vaccine hesitancy; determinants of health; gender; health behaviour change; Health Belief Model; Precaution Adoption Process Model

Highlights

- Canadian parents report suboptimal HPV vaccination rates for their children
- An integrated conceptual framework used the Health Belief Model (HBM) and Precaution Adoption Process Model (PAPM)
- Parents of boys were more likely to report being in earlier PAPM stages
- PAPM stages were associated with different psychosocial correlates
- Tailored health messages could be developed to target individuals at different PAPM stages

Introduction

Human papillomavirus (HPV) can cause a number of anogenital and oropharyngeal cancers in men and women.^{1,2} To prevent morbidity and mortality, three vaccines have been licensed and recommended for use.^{3,4} Currently, over 80 countries have implemented national HPV vaccination programs.^{1,5,6} In Canada, provinces and territories have implemented publicly funded school-based HPV vaccine programs. All Canadian jurisdictions implemented programs for girls, from 2007-2010.^{7,8} As of 2018, all jurisdictions also offer programs for boys in schools;⁹ however, the roll out of these programs (since 2013) has been staggered and HPV vaccination rates in Canada remain suboptimal.^{3,8,10}

Because HPV vaccination targets children,³ parental acceptance is critical to ensuring uptake. Previous research has indicated common themes associated with uptake, such as the importance of parents believing in the benefits of vaccination and perceiving few barriers.^{11,12} Unsurprisingly, parents are less likely to vaccinate their children if they are not aware of, or do not know enough about, HPV vaccination. Parents are also less likely to vaccinate their child if they believe that HPV vaccination can cause harm, or that vaccination is not accessible or affordable.¹³ Furthermore, positive attitudes towards vaccines in general are related to HPV vaccine acceptance.^{14,15} Notably, a strong health care provider (HCP) recommendation significantly improves parental vaccine acceptance.^{4,11,16} Other social influences, including by a partner, family, friends, or online social network, can also influence parents' decision.^{13,16-18}

It is likely that these factors have varying impact on parents depending on where they are in the decision-making process, which is obscured in much previous research investigating vaccination as a binary outcome (vaccinated or not). Literature on vaccine hesitancy highlights many reasons a parent may delay or refuse vaccination for their child.¹⁹ A theoretical stage-based model, the Precaution Adoption Process Model (PAPM), allows for a nuanced examination of which modifying factors and individual health beliefs are important for each stage of decision-making (Figure 1).²⁰⁻²² The PAPM identifies six stages involved in making a health decision and clarifies what factors lead individuals to move from one health behaviour decision-making stage to the next.²³ A stage-based understanding of HPV vaccine decision-making is important for identifying the psychosocial correlates for each stage and how to best intervene for parents at different stages. Nevertheless, few studies have examined the stages of HPV vaccine decision-

1
2
3
4 making in college students and parents of only boys,^{16,24-26} and no study has compared the stages
5 of decision-making of parents of girls to parents of boys. Previous studies have found that
6 college students and parents of boys were in the earliest stages of HPV vaccine decision-
7 making.^{16,24-26} Given HPV vaccine programs and policies have differentially targeted boys and
8 girls, it is important to examine differences in decision-making stage between parents of girls and
9 boys.

10
11
12
13
14
15 This study will identify and compare parents' stage of decision-making by gender for
16 their school-aged daughters and sons, examine differences in parents' HPV vaccine knowledge
17 and attitudes by PAPM stage, and investigate the psychosocial correlates of parents' PAPM
18 stages.
19
20
21
22
23
24
25

26 **Methods**

27 **Survey design and participants**

28
29
30
31
32
33 Details of the methodology are presented in the protocol paper.²⁰ This study used a cross-
34 sectional design to collect self-reported online survey data from a national sample of Canadian
35 parents. Data presented here were part of a larger two-wave protocol and were collected from
36 August 17 to September 11, 2016 (i.e. Time one). All Canadian jurisdictions at this time had
37 publicly funded, school-based HPV vaccination programs for girls but only three provinces (i.e.
38 Alberta, Prince Edward Island, and Nova Scotia) had programs for boys.
39
40
41
42
43

44 This study targeted parents and/or guardians (hereafter referred to as parents) of 9-16
45 year-old boys and girls. Parents with more than one child were asked to answer the questionnaire
46 in reference to the child who had the most recent birthday to ensure randomization. The online
47 survey was offered in English and French (i.e. Canada's two official languages). Participants
48 were recruited using, Leger-*The Research Intelligence Group*, which maintains a nationally
49 representative panel of 400,000 Canadians.²⁷ This study received Research Ethics Board
50 approval from the Research Review Office, Integrated Health and Social Services University
51 Network for West-Central Montreal (CODIM-FLP-16-219).²⁰
52
53
54
55
56
57
58
59
60

Measures

The dependent variable was parents' PAPM stage, which categorizes parents' stage of decision-making regarding HPV vaccination into six stages (Figure 1).²¹

Potential psychosocial predictors of HPV vaccine decision-making included socio-demographics, HCP recommendation, as well as validated scales to assess HPV and HPV vaccine knowledge, HPV vaccine attitudes, and general vaccine attitudes. HCP recommendation was assessed by asking parents, 'did a health care provider (e.g. a doctor, pediatrician, or nurse) recommend that [child's name] receive the HPV vaccine within the last 12 months?'. Parents were only administered this question if they had answered affirmatively that they had seen a HCP and discussed their child receiving the HPV vaccine with a HCP.

Two validated scales were used to measure parents' knowledge of HPV and HPV vaccine.^{28,29} The 23-item HPV General Knowledge Scale ($\alpha=.94$) and the 11-item HPV Vaccine Knowledge (VK) Scale ($\alpha=.88$) (Appendix 1). To each item, respondents answered 'true', 'false', or 'don't know', for which a total score was calculated based on correct answers (higher scores indicate greater knowledge on both scales).

HPV vaccine attitudes were assessed using constructs from the Health Belief Model (HBM) including perceived benefits of, and barriers to, HPV vaccination; perceived severity of, and susceptibility to, HPV infection and disease; external influences prompting HPV vaccine uptake (i.e. cues to action), and the ability to exert change (i.e. self-efficacy). Sub-scales from the psychometrically validated HPV vaccination Attitudes and Beliefs Scale (HABS) were used to evaluate constructs from the HBM using a 7-point Likert-type rating scale ranging from 1 (strongly disagree) to 7 (strongly agree).²⁹ Sub-scales were evaluated for internal consistency using Cronbach's α . HBM constructs, assessed using HABS subscales, included perceived susceptibility of child to HPV and its consequences (3 items, $\alpha=.92$), perceived severity of HPV and its consequence (3 items, $\alpha=.84$), perceived benefits of HPV vaccine (10 items, $\alpha=.94$), perceived barriers to HPV vaccine (6 items to measure harms, $\alpha=.93$; 4 items to measure accessibility, $\alpha=.79$; and 3 items to measure affordability, $\alpha=.87$), cues to action (8 items, $\alpha=.91$), and self-efficacy (4 items $\alpha=.89$) (Appendix 1).¹

¹ All scales are subscales the HABS except self-efficacy, which is a construct of the HBM but was not included as a subscale in the HABS.

1
2
3
4 General vaccine attitudes were assessed using two psychometrically validated scales: the
5
6 Vaccine Conspiracy Beliefs Scale (VCBS) and the Vaccine Hesitancy Scale (VHS).^{14,15} The
7
8 VCBS has seven items assessed on a 7-point Likert-type rating scale ($\alpha=.95$). The VHS was
9
10 developed by the World Health Organization Sage Working Group on Vaccine Hesitancy,³⁰ and
11
12 psychometrically validated by our research group.¹⁵ The VHS was found to have two underlying
13
14 factors (i.e. 'lack of confidence', $\alpha=.92$; and 'risks', $\alpha=.64$) and items are assessed on a 5-point
15
16 Likert-type rating scale (Appendix 1).
17

18 19 **Analysis**

20
21
22 This study reports parents' HPV vaccine decision-making in percentages based on the six
23
24 PAPM stages. For assessing significant differences in PAPM stage based on child's gender, a
25
26 chi-square test was used.
27

28 To examine differences between reported vaccine knowledge and attitudes by PAPM
29
30 stage, one-way ANOVA and Tukey Honest Significant Difference (HSD) post-hoc tests were
31
32 conducted.

33
34 Multinomial logistic regression analyses were used to calculate the odds ratios of being in
35
36 one of the first five PAPM stages compared to PAPM stage 6 (i.e. vaccinated, reference
37
38 category). PAPM stage was the dependent variable. First, we conducted bivariate multinomial
39
40 logistic regression analyses and estimated the associations for each independent variable
41
42 individually. Subsequently, we performed multivariate multinomial logistic regression analyses
43
44 by including 14 independent variables in a single model. In order to select variables that would
45
46 ensure the most parsimonious and theory-driven multivariate model, variables were included
47
48 based on attitudes predicted to be associated with behavioural change (according to the HBM)
49
50 and significant modifying factors in the literature (see Figure 1).^{16,31} Odds ratios and 95%
51
52 confidence intervals (CI) were calculated. To assess multicollinearity of the multivariate
53
54 multinomial logistic regression models, the Variation Inflation Factor (VIF) was calculated for
55
56 all predictors. Model fit diagnostics were reported based on following criteria: (a) Cox-Snell R^2 ,
57
58 (b) Cragg-Uhler R^2 , and (c) McFadden R^2 .

59
60 Statistical analyses were performed using IBM SPSS V.23 and R 3.3.2.

Results

Sample Demographics

A total of 4,606 parents completed the survey. The response rate, calculated based on completion by participants who initiated the questionnaire ($N=6,789$) was 67.9%. Overall, 827 (18.0%) participants were excluded as these participants were detected to be inattentive or unmotivated respondents based on data cleaning (i.e. the use of two bogus items and index of psychometric synonyms).²⁰ Sociodemographic characteristics for the final sample ($N=3,779$) are presented in Table 1. At the time of data collection, only 7% were parents of boys living in provinces where there was a publicly funded HPV vaccine program for boys available ($n = 252$) and fewer still would have been eligible for the program depending upon their child's age.

Identifying Canadian parents' stage of decision-making by child's gender

Table 2 shows the numbers of parents of boys and parents of girls across the six PAPM stages. HPV vaccine uptake of Canadian children was low, with only 801 (41.0%) parents of girls reporting that their daughters were vaccinated and only 160 (8.8%) parents of boys reporting that their sons were vaccinated. There was a significant and large association between child's gender and PAPM stage of decision making ($\chi^2(5)=735.25, p<.001, \phi_c=.44$) with parents of boys more likely to be in earlier stages.³²

Comparison of knowledge and attitudes for HPV vaccine PAPM stages

One-way ANOVA found that there was a significant effect of PAPM stage on all vaccine knowledge and attitude scales (Table 3). The greatest effect sizes were for cues to action, benefits, and barriers-harms.

Post hoc analyses found that knowledge (both HPV vaccine knowledge and HPV general knowledge) was significantly lower for PAPM stages 1-3 (unaware, unengaged, and undecided) compared to later PAPM stages. In addition, parents who were unaware reported significantly

1
2
3
4 lower perceived vaccine affordability compared to parents who were unengaged and undecided,
5 while parents who were unaware and unengaged reported significantly lower perceived harms
6 and VHS-risk compared to parents who were undecided.
7
8

9
10 Parents who decided not to vaccinate (Stage 4) significantly differed from all other PAPM
11 stages on all scales except barriers-not accessible, self-efficacy, and knowledge scales (VK and
12 GK); however, on these four scales, parents who decided not to vaccinate responded similarly to
13 parents who decided to vaccinate or already vaccinated their child (Stages 5 and 6).
14
15

16
17 Parents who decided to vaccinate their child (Stage 5) reported significantly higher perceived
18 benefits, greater perceived barriers of accessibility and affordability, and fewer cues to action
19 compared to those who already vaccinated their child (Stage 6).
20
21
22

23 24 **Examination of correlates of PAPM stage**

25
26
27
28 The bivariate and multivariate multinomial logistic regression analyses of parents' PAPM
29 stage can be found in Tables 4 and 5, respectively. Appendix 2 contains exploratory analyses of
30 additional variables as well as all analyses conducted separately for parents of boys and girls. All
31 earlier stages of PAPM were compared to the reference group (Stage 6-Vaccinated).
32
33
34

35 36 37 ***Bivariate multinomial logistic regression***

38
39 Parents of daughters, older children, and parents with a HCP recommendation had
40 decreased odds of being in any earlier PAPM stage as compared to the last PAPM stage (i.e.
41 vaccinated). Higher HPV vaccine knowledge was significantly associated with decreased odds of
42 being unaware, unengaged, undecided, or decided to vaccinate.
43
44
45

46
47 Parents who had decided not to vaccinate their child had significantly stronger vaccine
48 conspiracy beliefs (OR=3.10; 95% CI 2.78;3.46), lack of confidence in vaccines (OR=9.21; 95%
49 CI 7.50;11.31), and higher perception of vaccine risks (OR=5.19; 95% CI 4.38;6.16) compared
50 to parents who vaccinated their child.
51
52

53
54 Parents living in provinces with HPV vaccination programs for boys had significantly
55 lower odds of being unaware (OR=0.34; 95% CI 0.24;0.48), unengaged (OR=0.49; 95% CI
56 0.34;0.70), or undecided (OR=0.50; 95% CI 0.37;0.67). Further analysis by child's gender
57 indicated that this effect was not significant in the model examining parents of girls; however,
58
59
60

1
2
3
4 parents of boys living in provinces with HPV vaccine funding for boys had significantly lower
5 odds of being in any earlier PAPM stage compared to vaccinated (Tables B4 and B7).
6
7

8 9 ***Multivariate multinomial logistic regression***

10
11 Parents of daughters and older children had significantly decreased odds of reporting that
12 their child was in any earlier PAPM stage compared to vaccinated . Parents answering the survey
13 in French had lower odds of being unengaged (OR=0.67; 95% CI 0.47;0.97), undecided
14 (OR=0.61; 95% CI 0.44;0.84) and decided to vaccinate (OR=0.51; 95% CI 0.38;0.67).
15
16

17
18 Parents who received a HCP recommendation for HPV vaccination had lower odds of
19 being unaware (OR=0.04; 95% CI 0.01;0.16), unengaged (OR=0.21; 95% CI 0.10;0.43),
20 undecided (OR=0.55; 95% CI 0.36;0.84), and decided not to vaccinate (OR=0.30; 95% CI
21 0.15;0.61). There was no significant difference between the decided to vaccinate and vaccinated
22 groups. Higher HPV vaccine knowledge was significantly associated with decreased odds of
23 being unaware (OR=0.75; 95% CI 0.71;0.79), unengaged (OR=0.86; 95% CI 0.81;0.91), and
24 increased odds of having decided not to vaccinate (OR=1.12; 95% CI 1.04;1.22).
25
26

27
28 A higher perception of susceptibility and severity were only significantly associated with
29 decreased odds of being in the stage decided not to vaccinate (OR=0.68; 95% CI 0.55;0.84 and
30 OR=0.66; 95% CI 0.54;0.80). Higher perception of the benefits of vaccination was significantly
31 associated with increased odds of being unaware (OR=1.89; 95% CI 1.48;2.41), unengaged
32 (OR=1.62; 95% CI 1.26;2.08), undecided (OR=1.40; 95% CI 1.12;1.74) and decided to vaccinate
33 (OR=2.10; 95% CI 1.72;2.55), and decreased odds of being decided not to vaccinate (OR=0.60;
34 95% CI 0.45;0.81).
35
36

37
38 Compared to those who vaccinated their child, parents in all other stages had significantly
39 increased odds of reporting greater barriers (including affordability, accessibility, and perceived
40 harms). However, perceived accessibility of parents who vaccinated their children did not differ
41 significantly with parents who were undecided, decided not to vaccinate their child, or decided to
42 vaccinate their child. Of note, parents who had a higher score on perceived harms had higher
43 odds of being in Stage 4 (decided not to vaccinate) (OR=3.50; 95% CI 2.85;4.28). Greater
44 perceived influence of others (cues to action) was associated with lower odds of being in any of
45 the earlier stages (compared to vaccinated) (OR range of 0.20 to 0.51).
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

This study examined six distinct stages of HPV vaccine decision-making using the PAPM framework in a national survey of Canadian parents of 9-16 year-old boys and girls. Only 41.0% of girls and 8.8% of boys were in the final PAPM stage (Stage 6-Vaccinated). This is a lower proportion of vaccinated children than reported by other Canadian studies,^{3,33} which may be due to this study's design, which included data from jurisdictions without male HPV vaccination programs during data collection, relied on parental report of vaccination status, and evaluated a larger age range of children (including children before they were offered the HPV vaccine in funded school-based programs).

In a 2014 study using the PAPM to examine Canadian parents of boys, the majority of parents were unaware of HPV vaccination for their sons (Stage 1, 57.0%), while exceptionally few had decided to vaccinate their son (Stage 5, 5.0%) or had sons who had received the HPV vaccine (Stage 6, 1.1%).¹⁶ Data from the present study indicates that two years later and with two additional Canadian jurisdictions with male HPV vaccine programs (Alberta and Nova Scotia), fewer parents were unaware (25.6%), and more parents had decided to vaccinate (19.1%) or already vaccinated (8.8%) their son (Table 2). By comparison, a survey of parents of boys conducted around the same time in the UK (when there was no funded program for boys) found that 46.8% were unaware.³⁴ This emphasizes the importance of publicly funded vaccine programs and the associated educational campaigns in increasing awareness and uptake.³⁵⁻³⁷

Two further studies have used the PAPM to evaluate HPV vaccine decision-making.^{26,38} Though these studies were conducted in college students, both found males to be overwhelmingly unaware or unengaged (90% or 85.7%, respectively),^{26,38} and Barnard et al. also found females to be predominantly in these stages (62.9%).³⁸

The present study was unique in evaluating PAPM stages for both parents of girls and boys. The literature has primarily focused on parents of girls,^{35,37-44} with none using the PAPM. We found a significant association between child's gender and PAPM stage of decision-making, with parents of boys more likely to be in earlier PAPM stages, a finding similar to results from other studies.^{35,37,39-41}

Multinomial analyses highlighted some important correlates across all PAPM stages as well as some correlates that are particularly important for specific stages. Overall, this study

1
2
3
4 found that parents of daughters (compared to sons), of older children, and parents with a HCP
5 recommendation had decreased odds of being in any earlier PAPM stage as compared to the last
6 PAPM stage (i.e. vaccinated). The importance of a HCP recommendation in making the decision
7 regarding HPV vaccination is a well-established finding.^{11,12,16,35,37,41,45} This study contributes to
8 the literature by highlighting that a HCP recommendation is a significant and important
9 differential factor between parents who are ‘hesitant’ (unengaged, undecided, and decided not;
10 Stages 2-4) and ‘acceptors’ (decided to vaccinate and vaccinated; Stages 5-6), but not a
11 significant differential factors between acceptor groups (Stages 5 and 6). This suggests that while
12 a HCP recommendation may increase the likelihood that a parent accepts HPV vaccination; a
13 HCP recommendation alone may not be sufficient to move parents from deciding to vaccinate
14 their child (Stage 5) to having vaccinated their child (Stage 6).

15
16
17
18
19
20
21
22
23
24 The relationship between knowledge and uptake has previously yielded mixed results (as
25 high and low knowledge have been associated with vaccination).^{12,46} Application of the PAPM
26 model shows that low HPV vaccine knowledge is an important correlate of early stages of
27 decision-making (as compared to vaccinated). Interestingly, in the multivariate analysis parents
28 who had decided not to vaccinate their child had significantly higher HPV vaccine knowledge
29 than parents who vaccinated their child. Accordingly, education interventions alone may not be
30 sufficient for HPV vaccination.

31
32
33
34
35
36
37 In line with previous research, this study found perceived benefits, barriers-harms, and
38 cues to action were key correlates of PAPM stage.¹² Previous research using a binary outcome
39 has reported mixed findings regarding the relationship between HPV vaccination with perceived
40 severity and susceptibility of HPV infection and associated disease.^{12,31} By using a nuanced
41 framework of decision-making, this study highlights that, when taking all other variables into
42 account, susceptibility and severity were not significant correlates of earlier PAPM stages except
43 for the ‘decided not to vaccinate’ stage (Stage 4). Future research should use these findings to
44 investigate theoretically informed interventions to specifically target subsets of the population
45 (by child’s gender and decision-making stage), with particular attention towards addressing
46 knowledge gaps, perceived barriers, and concerns of parents.

47
48
49
50
51
52
53
54
55
56
57
58
59
60
Interestingly, parents of boys (but not parents of girls) living in provinces with HPV
vaccination programs for boys had significantly lower odds of being in early PAPM stages,
emphasizing the importance of publicly funded HPV vaccine programs for boys.

Study strengths and limitations

This study is unique in examining and comparing HPV vaccine decision-making in Canadian parents of boys and girls using two well-established theoretical frameworks (i.e. the HBM and the PAPM), which captures decision-making in a nuanced and precise way. This is the first study to use the PAPM to evaluate HPV vaccine decision-making in parents of girls and boys. Strengths of the study also include a large sample size, data cleaning techniques that eliminated careless responders, and a nationally representative sample.²⁰ This study's questionnaire benefited from the use of intelligent programming to personalize survey items, administration in either English or French, avoiding any missing data, the use of psychometrically validated scales, and the randomization of items within scales to reduce the possibility of an order effect.²⁰


This study is not without limitations. The cross-sectional design makes it impossible to determine causality. Although we had a reasonable response rate (67.9%), there remains the potential that there were similarities between non-responders that could influence the representativeness of the sample. Furthermore, although the sample was generally representative of the Canadian population, it was slightly wealthier, more educated, and White (as compared to the 2016 Census).⁴⁷ This study was also not able to recruit many participants living in Canada's territories, due to constraints in Leger's panel. Future research is needed to replicate this study's conceptual framework and findings to specifically investigate HPV vaccination in disadvantaged populations, as well as in other countries or in cross-country comparisons.

It is also possible that the measurement of HCP recommendation in this study overlooked parents who were recommended the vaccine by a HCP but had not seen or discussed the HPV vaccine with a HCP (e.g. via a letter that was sent home). Moreover, the scope of variables assessed is limited, there are other familial, sociological, environmental, and communication factors that were not included. Specifically, future research should consider the impact of having an older child who was eligible for, or received, the HPV vaccine. Future research should also further examine the impact of publicly funded HPV vaccination programs on parents' decision-making.

Conclusions

HPV vaccination remains low in Canada. Using a stage-based model of decision-making, this study found that only a quarter of parents were in the final PAPM stage. Parents of daughters, older children, and those with a HCP recommendation had decreased odds of being in any earlier PAPM stage. Individual health beliefs as well as cues to action were key correlates of PAPM stage overall; however, the combinations and importance of correlates varied by PAPM stage and child's gender. These findings indicate that it may be important for future interventions to target and tailor health messaging for different groups depending on their stage of decision-making.

Figure 1. An integrated conceptual framework of HPV vaccination



Note. The PAPM, as applied to HPV vaccination, identifies individuals along six stages of decision-making: 1) *unaware* of the vaccine (“I was unaware that the HPV vaccine could be given to CHILD*”); 2) *unengaged* in the decision to vaccinate their child (“I have never thought about vaccinating CHILD* against HPV”); 3) *undecided* about whether to vaccinate their child (“I am undecided about vaccinating CHILD* against HPV”); 4) *decided not to act* (i.e. decided not to vaccinate their child, “I have decided I DO NOT want to vaccinate CHILD* against HPV”); 5) *decided to act* (i.e. decided to vaccinate their child, “I have decided I DO want to vaccinate CHILD* against HPV”); and 6) *acted* (i.e. vaccinated their child, “CHILD* has already received the HPV vaccine”). *To increase the personalization of this questionnaire, intelligent programming allowed for each question with “CHILD” to specifically include their child’s name.

Table 1. Sample Characteristic ($N = 3,779$)

Participant Characteristics	N (%)
<i>Parent's gender</i>	
Men	1,311 (34.69)
Women	2,468 (65.31)
<i>Parent's age</i>	
Range (years)	18-81
Mean (SD)	43.51 (6.86)
<i>Language in which parents answered the survey</i>	
English	2,801 (74.12)
French	978 (25.88)
<i>Marital status</i>	
Single/Separated/divorced/widowed	760 (20.11)
Married/common law	3,019 (79.89)
<i>Parent's level of education</i>	
Elementary or high school	659 (17.44)
Trade technical or university	3,120 (82.56)
<i>Parent's employment status</i>	
Employed	3,057 (80.89)
Not employed	722 (19.11)
<i>Born in Canada</i>	
Yes	3,214 (85.05)
No	565 (14.95)
<i>Parent's ethnicity</i>	
White	3,224 (85.31)
Other	555 (14.69)
<i>Parent's religion</i>	
Any religious affiliation	2,493 (65.97)
No religious affiliation	1,286 (34.03)
<i>Household income</i>	
<100 K	1,973 (52.21)
≥ 100 K	1,409 (37.28)
Prefer not to answer	397 (10.51)
<i>Child's gender</i>	
Boys (sons)	1,826 (48.32)
Girls (daughters)	1,953 (51.68)
<i>Child's age</i>	
Range (years)	9-16
Mean (SD)	12.58 (2.31)
<i>Child's school's religion</i>	
No affiliation	2,821 (74.65)
Any religious affiliation	958 (25.35)
<i>Child's school's language</i>	
English	2,647 (70.04)
French and Other	1,132 (29.96)

Province with funded HPV vaccination for boys

No	3,225 (85.34)
Yes	554 (14.66)

Size of city

<100K	1,799 (47.61)
≥ 100K	1,980 (52.39)

Number of children in the family

1	868 (22.97)
2	1,747 (46.23)
≥3	1,164 (30.80)

Child's sexual orientation

Heterosexual	3,301 (87.35)
Other	478 (12.65)

HCP recommendation

No	3,346 (88.54)
Yes	433 (11.46)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. Parents' PAPM stage by child's gender

<i>PAPM Stage</i>	Parents of girls (<i>n</i> = 1,953)	Parents of boys (<i>n</i> = 1,826)	Test of proportions girls versus boys	All parents (<i>N</i> = 3,779)
	<i>n</i> (%)	<i>n</i> (%)	95% CI	<i>N</i> (%)
<i>Stage 1- Unaware</i>	136 (7.0)	468 (25.6)	-20.96; -16.37	604 (16.0)
<i>Stage 2- Unengaged</i>	97 (5.0)	298 (16.3)	-13.30; -9.40	395 (10.5)
<i>Stage 3- Undecided</i>	291 (14.9)	389 (21.3)	-8.86; -3.94	680 (18.0)
<i>Stage 4- Decided NOT</i>	191 (9.8)	162 (8.9)	-0.95; 2.76	353 (9.3)
<i>Stage 5- Decided YES</i>	437 (22.4)	349 (19.1)	0.68; 5.85	786 (20.8)
<i>Stage 6- Vaccinated</i>	801 (41.0)	160 (8.8)	29.71; 34.79	961 (25.4)

Table 3. Comparison of parents' vaccine knowledge and attitudes by PAPM stage

Scale	Total <i>N</i> = 3,779 <i>M</i> (<i>SD</i>)	Stage 1- Unaware <i>n</i> = 604 <i>M</i> (<i>SD</i>)	Stage 2- Unengaged <i>n</i> = 395 <i>M</i> (<i>SD</i>)	Stage 3- Undecided <i>n</i> = 680 <i>M</i> (<i>SD</i>)	Stage 4- Decided NOT <i>n</i> = 353 <i>M</i> (<i>SD</i>)	Stage 5- Decided YES <i>n</i> = 786 <i>M</i> (<i>SD</i>)	Stage 6- Vaccinated <i>n</i> = 961 <i>M</i> (<i>SD</i>)	ANOVA <i>F</i> test- statistic	Effect size (ω)
<i>HPV Vaccine Knowledge</i>									
HPV General Knowledge	12.79 (6.07)	8.53 (6.51) ^a	11.34 (5.97) ^b	12.95 (5.77) ^c	14.88 (5.45) ^d	14.33 (5.39) ^d	13.93 (5.32) ^d	101.32	0.34
HPV Vaccine Knowledge	6.01 (2.92)	3.76 (3.07) ^a	4.85 (2.97) ^b	5.89 (2.75) ^c	6.88 (2.55) ^d	6.74 (2.48) ^d	7.06 (2.37) ^d	149.95	0.41
<i>HPV-specific HBM attitudes</i>									
Susceptibility	4.93 (1.38)	4.73 (1.16) ^a	4.61 (1.16) ^{a,b}	4.50 (1.08) ^b	3.00 (1.37) ^c	5.65 (1.09) ^d	5.64 (1.05) ^d	382.00	0.58
Severity	5.91 (1.08)	5.84 (1.08) ^a	5.82 (1.05) ^a	5.83 (1.06) ^a	5.18 (1.39) ^b	6.15 (0.94) ^c	6.13 (0.96) ^c	53.11	0.25
Benefits	4.90 (1.14)	4.87 (0.95) ^a	4.75 (0.93) ^{a,b}	4.58 (0.86) ^b	3.10 (1.06) ^c	5.57 (0.86) ^d	5.33 (0.91) ^e	421.15	0.60
Barriers-not affordable	3.61 (1.70)	4.70 (1.33) ^a	4.35 (1.37) ^b	4.16 (1.52) ^b	3.09 (1.56) ^c	3.78 (1.64) ^d	2.30 (1.31) ^e	275.65	0.52
Barriers-not accessible	2.85 (1.15)	3.57 (0.92) ^a	3.35 (0.98) ^{a,b}	3.18 (1.02) ^b	2.86 (1.16) ^c	2.65 (1.12) ^c	2.15 (0.99) ^d	188.42	0.45
Barriers-harms	3.54 (1.42)	3.74 (1.07) ^a	3.82 (1.10) ^a	4.18 (1.12) ^b	5.50 (1.21) ^c	2.85 (1.17) ^d	2.68 (1.12) ^d	433.92	0.60
Cues to action	4.62 (1.16)	4.14 (0.88) ^a	3.98 (0.88) ^a	4.06 (0.79) ^a	3.43 (1.02) ^b	5.11 (0.94) ^c	5.61 (0.85) ^d	554.38	0.65
Self efficacy	6.00 (1.01)	5.63 (1.10) ^a	5.65 (1.09) ^a	5.67 (1.03) ^a	6.29 (1.00) ^b	6.27 (0.82) ^b	6.28 (0.83) ^b	79.45	0.31
<i>General Vaccine Attitudes</i>									
VCBS	3.23 (1.44)	3.46 (1.36) ^a	3.40 (1.38) ^a	3.56 (1.35) ^a	4.70 (1.35) ^b	2.66 (1.31) ^c	2.73 (1.21) ^c	160.04	0.42
VHS-Lack of Confidence	1.98 (0.72)	2.04 (0.66) ^a	2.07 (0.64) ^a	2.13 (0.65) ^a	2.84 (0.91) ^b	1.69 (0.57) ^c	1.74 (0.54) ^c	196.69	0.45
VHS-Risks	3.07 (0.95)	3.15 (0.85) ^a	3.14 (0.86) ^a	3.38 (0.87) ^b	3.90 (0.83) ^c	2.75 (0.94) ^d	2.74 (0.85) ^d	132.25	0.38

Note. All skewness and kurtosis are less than 2. VHS scales are measured 1-5; all other scales are measured 1-7. All one-way independent groups ANOVA were significant (all $p < .001$). Effect sizes (ω) are presented for each ANOVA analysis. Post hoc tests were conducted using Tukey HSD. For each scale, groups that were not significantly different ($p < .01$) from each other in post hoc tests are in the same group (notated using a superscript, e.g. ^a).

Table 4. Bivariate multinomial logistic regression analysis of parents' PAPM stage ($N = 3,779$)

Variables	Stage 1-Unaware OR (95% CI) $n = 604$	Stage 2-Unengaged OR (95% CI) $n = 395$	Stage 3-Undecided OR (95% CI) $n = 680$	Stage 4-Decided NOT OR (95% CI) $n = 353$	Stage 5-Decided YES OR (95% CI) $n = 786$
<i>Child's gender</i>					
Male	(reference)	(reference)	(reference)	(reference)	(reference)
Female	0.06*** (0.04; 0.07)	0.07*** (0.05; 0.09)	0.15*** (0.12; 0.19)	0.24*** (0.18; 0.31)	0.25*** (0.20; 0.31)
<i>Child's age (One-year increase)</i>	0.73*** (0.69; 0.76)	0.80*** (0.76; 0.84)	0.76*** (0.73; 0.79)	0.81*** (0.76; 0.85)	0.71*** (0.68; 0.75)
<i>Language parents answered the survey</i>					
English	(reference)	(reference)	(reference)	(reference)	(reference)
French	1.26 (1.02; 1.57)	0.65* (0.49; 0.85)	0.56*** (0.45; 0.71)	0.87 (0.67; 1.14)	0.48*** (0.38; 0.60)
<i>Parent's ethnicity</i>					
Other	(reference)	(reference)	(reference)	(reference)	(reference)
White	0.64* (0.48; 0.86)	0.61* (0.44; 0.85)	0.75 (0.56; 0.99)	0.90 (0.62; 1.31)	0.71 (0.54; 0.94)
<i>HCP recommendation</i>					
No	(reference)	(reference)	(reference)	(reference)	(reference)
Yes	0.01*** (0.01; 0.05)	0.09*** (0.04; 0.17)	0.32*** (0.24; 0.45)	0.43*** (0.29; 0.62)	0.74 (0.58; 0.95)
<i>HPV Vaccine knowledge (One-unit increase)</i>	0.66*** (0.64; 0.69)	0.74*** (0.71; 0.78)	0.84*** (0.81; 0.87)	0.97 (0.92; 1.02)	0.95* (0.91; 0.98)
<i>Susceptibility (One-unit increase)</i>	0.47*** (0.43; 0.52)	0.43*** (0.39; 0.48)	0.40*** (0.36; 0.44)	0.16*** (0.14; 0.18)	1.01 (0.92; 1.10)
<i>Severity (One-unit increase)</i>	0.75*** (0.68; 0.83)	0.74*** (0.66; 0.82)	0.74*** (0.67; 0.82)	0.48*** (0.43; 0.54)	1.03 (0.93; 1.14)
<i>Benefits (One-unit increase)</i>	0.57*** (0.50; 0.63)	0.49*** (0.42; 0.56)	0.39*** (0.35; 0.44)	0.08*** (0.07; 0.1)	1.36*** (1.22; 1.51)
<i>Barriers-Affordability (One-unit increase)</i>	3.03*** (2.78; 3.30)	2.59*** (2.37; 2.83)	2.38*** (2.20; 2.57)	1.49*** (1.37; 1.63)	2.02*** (1.88; 2.17)
<i>Barriers-Accessibility (One-unit increase)</i>	3.74*** (3.34; 4.20)	3.02*** (2.67; 3.42)	2.60*** (2.34; 2.88)	1.95*** (1.73; 2.19)	1.60*** (1.46; 1.76)
<i>Barriers-Harms (One-unit increase)</i>	2.22*** (2.02; 2.44)	2.36*** (2.11; 2.63)	3.12*** (2.82; 3.45)	8.85*** (7.60; 10.31)	1.14* (1.05; 1.24)
<i>Cues to action (One-unit increase)</i>	0.15*** (0.13; 0.17)	0.12*** (0.10; 0.14)	0.13*** (0.11; 0.15)	0.06*** (0.05; 0.07)	0.55*** (0.50; 0.62)
<i>Self-Efficacy (One-unit increase)</i>	0.50*** (0.45; 0.56)	0.51*** (0.45; 0.57)	0.51*** (0.46; 0.57)	1.02 (0.88; 1.19)	0.99 (0.88; 1.11)
<i>Vaccine conspiracy beliefs (One-unit increase)</i>	1.51*** (1.40; 1.64)	1.47*** (1.34; 1.60)	1.60*** (1.48; 1.73)	3.10*** (2.78; 3.46)	0.96 (0.89; 1.03)
<i>Hesitancy-Lack of Confidence (One-unit increase)</i>	2.34* (1.97; 2.79)	2.49*** (2.05; 3.03)	2.86*** (2.42; 3.38)	9.21*** (7.50; 11.31)	0.86 (0.72; 1.02)
<i>Hesitancy-Risks (One-unit increase)</i>	1.69*** (1.50; 1.90)	1.65*** (1.44; 1.89)	2.31*** (2.05; 2.61)	5.19*** (4.38; 6.16)	1.01 (0.91; 1.12)

Note. The reference category for PAPM stage is 'Stage 6-Vaccinated' ($n = 961$). Bold integers indicates significant OR at $p < .05$. * $p < .01$, ** $p < .001$, *** $p < .0001$. (reference) = reference category. OR = odds ratio. 95% CI = 95% confidence interval. HCP = health care provider.

Table 5. Multivariate multinomial logistic regression analysis of parents' PAPM stage ($N = 3,779$)

Variables	Stage 1-Unaware AOR (95% CI) $n = 604$	Stage 2-Unengaged AOR (95% CI) $n = 395$	Stage 3-Undecided AOR (95% CI) $n = 680$	Stage 4-Decided NOT AOR (95% CI) $n = 353$	Stage 5-Decided YES AOR (95% CI) $n = 786$
<i>Child's gender</i>					
Male	(reference)	(reference)	(reference)	(reference)	(reference)
Female	0.15*** (0.11; 0.21)	0.15*** (0.11; 0.22)	0.33*** (0.24; 0.45)	0.36*** (0.23; 0.57)	0.49*** (0.37; 0.65)
<i>Child's age</i> (One-year increase)	0.63*** (0.59; 0.67)	0.69*** (0.65; 0.74)	0.68*** (0.64; 0.73)	0.80*** (0.74; 0.88)	0.64*** (0.60; 0.67)
<i>Language parents answered the survey</i>					
English	(reference)	(reference)	(reference)	(reference)	(reference)
French	1.21 (0.87; 1.69)	0.67 (0.47; 0.97)	0.61* (0.44; 0.84)	0.91 (0.58; 1.43)	0.51*** (0.38; 0.67)
<i>Parent's ethnicity</i>					
Other	(reference)	(reference)	(reference)	(reference)	(reference)
White	1.37 (0.91; 2.06)	1.28 (0.84; 1.96)	1.62 (1.10; 2.39)	1.16 (0.65; 2.05)	1.31 (0.93; 1.84)
<i>HCP recommendation</i>					
No	(reference)	(reference)	(reference)	(reference)	(reference)
Yes	0.04*** (0.01; 0.16)	0.21*** (0.10; 0.43)	0.55* (0.36; 0.84)	0.30** (0.15; 0.61)	0.89 (0.66; 1.20)
<i>HPV Vaccine knowledge</i> (One-unit increase)	0.75*** (0.71; 0.79)	0.86*** (0.81; 0.91)	0.98 (0.93; 1.04)	1.12* (1.04; 1.22)	0.96 (0.91; 1.01)
<i>Susceptibility</i> (One-unit increase)	0.97 (0.81; 1.17)	0.98 (0.81; 1.18)	0.89 (0.75; 1.05)	0.68** (0.55; 0.84)	1.06 (0.91; 1.23)
<i>Severity</i> (One-unit increase)	1.06 (0.91; 1.25)	1.06 (0.90; 1.25)	1.08 (0.93; 1.25)	0.66*** (0.54; 0.80)	1.02 (0.89; 1.16)
<i>Benefits</i> (One-unit increase)	1.89*** (1.48; 2.41)	1.62** (1.26; 2.08)	1.40* (1.12; 1.74)	0.60** (0.45; 0.81)	2.10*** (1.72; 2.55)
<i>Barriers-Affordability</i> (One-unit increase)	1.99*** (1.76; 2.24)	1.62*** (1.43; 1.84)	1.70*** (1.52; 1.90)	1.22 (1.03; 1.43)	1.68*** (1.53; 1.84)
<i>Barriers-Accessibility</i> (One-unit increase)	1.50*** (1.27; 1.78)	1.26 (1.05; 1.50)	1.10 (0.94; 1.29)	1.07 (0.85; 1.34)	1.14 (0.99; 1.31)
<i>Barriers-Harms</i> (One-unit increase)	1.53*** (1.31; 1.78)	1.68*** (1.43; 1.97)	2.19*** (1.90; 2.52)	3.50*** (2.85; 4.28)	1.15 (1.01; 1.30)
<i>Cues to action</i> (One-unit increase)	0.27*** (0.22; 0.34)	0.20*** (0.16; 0.25)	0.24*** (0.19; 0.29)	0.21*** (0.16; 0.28)	0.51*** (0.43; 0.61)
<i>Self-Efficacy</i> (One-unit increase)	1.10 (0.91; 1.33)	1.16 (0.96; 1.40)	1.11 (0.93; 1.32)	1.87*** (1.49; 2.35)	1.24 (1.04; 1.48)

Note. The reference category for PAPM stage is 'Stage 6-Vaccinated' ($n = 961$). Bold integers indicates significant AOR at $p < .05$. * $p < .01$, ** $p < .001$, *** $p < .0001$. Cox-Snell $R^2 = 0.72$. Cragg Uhler $R^2 = 0.74$. McFadden $R^2 = 0.37$. (reference) = reference category. AOR = adjusted odds ratio. 95% CI = 95% confidence interval. HCP = health care provider.

References

1. Brotherton JML, Zuber F, Bloem PJN. Primary Prevention of HPV through Vaccination: Update on Current Global Status. *Curr Obst Gynecol Reports*. 2016;5:210-224.
2. Canadian Cancer Society. *Canadian Cancer Statistics 2016*. 2016.
3. Shapiro GK, Guichon J, Kelaher M. Canadian school-based HPV vaccine programs and policy considerations. *Vaccine*. 2017; 35:5700-5707.
4. Blake DR, Middleman AB. Human Papillomavirus Vaccine Update. *Pediatric clinics of North America*. Apr 2017;64(2):321-329.
5. Brotherton JML, Bloem PJN. HPV Vaccination: Current Global Status. *Curr Obstet Gynecol Rep*. 2015;4:220-233.
6. Cervical Cancer Action. Global Progress in HPV Vaccination. 2017; <http://www.cervicalcanceraction.org/comments/comments3.php>. Accessed July 20, 2018.
7. Shapiro GK, Perez S, Rosberger Z. Including males in Canadian human papillomavirus vaccination programs: a policy analysis. *Canadian Medical Association Journal*. 2016;188(12):881-886.
8. Shapiro GK, Guichon J, Prue G, Perez S, Rosberger Z. A Multiple Streams analysis of the decisions to fund gender-neutral HPV vaccination in Canada. *Prev Med*. Apr 20 2017;100:123-131.
9. Public Health Agency of Canada. Canada's Provincial and Territorial Routine (and Catch-up) Vaccination Programs for Infants and Children. In: Canadian Immunization Guide, ed2017.
10. Shapiro G, Guichon J, Perez S, Rosberger Z. British Columbia's flawed HPV vaccination policy. *Policy Options* [Internet]. 2015 Sep 25; <http://policyoptions.irpp.org/issues/september-2015/british-columbias-flawed-hpv-vaccination-policy/>.
11. Holman DM, Benard V, Roland KB, Watson M, Liddon N, Stokley S. Barriers to Human Papillomavirus Vaccination Among US Adolescents: A Systematic Review of the Literature. *JAMA pediatrics*. 2014; 168 (1):76-82.
12. Radisic G, Chapman J, Flight I, Wilson C. Factors associated with parents' attitudes to the HPV vaccination of their adolescent sons : A systematic review. *Prev Med*. Feb 2017;95:26-37.
13. Gerend MA, Shepherd JE. Predicting human papillomavirus vaccine uptake in young adult women: comparing the health belief model and theory of planned behavior. *Annals of behavioral medicine : a publication of the Society of Behavioral Medicine*. Oct 2012;44(2):171-180.
14. Shapiro GK, Holding A, Perez S, Amsel R, Rosberger Z. Validation of the vaccine conspiracy beliefs scale. *Papillomavirus Research*. 12// 2016;2:167-172.
15. Shapiro GK, Tatar O, Dube E, et al. The Vaccine Hesitancy Scale: Psychometric properties and validation. *Vaccine*. 2018;36:545-552.
16. Perez S, Tatar O, Gilca V, et al. Untangling the psychosocial predictors of HPV vaccination decision-making among parents of boys. *Vaccine*. Aug 24 2017;35(36):4713-4721.
17. Perez S, Shapiro GK, Tatar O, Joyal-Desmarais K, Rosberger Z. Development and

- validation of the HPV Attitudes and Beliefs Scale (HABS) in a National Canadian sample. *Sexually Transmitted Diseases*. 2016;43(10):626-632.
18. Shapiro GK, Surian D, Dunn A, Perry R, Kelaheer M. HPV vaccine concern on Twitter: A cross-country analysis in Australia, Canada and the UK. *BMJ Open*. 2017;7:e016869.
 19. Dube E, Laberge C, Guay M, Bramadat P, Roy R, Bettinger J. Vaccine hesitancy: An overview. *Human Vaccines & Immunotherapeutics*. 2013;9(8):1763–1773.
 20. Shapiro GK, Perez S, Naz A, et al. Investigating Canadian parents' HPV vaccine knowledge, attitudes, and behaviour: A study protocol for a longitudinal national online survey. *BMJ Open*. 2017;7:e017814.
 21. Weinstein ND. The precaution adoption process. *Health Psychol*. 1988;7(4):355-386.
 22. Prue G, Santin O. HPV vaccine acceptance in male adolescents. *Psychooncology*. Aug 28 2015;24(10):1327-1329.
 23. Weinstein ND, Sandman PM. A model of the precaution adoption process: evidence from home radon testing. *Health Psychol*. 1992;11(3):170-180.
 24. Barnard M, George P, Perryman ML, Wolff LA. Human papillomavirus (HPV) vaccine knowledge, attitudes, and uptake in college students: Implications from the Precaution Adoption Process Model. *PloS one*. 2017;12(8):e0182266.
 25. Perez S, Tatar O, Shapiro GK, et al. Psychosocial determinants of parental human papillomavirus (HPV) vaccine decision-making for sons: Methodological challenges and initial results of a pan-Canadian longitudinal study. *BMC Public Health*. Dec 05 2016;16(1):1223.
 26. Tatar O, Perez S, Naz A, Shapiro GK, Rosberger Z. Psychosocial correlates of HPV vaccine acceptability in college males: A cross-sectional exploratory study. *Papillomavirus research (Amsterdam, Netherlands)*. Dec 2017;4:99-107.
 27. Leger. Marketing Research and Surveys. 2018; <http://leger360.com/en-ca/servicesurveyadvice>. Accessed April 24, 2018.
 28. Waller J, Ostini R, Marlow LA, McCaffery K, Zimet G. Validation of a measure of knowledge about human papillomavirus (HPV) using item response theory and classical test theory. *Preventive medicine*. 2013;56(1):35-40.
 29. Perez S, Tatar O, Ostini R, et al. Extending and validating a human papillomavirus (HPV) knowledge measure in a national sample of Canadian parents of boys. *Prev Med*. Oct 2016;91:43-49.
 30. Larson HJ, Jarretta C, Schulza WS, et al. Measuring vaccine hesitancy: The development of a survey tool. *Vaccine*. 2015;33(34):4165-4175.
 31. Krawczyk A, Knauper B, Gilca V, et al. Parents' decision-making about the human papillomavirus vaccine for their daughters: I. Quantitative results. *Human Vaccines & Immunotherapeutics*. 2015;11(2):322-329.
 32. Cohen J. Quantitative methods in psychology: A power primer. *Psychological Bulletin*. 1992;112(1):155-159.
 33. Gilbert NL, Gilmour H, Dubé E, Wilson SE, Laroche J. Estimates and determinants of HPV non-vaccination and vaccine refusal in girls 12 to 14 years of age in Canada: Results from the Childhood National Immunization Coverage Survey, 2013. *Human Vaccines & Immunotherapeutics*. 2016;12(6):1484-1490.
 34. Sherman SM, Nailor E. Attitudes towards and knowledge about Human

- Papillomavirus (HPV) and the HPV vaccination in parents of teenage boys in the UK. *PloS one*. 2018;13(4):e0195801.
35. Loke AY, Kwan ML, Wong YT, Wong AKY. The Uptake of Human Papillomavirus Vaccination and Its Associated Factors Among Adolescents: A Systematic Review. *Journal of primary care & community health*. Oct 2017;8(4):349-362.
36. Bird Y, Obidiya O, Mahmood R, Nwankwo C, Moraros J. Human Papillomavirus Vaccination Uptake in Canada: A Systematic Review and Meta-analysis. *International journal of preventive medicine*. 2017;8:71.
37. Johnson KL, Lin MY, Cabral H, Kazis LE, Katz IT. Variation in Human Papillomavirus Vaccine Uptake and Acceptability Between Female and Male Adolescents and Their Caregivers. *Journal of community health*. Jun 2017;42(3):522-532.
38. Barnard M. Human papillomavirus (HPV) vaccine knowledge, attitudes, and uptake in college students: Implications from the Precaution Adoption Process Model. *Hum Vaccin Immunother*. 2017;12(8):e0182266.
39. Finney Rutten LJ, St Sauver JL, Beebe TJ, et al. Association of both consistency and strength of self-reported clinician recommendation for HPV vaccination and HPV vaccine uptake among 11- to 12-year-old children. *Vaccine*. Oct 27 2017;35(45):6122-6128.
40. Hanson KE, Koch B, Bonner K, McRee AL, Basta NE. National Trends in Parental HPV Vaccination Intentions and Reasons for Hesitancy, 2010-2015. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. Mar 27 2018.
41. Lindley MC, Jeyarajah J, Yankey D, Curtis CR, Markowitz LE, Stokley S. Comparing human papillomavirus vaccine knowledge and intentions among parents of boys and girls. *Hum Vaccin Immunother*. Jun 02 2016;12(6):1519-1527.
42. Mohammed KA, Vivian E, Loux TM, Arnold LD. Factors associated with parents' intent to vaccinate adolescents for human papillomavirus: Findings from the 2014 National Immunization Survey-Teen. *Preventing chronic disease*. Jun 8 2017;14:E45.
43. VanWormer JJ, Bendixsen CG, Vickers ER, et al. Association between parent attitudes and receipt of human papillomavirus vaccine in adolescents. *BMC Public Health*. Oct 2 2017;17(1):766.
44. Wang Z, Wang J, Fang Y, et al. Parental acceptability of HPV vaccination for boys and girls aged 9-13 years in China: A population-based study. *Vaccine*. Mar 29 2018.
45. Gilkey MB, Moss JL, McRee AL, Brewer NT. Do correlates of HPV vaccine initiation differ between adolescent boys and girls? *Vaccine*. 2012;30:5928-5934.
46. Nickel B, Dodd RH, Turner RM, et al. Factors associated with the human papillomavirus (HPV) vaccination across three countries following vaccination introduction. *Preventive medicine reports*. Dec 2017;8:169-176.
47. Statistics Canada. Highlight tables, 2016 Census. In: Statistics Canada Catalogue no. 98-316-X2016001, ed. Ottawa2017.

***Conflict of Interest Form**

[Click here to download Conflict of Interest Form: CorrespondingAuthorDeclarationFormPM.pdf](#)

Appendices

Appendix A. Validated scales

Please find all the psychometrically validated scales that were used in this study in the tables below (Tables A1-A3). We hope that the inclusion of these scales in this appendix will facilitate replication, further validation, and exploration of these scales in future research.

Table A1. HPV and HPV vaccine knowledge

HPV General Knowledge Scale		True/ False/ Don't Know
Question		
1	HPV is very rare	F
2	HPV always has visible signs or symptoms	F
3	HPV can cause cervical cancer	T
4	HPV can be transmitted through genital skin-to-skin contact	T
5	There are many types of HPV	T
6	HPV can cause HIV/AIDS	F
7	HPV can be passed on during sexual intercourse	T
8	HPV can cause genital warts	T
9	Men cannot get HPV	F
10	Using condoms reduces the chances of HPV transmission	T
11	HPV can be cured with antibiotics	F
12	Having many sexual partners increases the risk of getting HPV	T
13	Most sexually active people will get HPV at some point in their lives	T
14	A person could have HPV for many years without knowing it	T
15	Having sex at an early age increases the risk of getting HPV	T
16	HPV can cause anal cancer	T
17	HPV is a bacterial infection	F
18	HPV can be transmitted through oral sex	T
19	HPV can cause cancer of the penis	T
20	HPV can be transmitted through anal sex	T
21	HPV infections always lead to health problems	F
22	HPV can cause oral cancer	T
23	A person with no symptoms cannot transmit the HPV infection	F
HPV Vaccine Knowledge Scale		

Question	True/ False/ Don't Know
1 The HPV vaccine requires at least 2 doses	T
2 The HPV vaccines offers protection against all sexually transmitted infections	F
3 The HPV vaccines are most effective if given to people who've never had sex	T
4 Someone who has had the HPV vaccine cannot develop cervical cancer	F
5 The HPV vaccines offer protection against most cervical cancers	T
6 The HPV vaccine offers protection against genital warts	T
7 Girls who have had an HPV vaccine do not need a Pap test when they are older	F
8 The HPV vaccine protects you from every type of HPV	F
9 You can cure HPV by getting the HPV vaccine	F
10 The HPV vaccine is approved and recommended by Health Canada for females aged 9-45 years	T
11 The HPV vaccine is approved and recommended by Health Canada for males aged 9-26 years	T

Source: Perez, S., Tatar, O., Ostini, R., Shapiro, G. K., Waller, J., Zimet, G., & Rosberger, Z. (2016). Extending and validating a human papillomavirus (HPV) knowledge measure in a national sample of Canadian parents of boys. *Preventive Medicine*, 91, 43-49. doi: 10.1016/j.ypmed.2016.07.017

Note. As recommended by the authors, two items from the original GK scale administration ('HPV usually doesn't need any treatment' and 'HPV can cause herpes') were deleted from this study's because the psychometric evaluation found that when removed the reliability of the scale improved. Further, in this study's administration of the VK Scale, 'vaccines' was changed to 'vaccine' to make the measure consistently in the singular. Slight adaptations were also made to ensure the items were gender-neutral (rather than directed at parents of males only) and updated based on policy recommendations and current generation vaccines.

Table A2. HPV-specific vaccine attitudes

Severity (HABS 'Perceived Threat')	
1	<i>...it would be serious if [CHILD] contracted HPV later in life</i>
2	<i>...it would be serious if [CHILD] contracted genital warts later in life</i>
3	<i>...it would be serious if [CHILD] contracted an HPV-related cancer later in life</i>
Susceptibility (HABS 'Risk')	
1	<i>...without the HPV vaccine, [CHILD] would be at risk of getting HPV later in life.</i>
2	<i>...without the HPV vaccine, [CHILD] would be at risk of getting genital warts later in life.</i>
3	<i>...without the HPV vaccine, [CHILD] would be at risk of getting an HPV-related cancer later in life.</i>
Benefits (HABS 'Benefits')	
1	<i>...the HPV vaccine has many benefits</i>
2	<i>...the HPV vaccine will protect [CHILD]'s sexual health</i>
3	<i>...the HPV vaccine works well</i>
4	<i>...the HPV vaccine is effective in preventing HPV</i>
5	<i>...the HPV vaccine is effective in preventing genital warts</i>
6	<i>...vaccinating [CHILD] against HPV may be a good thing to do for [CHILD]'s health</i> <i>...vaccinating [CHILD] against HPV was a good thing to do for [CHILD]'s health*</i>
7	<i>...vaccinating [CHILD] against HPV would give me peace of mind about [CHILD]'s sexual health</i> <i>...having vaccinated [CHILD] against HPV gives me peace of mind about [CHILD]'s sexual health*</i>
8	<i>...the HPV vaccine is effective in preventing HPV-related cancers</i>
9	<i>...vaccinating [CHILD] against HPV would protect [CHILD]'s current/future partner from getting infected with HPV</i> <i>...having vaccinated [CHILD] against HPV protects [CHILD]'s current/future partner from getting infected*</i>
10	<i>...getting [CHILD] the HPV vaccine would protect [CHILD]'s current/future partner against cancer</i> <i>...having gotten [CHILD] the HPV vaccine protects [CHILD]'s current/future partner against cancer*</i>
Barriers Accessibility (HABS 'Accessibility')	
1	<i>...it is hard to find a clinic that would be easy to access for getting the HPV vaccine for [CHILD] <input type="checkbox"/></i> <i>...it was hard to find a clinic that was easy to access for getting the HPV vaccine for [CHILD] * <input type="checkbox"/></i>
2	<i>...it is hard to find a provider or clinic where I would not have to wait a long time to get an appointment for [CHILD] to get vaccinated <input type="checkbox"/></i>
3	<i>...dealing with getting the HPV vaccine for [CHILD] would be simple</i> <i>...dealing with getting the HPV vaccine for [CHILD] was simple*</i>
4	<i>...the process of actually getting the HPV vaccine for [CHILD] would be easy</i> <i>...the process of actually getting the HPV vaccine for [CHILD] was easy*</i>
Barriers Affordability (HABS 'Affordability')	
1	<i>...the HPV vaccine is too expensive</i> <i>...the HPV vaccine was too expensive*</i>
2	<i>...my/our insurance does not cover enough of the cost of the HPV vaccine for [CHILD]</i> <i>...my/our insurance did not cover enough of the cost of the HPV vaccine for [CHILD] *</i>
3	<i>...the HPV vaccine costs more than I can afford</i>

...the HPV vaccine cost more than I could afford*

Barriers Harms (HABS ‘Harms’)

- 1 ...*the HPV vaccine is being pushed to make money for pharmaceutical companies*
 - 2 ...giving [CHILD] the HPV vaccine would be like performing an experiment on [CHILD]
 - 2 ...giving [CHILD] the HPV vaccine was like performing an experiment on [CHILD]*
 - 3 ...*the HPV vaccine may lead to long-term health problems*
 - 4 ...*the HPV vaccine is too new*
 - 5 ...*there has not been enough research done on the HPV vaccine*
 - 6 ...*the HPV vaccine is unsafe*
-

Cues to Action (HABS ‘Influence’)

- 1 ...*other parents in my community are getting their children the HPV vaccine*
 - 2 ...*my friends are getting their children vaccinated with the HPV vaccine*
 - 3 ...*other children around [CHILD]'s age are getting vaccinated with the HPV vaccine*
 - 4 ...it is expected of me that I should vaccinate [CHILD] against HPV
 - 4 ...it was expected of me that I should vaccinate [CHILD] against HPV*
 - 5 ...most of my friends think vaccinating [CHILD] against HPV is a good idea
 - 5 ...most of my friends think vaccinating [CHILD] against HPV was a good idea*
 - 6 ...*my [CHILD]'s other parent believes we should get the HPV vaccine for [CHILD]*
 - 6 ...my [CHILD]'s other parent believes in having gotten the HPV vaccine for [CHILD] *
 - 7 ...*doctors/health care providers believe vaccinating [CHILD] against HPV is a good idea*
 - 8 ...my family thinks it is a good idea to vaccinate [CHILD] against HPV
 - 8 ...my family thinks it was a good idea to vaccinate [CHILD] against HPV*
-

Self Efficacy +

- 1 ...*competent to make decisions about the vaccines [CHILD] receives*
 - 2 ...*competent to make decisions about [CHILD]'s health*
 - 3 ...*confident in making the decision about whether [CHILD] receives the HPV vaccine*
 - 4 ...*capable to make the decision about whether [CHILD] receives the HPV vaccine*
-

Source: Perez S, Shapiro GK, Tatar O, Joyal-Desmarais K, Rosberger Z. Development and validation of the human papillomavirus attitudes and beliefs scale in a National Canadian sample. *Sexually Transmitted Diseases* 2016; 43(10): 626-32.

Note. All items were modified to make them gender neutral (i.e. [CHILD] instead of ‘my son’). □ Signifies reverse coded items. * Signifies that this item has two possible stem phrasing depending on PAPM stage (PAPM Group 1-5, vs. 6). + items were developed to measure ‘self-efficacy’ (i.e. items not measured by Perez et al. 2016).

Table A3. General vaccine attitudes

Vaccine Conspiracy Beliefs Scale ¹	
1	Vaccine safety data is often fabricated
2	Negative vaccination effects are covered up *
3	Pharmaceutical companies cover up the dangers of vaccines
4	People are deceived about vaccine efficacy
5	Vaccine efficacy data is often fabricated
6	People are deceived about vaccine safety
7	The government is trying to cover up the link between vaccines and autism.
Vaccine Hesitancy Scale -Lack of Confidence ²	
1	Childhood vaccines are important for my child's health (R)
2	Childhood vaccines are effective (R)
3	Having my child vaccinated is important for the health of others in my community (R)
4	All childhood vaccines offered by the government program in my community are beneficial (R)
5	The information I receive about vaccines from the vaccine program is reliable and trustworthy (R)
6	Getting vaccines is a good way to protect my child/children from disease (R)
7	Generally I do what my doctor or health care provider recommends about vaccines for my child/children (R)
Vaccine Hesitancy Scale -Risks ²	
1	I am concerned about serious adverse effects of vaccines
2	New vaccines carry more risks than older vaccines

Sources: ¹ Shapiro GK, Holding A, Perez S, Amsel R, Rosberger Z. Validation of the vaccine conspiracy belief scale. *Papillomavirus Research* 2016; **2**: 167-72. ² Shapiro GK, Tatar O, Dube E, et al. The Vaccine Hesitancy Scale: Psychometric properties and validation. *Vaccine* 2018; **36**: 545-52.

Note. * One item from the original scale ('immunizing children is harmful and this fact is covered up') was modified slightly in this study's administration (to 'negative vaccination effects are covered up') as this was a double-barrelled question.

(R) Indicates items that were reverse coded.

Appendix B. Additional multinomial logistic regression analyses of parents' PAMP stage

1. Exploratory bivariate multinomial logistic regression analyses of parents of boys and girls (N = 3,779)

Exploratory bivariate multinomial logistic regression analyses were conducted to examine the relationship between additional variables of interest and parents' PAMP stage (Table B1).

Table B1. Bivariate multinomial logistic regression analysis of parents' PAMP stage (N = 3,779) for additional variables

Variables	Stage 1-Unaware OR (95% CI) n = 604	Stage 2-Unengaged OR (95% CI) n = 395	Stage 3-Undecided OR (95% CI) n = 680	Stage 4-Decided NOT OR (95% CI) n = 353	Stage 5-Decided YES OR (95% CI) n = 786
<i>Parent's gender</i>					
Male	(reference)	(reference)	(reference)	(reference)	(reference)
Female	0.58*** (0.47; 0.71)	0.49*** (0.38; 0.62)	0.81 (0.66; 1.00)	1.28 (0.97; 1.68)	0.84 (0.68; 1.03)
<i>Parent's age (one year increase)</i>					
	0.97*** (0.95; 0.98)	0.98 (0.96; 0.99)	0.98 (0.97; 0.99)	0.97** (0.95; 0.99)	0.97** (0.96; 0.99)
<i>Household income</i>					
< 100,000 CAD\$	(reference)	(reference)	(reference)	(reference)	(reference)
≥100,000 CAD\$	0.73* (0.58; 0.91)	0.82 (0.64; 1.05)	0.83 (0.67; 1.03)	0.72 (0.55; 0.95)	1.11 (0.91; 1.35)
Prefer not to answer	0.82 (0.59; 1.16)	0.64 (0.42; 0.98)	1.06 (0.77; 1.45)	1.25 (0.87; 1.81)	0.60* (0.42; 0.85)
<i>Parent's religion</i>					
No religious affiliation	(reference)	(reference)	(reference)	(reference)	(reference)
Any religious affiliation	1.09 (0.88; 1.36)	0.95 (0.74; 1.22)	0.89 (0.73; 1.10)	0.95 (0.73; 1.22)	0.99 (0.81; 1.21)
<i>Born in Canada</i>					
Yes	(reference)	(reference)	(reference)	(reference)	(reference)
No	1.82*** (1.37; 2.42)	1.55* (1.12; 2.16)	1.51* (1.14; 2.01)	1.37 (0.96; 1.95)	1.26 (0.95; 1.67)
<i>Child's school's religion</i>					
No affiliation	(reference)	(reference)	(reference)	(reference)	(reference)
Any religious affiliation	1.01 (0.80; 1.27)	0.97 (0.74; 1.27)	1.02 (0.82; 1.28)	0.90 (0.68; 1.20)	0.95 (0.76; 1.18)
<i>Child's school's language</i>					
English	(reference)	(reference)	(reference)	(reference)	(reference)
French and other	1.27 (1.03; 1.56)	0.71* (0.54; 0.92)	0.68** (0.55; 0.84)	1.04 (0.81; 1.35)	0.56*** (0.45; 0.69)
<i>Size of city</i>					
<100,000 inhabitants	(reference)	(reference)	(reference)	(reference)	(reference)
≥100,000 inhabitants	0.93 (0.76; 1.14)	1.17 (0.92; 1.48)	1.24 (1.02; 1.51)	1.00 (0.79; 1.28)	1.27 (1.05; 1.53)

Variables	Stage 1-Unaware OR (95% CI) <i>n</i> = 604	Stage 2-Unengaged OR (95% CI) <i>n</i> = 395	Stage 3-Undecided OR (95% CI) <i>n</i> = 680	Stage 4-Decided NOT OR (95% CI) <i>n</i> = 353	Stage 5-Decided YES OR (95% CI) <i>n</i> = 786
<i>Parent's level of education</i>					
Elementary or high school	(reference)	(reference)	(reference)	(reference)	(reference)
Trade technical or university	1.10 (0.85; 1.43)	1.17 (0.86; 1.59)	1.23 (0.95; 1.60)	1.00 (0.74; 1.36)	1.45* (1.12; 1.87)
<i>Parent's employment status</i>					
Employed	(reference)	(reference)	(reference)	(reference)	(reference)
Not employed	1.00 (0.77; 1.30)	0.77 (0.56; 1.06)	1.09 (0.85; 1.39)	1.35 (1.01; 1.80)	0.98 (0.77; 1.25)
<i>Marital status</i>					
Single, separated, divorced, widowed	(reference)	(reference)	(reference)	(reference)	(reference)
Married or common law	0.97 (0.76; 1.25)	1.30 (0.96; 1.77)	0.97 (0.76; 1.24)	0.93 (0.69; 1.26)	1.04 (0.82; 1.32)
<i>Number of children in the family</i>					
Three or more children	(reference)	(reference)	(reference)	(reference)	(reference)
One child	1.05 (0.79; 1.38)	0.97 (0.71; 1.34)	1.12 (0.85; 1.47)	0.72 (0.51; 1.00)	0.84 (0.64; 1.09)
Two children	0.90 (0.71; 1.14)	0.86 (0.66; 1.13)	1.05 (0.83; 1.33)	0.75 (0.57; 0.99)	1.02 (0.82; 1.27)
<i>Child's sexual orientation</i>					
Heterosexual	(reference)	(reference)	(reference)	(reference)	(reference)
Other sexual orientation	1.26 (0.93; 1.71)	1.17 (0.82; 1.67)	1.16 (0.86; 1.57)	1.33 (0.93; 1.91)	1.13 (0.84; 1.51)
<i>HPV General knowledge</i> (One-unit increase)					
	0.87*** (0.85; 0.88)	0.93*** (0.91; 0.95)	0.97** (0.95; 0.99)	1.03* (1.01; 1.06)	1.01 (0.99; 1.03)
<i>Province with funded HPV vaccination for boys</i>					
No	(reference)	(reference)	(reference)	(reference)	(reference)
Yes	0.34*** (0.24; 0.48)	0.49** (0.34; 0.70)	0.50*** (0.37; 0.67)	0.90 (0.66; 1.23)	0.92 (0.72; 1.17)

Note. The reference category for PAPM stage is 'Stage 6-Vaccinated' (*n* = 961). Bold integers indicates significant OR at $p < .05$. * $p < .01$, ** $p < .001$, *** $p < .0001$. (reference) = reference category. OR = odds ratio. 95% CI = 95% confidence interval. HCP = health care provider.

2. Bivariate and multivariate multinomial logistic regression analyses for parents of boys ($n = 1,826$)

Additional analyses were conducted to examine the correlates of PAMP stage for parents of boys. Table B2 reports findings from the bivariate multinomial logistic regression analysis of parents of boys' PAMP stage. Table B3 reports findings from the multivariate multinomial logistic regression analysis of parents of boys' PAMP stage. Finally, Table B4 reports on exploratory bivariate multinomial logistic regression analyses that were conducted to examine the relationship between additional variables of interest and parents of boys' PAMP stage.

Table B2. Bivariate multinomial logistic regression analysis of parents of boys' PAMP stage ($n = 1,826$)

Variables	Stage 1-Unaware OR (95% CI) $n = 468$	Stage 2-Unengaged OR (95% CI) $n = 298$	Stage 3-Undecided OR (95% CI) $n = 389$	Stage 4-Decided NOT OR (95% CI) $n = 162$	Stage 5-Decided YES OR (95% CI) $n = 349$
<i>Child's age</i> (One-year increase)	0.77*** (0.71; 0.84)	0.86** (0.79; 0.94)	0.84*** (0.78; 0.92)	0.82** (0.74; 0.91)	0.79*** (0.72; 0.86)
<i>Language parents answered the survey</i>	(reference)	(reference)	(reference)	(reference)	(reference)
English					
French	3.08*** (1.97; 4.82)	1.45 (0.89; 2.36)	1.28 (0.80; 2.06)	1.65 (0.96; 2.83)	0.84 (0.51; 1.39)
<i>Parent's ethnicity</i>					
Other	(reference)	(reference)	(reference)	(reference)	(reference)
White	0.82 (0.48; 1.39)	0.81 (0.46; 1.40)	0.94 (0.54; 1.61)	1.21 (0.62; 2.37)	0.83 (0.50; 1.42)
<i>HCP Recommendation</i>					
No	(reference)	(reference)	(reference)	(reference)	(reference)
Yes	0.01 (0.01; 243)	0.07*** (0.03; 0.16)	0.21*** (0.12; 0.35)	0.09*** (0.04; 0.24)	0.46* (0.29; 0.74)
<i>HPV Vaccine knowledge</i> (One-unit increase)	0.69*** (0.64; 0.74)	0.76*** (0.71; 0.82)	0.86*** (0.80; 0.93)	0.93 (0.85; 1.01)	0.98 (0.90; 1.05)
<i>Susceptibility</i> (One-unit increase)	0.45*** (0.38; 0.54)	0.40*** (0.33; 0.49)	0.40*** (0.33; 0.48)	0.15*** (0.12; 0.18)	0.96 (0.80; 1.15)
<i>Severity</i> (One-unit increase)	0.74* (0.61; 0.89)	0.71** (0.58; 0.87)	0.71** (0.58; 0.86)	0.42*** (0.34; 0.52)	1.04 (0.85; 1.28)
<i>Benefits</i> (One-unit increase)	0.50*** (0.41; 0.62)	0.41*** (0.33; 0.51)	0.39*** (0.31; 0.48)	0.08*** (0.05; 0.10)	1.26 (1.02; 1.56)
<i>Barriers-Affordability</i> (One-unit increase)	2.56*** (2.23; 2.95)	2.27*** (1.96; 2.62)	2.39*** (2.08; 2.76)	1.50*** (1.28; 1.74)	2.17*** (1.88; 2.50)
<i>Barriers-Accessibility</i> (One-unit increase)	3.01*** (2.50; 3.63)	2.48*** (2.04; 3.01)	2.26*** (1.88; 2.72)	1.86*** (1.50; 2.30)	1.42** (1.19; 1.70)
<i>Barriers-Harms</i> (One-unit increase)	2.26*** (1.91; 2.67)	2.46*** (2.06; 2.95)	2.76*** (2.31; 3.29)	8.58*** (6.73; 10.93)	1.09 (0.92; 1.28)
<i>Cues to action</i> (One-unit increase)	0.16*** (0.13; 0.21)	0.12*** (0.09; 0.15)	0.13*** (0.10; 0.17)	0.05*** (0.04; 0.06)	0.50*** (0.41; 0.62)
<i>Self-Efficacy</i> (One-unit increase)	0.52*** (0.42; 0.64)	0.50*** (0.40; 0.63)	0.51*** (0.41; 0.63)	0.94 (0.72; 1.23)	0.96 (0.76; 1.20)

Note. The reference category for PAPM stage is 'Stage 6-Vaccinated' (n = 160). Bold integers indicates significant OR at $p < .05$, * $p < .01$, ** $p < .001$, *** $p < .0001$. (reference) = reference category. OR = odds ratio. 95% CI = 95% confidence interval. HCP = health care provider.

Table B3. Multivariate multinomial logistic regression analysis of parents of boys' PAPM stage ($n = 1,826$)

Variables	Stage 1-Unaware AOR (95% CI) $n = 468$	Stage 2-Unengaged AOR (95% CI) $n = 298$	Stage 3-Undecided AOR (95% CI) $n = 389$	Stage 4-Decided AOR (95% CI) $n = 162$	Stage 5-Decided YES AOR (95% CI) $n = 349$
<i>Child's age</i> (One-year increase)	0.75*** (0.67; 0.84)	0.84* (0.74; 0.94)	0.82** (0.73; 0.92)	0.88 (0.76; 1.01)	0.74*** (0.66; 0.82)
<i>Language parents answered the survey</i>	(reference)	(reference)	(reference)	(reference)	(reference)
English					
French	1.45 (0.81; 2.61)	0.74 (0.40; 1.37)	0.69 (0.37; 1.26)	0.81 (0.38; 1.73)	0.51 (0.28; 0.92)
<i>Parent's ethnicity</i>					
Other	(reference)	(reference)	(reference)	(reference)	(reference)
White	1.16 (0.59; 2.28)	1.20 (0.60; 2.40)	1.39 (0.70; 2.74)	1.17 (0.47; 2.95)	0.84 (0.44; 1.59)
<i>HCP Recommendation</i>					
No	(reference)	(reference)	(reference)	(reference)	(reference)
Yes	NA	0.17*** (0.07; 0.44)	0.40 (0.20; 0.81)	0.10* (0.02; 0.43)	0.62 (0.35; 1.10)
<i>HPV Vaccine knowledge</i> (One-unit increase)	0.81*** (0.74; 0.89)	0.92 (0.84; 1.01)	1.04 (0.95; 1.14)	1.14 (1.01; 1.28)	1.03 (0.94; 1.13)
<i>Susceptibility</i> (One-unit increase)	0.89 (0.66; 1.21)	0.90 (0.66; 1.22)	0.86 (0.63; 1.16)	0.68 (0.47; 0.99)	1.00 (0.75; 1.34)
<i>Severity</i> (One-unit increase)	0.99 (0.77; 1.30)	1.01 (0.77; 1.33)	1.01 (0.77; 1.31)	0.60* (0.44; 0.83)	1.04 (0.80; 1.34)
<i>Benefits</i> (One-unit increase)	1.83* (1.24; 2.71)	1.50 (1.01; 2.23)	1.34 (0.91; 1.98)	0.64 (0.39; 1.03)	2.08*** (1.43; 3.04)
<i>Barriers-Affordability</i> (One-unit increase)	2.10*** (1.75; 2.51)	1.87*** (1.56; 2.25)	2.05*** (1.71; 2.45)	1.52** (1.21; 1.92)	1.83*** (1.55; 2.15)
<i>Barriers-Accessibility</i> (One-unit increase)	1.39 (1.08; 1.80)	1.15 (0.88; 1.51)	1.04 (0.81; 1.35)	1.09 (0.77; 1.54)	1.02 (0.80; 1.30)
<i>Barriers-Harms</i> (One-unit increase)	1.28 (1.01; 1.63)	1.37 (1.07; 1.76)	1.55** (1.22; 1.97)	2.63*** (1.91; 3.62)	0.94 (0.74; 1.18)
<i>Cues to action</i> (One-unit increase)	0.21*** (0.15; 0.30)	0.15*** (0.11; 0.22)	0.18*** (0.12; 0.25)	0.15*** (0.10; 0.23)	0.35*** (0.25; 0.49)
<i>Self-Efficacy</i> (One-unit increase)	1.05 (0.75; 1.48)	1.04 (0.73; 1.47)	0.97 (0.69; 1.36)	1.68 (1.13; 2.52)	1.08 (0.76; 1.54)

Note. The reference category for PAPM stage is 'Stage 6-Vaccinated' ($n = 160$). Bold integers indicates significant AOR at $p < .05$, $*p < .01$, $**p < .001$, $***p < .0001$; NA when cell size = 0. Cox-Snell $R^2 = 0.60$. Cragg Uhler $R^2 = 0.62$. McFadden $R^2 = 0.27$. (reference) = reference category. AOR = adjusted odds ratio. 95% CI = 95% confidence interval. HCP = health care provider.

Table B4. Bivariate multinomial logistic regression analysis of parents of boys' PAPM stage ($n = 1,826$) for additional variables

Variables	Stage 1-Unaware OR (95% CI) $n = 468$	Stage 2-Unengaged OR (95% CI) $n = 298$	Stage 3-Undecided OR (95% CI) $n = 389$	Stage 4-Decided NOT OR (95% CI) $n = 162$	Stage 5-Decided YES OR (95% CI) $n = 349$
<i>Parent's gender</i>					
Male	(reference)	(reference)	(reference)	(reference)	(reference)
Female	0.80 (0.55; 1.17)	0.63 (0.42; 0.94)	1.04 (0.71; 1.54)	1.24 (0.78; 1.99)	1.04 (0.70; 1.54)
<i>Parent's age</i> (one year increase)	0.96** (0.93; 0.98)	0.97 (0.95; 1.00)	0.98 (0.95; 1.00)	0.95** (0.92; 0.98)	0.97 (0.94; 0.99)
<i>Household income</i>					
< 100,000 CAD\$	(reference)	(reference)	(reference)	(reference)	(reference)
>100,000 CAD\$	0.67 (0.45; 0.99)	0.79 (0.52; 1.20)	0.79 (0.53; 1.18)	0.57 (0.35; 0.93)	1.12 (0.75; 1.67)
Prefer not to answer	0.61 (0.35; 1.06)	0.41* (0.21; 0.77)	0.74 (0.42; 1.29)	0.71 (0.37; 1.36)	0.50 (0.27; 0.93)
<i>Parent's religion</i>					
No religious affiliation	(reference)	(reference)	(reference)	(reference)	(reference)
Any religious affiliation	1.43 (0.99; 2.08)	1.23 (0.83; 1.83)	1.06 (0.73; 1.54)	1.30 (0.82; 2.05)	1.39 (0.94; 2.05)
<i>Born in Canada</i>					
Yes	(reference)	(reference)	(reference)	(reference)	(reference)
No	1.63 (0.94; 2.81)	1.51 (0.85; 2.70)	1.55 (0.89; 2.72)	1.44 (0.75; 2.76)	1.41 (0.80; 2.50)
<i>Child's school's religion</i>					
No affiliation	(reference)	(reference)	(reference)	(reference)	(reference)
Any religious affiliation	1.24 (0.80; 1.90)	1.29 (0.82; 2.04)	1.25 (0.80; 1.94)	1.18 (0.70; 1.98)	1.39 (0.89; 2.17)
<i>Child's school's language</i>					
English	(reference)	(reference)	(reference)	(reference)	(reference)
French and other	2.86*** (1.86; 4.39)	1.39 (0.87; 2.22)	1.40 (0.90; 2.20)	2.11* (1.28; 3.50)	0.88 (0.55; 1.42)
<i>Size of city</i>					
<100,000 inhabitants	(reference)	(reference)	(reference)	(reference)	(reference)
>100,000 inhabitants	1.03 (0.72; 1.48)	1.38 (0.94; 2.03)	1.26 (0.87; 1.82)	0.95 (0.62; 1.47)	1.32 (0.90; 1.91)
<i>Parent's level of education</i>					
Elementary or high school	(reference)	(reference)	(reference)	(reference)	(reference)
Trade technical or university	0.98 (0.62; 1.57)	1.07 (0.65; 1.77)	1.40 (0.85; 2.30)	0.87 (0.50; 1.51)	1.18 (0.72; 1.94)
<i>Parent's employment status</i>					
Employed	(reference)	(reference)	(reference)	(reference)	(reference)
Not employed	0.77 (0.51; 1.18)	0.53 (0.33; 0.86)	0.75 (0.49; 1.16)	0.89 (0.53; 1.48)	0.79 (0.51; 1.23)
<i>Marital status</i>					
Single, separated, divorced, widowed	(reference)	(reference)	(reference)	(reference)	(reference)
Married or common law	0.96 (0.61; 1.49)	1.32 (0.81; 2.16)	0.83 (0.53; 1.29)	0.88 (0.52; 1.49)	1.07 (0.67; 1.71)
<i>Number of children in the family</i>					

Variables	Stage 1-Unaware OR (95% CI) n = 468	Stage 2-Unengaged OR (95% CI) n = 298	Stage 3-Undecided OR (95% CI) n = 389	Stage 4-Decided NOT OR (95% CI) n = 162	Stage 5-Decided YES OR (95% CI) n= 349
Three or more children	(reference)	(reference)	(reference)	(reference)	(reference)
One child	1.28 (0.78; 2.11)	1.25 (0.73; 2.13)	1.57 (0.94; 2.63)	0.99 (0.55; 1.82)	1.09 (0.64; 1.86)
Two children	0.99 (0.66; 1.49)	0.97 (0.62; 1.50)	1.35 (0.88; 2.06)	0.78 (0.47; 1.28)	1.23 (0.80; 1.87)
<i>Child's sexual orientation</i>					
Heterosexual	(reference)	(reference)	(reference)	(reference)	(reference)
Other sexual orientation	1.12 (0.64; 1.96)	1.08 (0.59; 1.98)	0.80 (0.44; 1.46)	0.86 (0.42; 1.76)	1.02 (0.57; 1.84)
<i>HPV General knowledge</i> (One-unit increase)	0.87*** (0.84; 0.90)	0.93*** (0.89; 0.96)	0.97 (0.93; 0.99)	1.01 (0.97; 1.05)	1.02 (0.98; 1.05)
<i>Vaccine conspiracy beliefs</i> (One-unit increase)	1.58*** (1.37; 1.82)	1.59*** (1.36; 1.84)	1.59*** (1.37; 1.83)	3.11*** (2.58; 3.74)	1.01 (0.87; 1.17)
<i>Hesitancy-Lack of Confidence</i> (7 items) (One-unit increase)	2.35*** (1.71; 3.22)	2.53*** (1.81; 3.54)	2.61*** (1.89; 3.61)	9.01*** (6.23; 13.03)	0.83 (0.59; 1.16)
<i>Hesitancy-Risks</i> (2 items) (One-unit increase)	1.93*** (1.58; 2.36)	1.91*** (1.54; 2.37)	2.38*** (1.92; 2.93)	5.06*** (3.83; 6.68)	1.12 (0.91; 1.37)
<i>Province with funded HPV vaccination for boys</i>					
No	(reference)	(reference)	(reference)	(reference)	(reference)
Yes	0.09*** (0.05; 0.15)	0.16*** (0.10; 0.26)	0.10*** (0.06; 0.17)	0.35*** (0.21; 0.57)	0.39*** (0.26; 0.59)

Note. The reference category for PAPM stage is 'Stage 6-Vaccinated' (n = 160). Bold integers indicates significant OR at $p < .05$, $*p < .01$, $**p < .001$, $***p < .0001$. (reference) = reference category. OR = odds ratio. 95% CI = 95% confidence interval.

3. Bivariate and multivariate multinomial logistic regression analyses for parents of girls ($n = 1,953$)

Additional analyses were conducted to examine the correlates of PAMP stage for parents of girls. Table B5 reports findings from the bivariate multinomial logistic regression analysis of parents of girls' PAMP stage. Table B6 reports findings from the multivariate multinomial logistic regression analysis of parents of girls' PAMP stage. Finally, Table B7 reports on exploratory bivariate multinomial logistic regression analyses that were conducted to examine the relationship between additional variables of interest and parents of girls' PAMP stage.

Table B5. Bivariate multinomial logistic regression analysis of parents of girls' PAMP stage ($n = 1,953$)

Variables	Stage 1-Unaware OR (95% CI) $n = 136$	Stage 2-Unengaged OR (95% CI) $n = 97$	Stage 3-Undecided OR (95% CI) $n = 291$	Stage 4-Decided NOT OR (95% CI) $n = 191$	Stage 5-Decided YES OR (95% CI) $n = 437$
<i>Child's age</i> (One-year increase)	0.62*** (0.57; 0.68)	0.65*** (0.59; 0.72)	0.66*** (0.62; 0.70)	0.80*** (0.74; 0.86)	0.65*** (0.62; 0.69)
<i>Language parents answered the survey</i>	(reference)	(reference)	(reference)	(reference)	(reference)
English					
French	0.63 (0.42; 0.96)	0.48* (0.29; 0.81)	0.45*** (0.32; 0.63)	0.84 (0.60; 1.19)	0.48*** (0.36; 0.63)
<i>Parent's ethnicity</i>					
Other	(reference)	(reference)	(reference)	(reference)	(reference)
White	0.45** (0.29; 0.72)	0.41** (0.25; 0.69)	0.65 (0.44; 0.95)	0.78 (0.49; 1.24)	0.68 (0.49; 0.96)
<i>HCP Recommendation</i>					
No	(reference)	(reference)	(reference)	(reference)	(reference)
Yes	0.06*** (0.01; 0.24)	0.08** (0.02; 0.34)	0.42*** (0.28; 0.65)	0.77 (0.51; 1.17)	0.92 (0.68; 1.23)
<i>HPV Vaccine knowledge</i> (One-unit increase)	0.58*** (0.54; 0.62)	0.69*** (0.64; 0.74)	0.80*** (0.76; 0.84)	1.01 (0.94; 1.09)	0.91** (0.87; 0.96)
<i>Susceptibility</i> (One-unit increase)	0.50*** (0.43; 0.59)	0.50*** (0.42; 0.60)	0.40*** (0.36; 0.46)	0.18*** (0.15; 0.21)	1.04 (0.93; 1.16)
<i>Severity</i> (One-unit increase)	0.74** (0.62; 0.87)	0.75* (0.62; 0.91)	0.77*** (0.68; 0.88)	0.54*** (0.47; 0.62)	1.00 (0.88; 1.13)
<i>Benefits</i> (One-unit increase)	0.50*** (0.40; 0.61)	0.51*** (0.40; 0.64)	0.33*** (0.28; 0.39)	0.08*** (0.06; 0.10)	1.32*** (1.15; 1.50)
<i>Barriers-Affordability</i> (One-unit increase)	2.91*** (2.52; 3.37)	2.13*** (1.82; 2.49)	1.98*** (1.79; 2.20)	1.25** (1.10; 1.41)	1.77*** (1.62; 1.95)
<i>Barriers-Accessibility</i> (One-unit increase)	3.60*** (2.96; 4.37)	2.75*** (2.23; 3.40)	2.39*** (2.08; 2.74)	1.74*** (1.49; 2.03)	1.53*** (1.36; 1.72)
<i>Barriers-Harms</i> (One-unit increase)	2.38*** (2.01; 2.82)	2.34*** (1.94; 2.84)	3.84*** (3.30; 4.47)	9.64*** (7.74; 12.00)	1.20** (1.08; 1.33)
<i>Cues to action</i> (One-unit increase)	0.17*** (0.13; 0.22)	0.15*** (0.12; 0.20)	0.15*** (0.12; 0.19)	0.08*** (0.06; 0.10)	0.67*** (0.58; 0.76)
<i>Self-Efficacy</i> (One-unit increase)	0.42*** (0.36; 0.50)	0.50*** (0.41; 0.62)	0.52*** (0.45; 0.60)	1.09 (0.90; 1.33)	1.01 (0.87; 1.16)

Note. The reference category for PAPM stage is ‘Stage 6-Vaccinated’ (n = 801). Bold integers indicates OR at $p < .05$, * $p < .01$, ** $p < .001$, *** $p < .0001$. (reference) = reference category. OR = odds ratio. 95% CI = 95% confidence interval. HCP = health care provider.

Table B6. Multivariate multinomial logistic regression analysis of parents of girls' PAPM stage ($n = 1,953$)

Variables	Stage 1-Unaware AOR (95% CI) $n = 136$	Stage 2-Unengaged AOR (95% CI) $n = 97$	Stage 3-Undecided AOR (95% CI) $n = 291$	Stage 4-Decided NOT AOR (95% CI) $n = 191$	Stage 5-Decided YES AOR (95% CI) $n = 437$
<i>Child's age</i> (One-year increase)	0.54*** (0.48; 0.60)	0.61*** (0.55; 0.69)	0.62*** (0.57; 0.67)	0.82* (0.72; 0.93)	0.60*** (0.56; 0.64)
<i>Language parents answered the survey</i>	(reference)	(reference)	(reference)	(reference)	(reference)
English					
French	0.80 (0.45; 1.40)	0.57 (0.31; 1.05)	0.62** (0.30; 0.74)	0.82 (0.51; 1.79)	0.60*** (0.33; 0.65)
<i>Parent's ethnicity</i>					
Other	(reference)	(reference)	(reference)	(reference)	(reference)
White	1.48 (0.80; 2.75)	1.11 (0.59; 2.07)	1.76 (1.06; 2.93)	0.96 (0.44; 2.10)	1.75* (1.15; 2.67)
<i>HCP Recommendation</i>					
No	(reference)	(reference)	(reference)	(reference)	(reference)
Yes	0.11* (0.02; 0.50)	0.15 (0.04; 0.66)	0.56 (0.32; 0.98)	0.39 (0.17; 0.91)	0.92 (0.65; 1.32)
<i>HPV Vaccine knowledge</i> (One-unit increase)	0.68*** (0.62; 0.75)	0.81*** (0.74; 0.89)	0.96 (0.89; 1.03)	1.16 (1.03; 1.30)	0.92* (0.87; 0.98)
<i>Susceptibility</i> (One-unit increase)	1.13 (0.82; 1.55)	1.08 (0.79; 1.48)	0.88 (0.71; 1.10)	0.64* (0.49; 0.85)	1.08 (0.90; 1.30)
<i>Severity</i> (One-unit increase)	1.20 (0.93; 1.56)	1.08 (0.83; 1.41)	1.10 (0.90; 1.34)	0.68* (0.52; 0.89)	0.99 (0.84; 1.17)
<i>Benefits</i> (One-unit increase)	1.97* (1.30; 2.97)	1.91* (1.25; 2.91)	1.45 (1.08; 1.96)	0.56* (0.38; 0.84)	2.18*** (1.72; 2.76)
<i>Barriers-Affordability</i> (One-unit increase)	2.35*** (1.90; 2.91)	1.48** (1.19; 1.83)	1.45*** (1.24; 1.70)	0.97 (0.75; 1.26)	1.65*** (1.46; 1.86)
<i>Barriers-Accessibility</i> (One-unit increase)	1.61* (1.20; 2.16)	1.49 (1.10; 2.01)	1.17 (0.93; 1.46)	1.08 (0.77; 1.52)	1.27* (1.07; 1.52)
<i>Barriers-Harms</i> (One-unit increase)	1.34 (1.03; 1.74)	1.60** (1.23; 2.08)	2.82*** (2.33; 3.42)	4.05*** (3.05; 5.38)	1.20 (1.03; 1.39)
<i>Cues to action</i> (One-unit increase)	0.30*** (0.21; 0.44)	0.21*** (0.14; 0.30)	0.27*** (0.21; 0.36)	0.24*** (0.16; 0.35)	0.62*** (0.50; 0.76)
<i>Self-Efficacy</i> (One-unit increase)	0.87 (0.66; 1.16)	1.15 (0.86; 1.53)	1.12 (0.90; 1.39)	1.86*** (1.38; 2.50)	1.26 (1.01; 1.56)

Note. The reference category for PAPM stage is 'Stage 6-Vaccinated' ($n = 801$). Bold integers indicates significant AOR at $p < .05$, * $p < .01$, ** $p < .001$, *** $p < .0001$. Cox-Snell $R^2 = 0.71$. Cragg Uhler $R^2 = 0.75$. McFadden $R^2 = 0.41$. (reference) = reference category. AOR = adjusted odds ratio. 95% CI = 95% confidence interval. HCP = health care provider.

Table B7. Bivariate multinomial logistic regression analysis of parents of girls' PAPM stage ($n = 1,953$) for additional variables

Variables	Stage 1-Unaware OR (95% CI) $n = 136$	Stage 2-Unengaged OR (95% CI) $n = 97$	Stage 3-Undecided OR (95% CI) $n = 291$	Stage 4-Decided NOT OR (95% CI) $n = 191$	Stage 5-Decided YES OR (95% CI) $n = 437$
<i>Parent's gender</i>					
Male	(reference)	(reference)	(reference)	(reference)	(reference)
Female	0.37*** (0.26; 0.54)	0.40*** (0.26; 0.61)	0.74 (0.56; 0.98)	1.53 (1.05; 2.24)	0.80 (0.62; 1.03)
<i>Parent's age</i> (one year increase)	0.96* (0.93; 0.99)	0.96* (0.93; 0.99)	0.97* (0.95; 0.99)	0.98 (0.95; 1.00)	0.97** (0.95; 0.98)
<i>Household income</i>					
< 100,000 CAD\$	(reference)	(reference)	(reference)	(reference)	(reference)
>100,000 CAD\$	0.61 (0.41; 0.91)	0.58 (0.36; 0.94)	0.76 (0.56; 1.02)	0.80 (0.56; 1.14)	1.02 (0.80; 1.30)
Prefer not to answer	0.62 (0.32; 1.22)	0.80 (0.39; 1.63)	1.11 (0.73; 1.71)	1.57 (0.99; 2.50)	0.54 (0.34; 0.87)
<i>Parent's religion</i>					
No religious affiliation	(reference)	(reference)	(reference)	(reference)	(reference)
Any religious affiliation	0.97 (0.66; 1.43)	0.89 (0.57; 1.39)	0.95 (0.71; 1.26)	0.85 (0.61; 1.18)	0.87 (0.68; 1.12)
<i>Born in Canada</i>					
Yes	(reference)	(reference)	(reference)	(reference)	(reference)
No	2.67*** (1.72; 4.15)	1.76 (1.01; 3.06)	1.48 (1.01; 2.17)	1.32 (0.84; 2.09)	1.16 (0.81; 1.65)
<i>Child's school's religion</i>					
No affiliation	(reference)	(reference)	(reference)	(reference)	(reference)
Any religious affiliation	1.11 (0.99; 1.02)	0.81 (1.01; 1.03)	1.05 (0.99; 1.02)	0.85 (1.00; 1.02)	0.81 (0.99; 1.01)
<i>Child's school's language</i>					
English	(reference)	(reference)	(reference)	(reference)	(reference)
French and other	0.76 (0.51; 1.12)	0.68 (0.42; 1.08)	0.59** (0.43; 0.79)	0.93 (0.66; 1.29)	0.59*** (0.45; 0.76)
<i>Size of city</i>					
<100,000 inhabitants	(reference)	(reference)	(reference)	(reference)	(reference)
>100,000 inhabitants	0.78 (0.54; 1.12)	0.82 (0.54; 1.25)	1.29 (0.99; 1.69)	1.09 (0.79; 1.49)	1.27 (1.01; 1.61)
<i>Parent's level of education</i>					
Elementary or high school	(reference)	(reference)	(reference)	(reference)	(reference)
Trade technical or university	1.22 (0.75; 1.97)	1.17 (0.67; 2.02)	0.97 (0.70; 1.36)	1.07 (0.71; 1.59)	1.62* (1.17; 2.24)
<i>Parent's employment status</i>					
Employed	(reference)	(reference)	(reference)	(reference)	(reference)
Not employed	0.71 (0.42; 1.21)	0.86 (0.48; 1.53)	1.19 (0.85; 1.67)	1.53 (1.05; 2.23)	0.94 (0.69; 1.28)
<i>Marital status</i>					
Single, separated, divorced, widowed	(reference)	(reference)	(reference)	(reference)	(reference)
Married or common law	1.08 (0.68; 1.71)	1.29 (0.74; 2.27)	1.26 (0.89; 1.79)	1.00 (0.67; 1.47)	1.03 (0.77; 1.38)
<i>Number of children in the family</i>					

Variables	Stage 1-Unaware OR (95% CI) n = 136	Stage 2-Unengaged OR (95% CI) n = 97	Stage 3-Undecided OR (95% CI) n = 291	Stage 4-Decided NOT OR (95% CI) n = 191	Stage 5-Decided YES OR (95% CI) n= 437
Three or more children	(reference)	(reference)	(reference)	(reference)	(reference)
One child	1.36 (0.82; 2.24)	1.04 (0.58; 1.86)	0.99 (0.69; 1.43)	0.65 (0.42; 1.02)	0.81 (0.58; 1.12)
Two children	1.16 (0.74; 1.81)	0.99 (0.60; 1.63)	0.91 (0.67; 1.25)	0.81 (0.56; 1.15)	0.98 (0.75; 1.29)
<i>Child's sexual orientation</i>					
Heterosexual	(reference)	(reference)	(reference)	(reference)	(reference)
Other sexual orientation	1.78 (1.09; 2.89)	1.45 (0.80; 2.61)	1.68* (1.16; 2.44)	1.77* (1.16; 2.72)	1.21 (0.85; 1.72)
<i>HPV General knowledge</i> (One-unit increase)	0.82*** (0.79; 0.85)	0.89*** (0.86; 0.92)	0.96** (0.93; 0.98)	1.04* (1.01; 1.08)	1.00 (0.98; 1.03)
<i>Vaccine conspiracy beliefs</i> (One-unit increase)	1.69*** (1.47; 1.95)	1.47*** (1.25; 1.72)	1.80*** (1.61; 2.01)	3.34*** (2.87; 3.89)	0.97 (0.89; 1.06)
<i>Hesitancy-Lack of Confidence</i> (7 items) (One-unit increase)	2.87*** (2.13; 3.86)	2.86*** (2.04; 4.01)	3.51*** (2.79; 4.42)	10.22*** (7.74;13.50)	0.92 (0.74; 1.14)
<i>Hesitancy-Risks</i> (2 items) (One-unit increase)	1.79*** (1.44; 2.22)	1.66*** (1.29; 2.12)	2.74*** (2.30; 3.27)	6.09*** (4.80; 7.73)	1.02 (0.90; 1.16)
<i>Province with funded HPV vaccination for boys</i>					
No	(reference)	(reference)	(reference)	(reference)	(reference)
Yes	0.90 (0.54; 1.52)	0.86 (0.47; 1.59)	1.07 (0.74; 1.55)	1.12 (0.73; 1.71)	1.04 (0.76; 1.44)

Note. The reference category for PAPM stage is 'Stage 6-Vaccinated' (n = 801). Bold integers indicates significant odds ratio (OR) at $p < .05$, * $p < .01$, ** $p < .001$, *** $p < .0001$. (reference) = reference category. OR = odds ratio. 95% CI = 95% confidence interval.