

**PURDUE UNIVERSITY
GRADUATE SCHOOL
Thesis/Dissertation Acceptance**

This is to certify that the thesis/dissertation prepared

By Michael P. Smoker

Entitled

Effect of Gabra2 Expression in the Central Nucleus of the Amygdala on Anxiety and Alcohol's Anxiolytic Capacity in C57BL/6J Mice.

For the degree of Master of Science

Is approved by the final examining committee:

Stephen L. Boehm II

Chair

Christopher C. Lapish

Cristine L. Czachowski

To the best of my knowledge and as understood by the student in the Thesis/Dissertation Agreement, Publication Delay, and Certification Disclaimer (Graduate School Form 32), this thesis/dissertation adheres to the provisions of Purdue University's "Policy of Integrity in Research" and the use of copyright material.

Approved by Major Professor(s): Stephen L. Boehm II

Approved by: Nicholas J. Grahame

Head of the Departmental Graduate Program

3/24/2016

Date

EFFECT OF CENTRAL AMYGDALA *GABRA2* EXPRESSION ON ANXIETY AND
ALCOHOL'S ANXIOLYTIC CAPACITY IN C57BL/6J MICE

A Thesis

Submitted to the Faculty

of

Purdue University

by

Michael P. Smoker

In Partial Fulfillment of the

Requirements for the Degree

of

Master of Science

May 2016

Purdue University

Indianapolis, Indiana

ACKNOWLEDGEMENTS

I would like to thank my committee, Drs. Steve Boehm, Chris Lapish, and Cris Czachowski, for their time and support. In addition, I'd like to thank Drs. Steve Boehm and Chris Lapish for their mentorship in my graduate studies. I would like to thank former lab member Dr. Caroline Quoilin and current lab members Brandon Fritz and Chelsea Kasten for their guidance and advice in research. Lastly, I'd like to acknowledge Yanping Zhang and William O'Neal for their assistance with various aspects of this project.

TABLE OF CONTENTS

	Page
LIST OF TABLES	v
LIST OF FIGURES	vi
ABSTRACT	vii
CHAPTER 1 INTRODUCTION	1
1.1 <i>GABRA2</i> & Alcohol-Related Behavior	1
1.2 Anxiety's Mediation of <i>GABRA2</i> & Alcohol-Related Behavior	3
1.3 <i>GABRA2</i> & Anxiety-Related Behavior	4
1.4 GABA _A Involvement in CeA-Mediated Behavior	6
1.5 <i>GABRA2</i> & the CeA	7
1.6 Specific Aims	9
CHAPTER 2 EFFECT OF CENTRAL AMYGDALA <i>GABRA2</i> EXPRESSION ON ANXIETY AND ALCOHOL'S ANXIOLYTIC CAPACITY	10
2.1 Materials and Methods	10
2.1.1 General Design	10
2.1.2 Subjects	10
2.1.3 Surgery	11
2.1.4. EPM Assessment	12
2.1.5. Histology	14
2.1.6. Statistical Analysis	15
2.2 Results	17
2.2.1. Aim 1	17
2.2.2. Histology	17
2.2.3. EPM Assessment	18

	Page
2.2.4. Brain-Behavior Correlations.....	19
CHAPTER 3 DISCUSSION.....	20
3.1. EPM & Alcohol's Anxiolytic Capacity	20
3.2. Virally-Mediated Gabra2 Knock-Down	20
3.3. CeA <i>Gabra2</i> Expression & Basal Anxiety	20
3.4. CeA <i>Gabra2</i> Expression & Alcohol's Anxiolytic Capacity.....	23
3.5. Conclusions, Limitations, & Future Directions	24
REFERENCES	29
TABLES	33
FIGURES.....	35
VITA.....	42

LIST OF TABLES

Table	Page
Table 1. Group sizes overall and for specific analyses.....	33
Table 2. Correlations between behavior and CeA $\alpha 2$ expression.....	34

LIST OF FIGURES

Figure	Page
Figure 1. RNA interference via shRNA.....	35
Figure 2. Sandwich ELISA.....	36
Figure 3. EPM behaviors in Aim 1	37
Figure 4. Lentiviral incorporation in the CeA.	38
Figure 5. CeA $\alpha 2$ protein concentrations	39
Figure 6. EPM arm entries	39
Figure 7. EPM total and average time	40
Figure 8. EPM head dips.....	41
Figure 9. Brain-behavior correlations	41

ABSTRACT

Smoker, Michael P. M.S., Purdue University, May 2016. Effect of Central Amygdala *Gabra2* Expression on Anxiety and Alcohol's Anxiolytic Capacity in C57BL/6J Mice. Major Professor: Stephen L. Boehm, II.

The *GABRA2* gene, which encodes the $\alpha 2$ subunit of GABA_A receptors, is one of the genes most frequently associated with alcohol-related behavior in human studies (Demers, Bogdan, & Agrawal, 2014). Polymorphisms in *GABRA2* have been found to be associated with alcohol dependence, changes in drinking frequency, and alcohol's stimulating and euphoric effects (Arias et al., 2014; Dick et al., 2014; Edenberg et al., 2004). However, the *GABRA2*-alcohol relationship may not be direct, as anxiety and impulsiveness have been found to be mediating factors (Enoch, Schwartz, Albaugh, Virkkunen, & Goldman, 2006; Villafuerte, Strumba, Stoltenberg, Zucker, & Burmeister, 2013).

Comorbidity of anxiety and alcohol use disorders is both prevalent and clinically relevant (J. P. Smith & Randall, 2012), and GABA_A receptors play a significant role in each. Benzodiazepines, primary pharmacologic treatments for anxiety disorders and alcohol withdrawal, facilitate signaling at GABA_A receptors, and their anxiolytic effects appear to depend on the presence of $\alpha 2$ subunits in these receptors (Low et al., 2000). The amygdala is widely implicated in both anxiety disorders as well as addiction (Janak & Tye, 2015), and its central nucleus is an important mediator of responses to both

alcohol- and stress-related stimuli (Roberto, Gilpin, & Siggins, 2012), some of which may be related to *GABRA2* expression within this region (Jin et al., 2014).

The aim of the current study was to explore the role of *Gabra2* (mouse ortholog of *GABRA2*) expression within the central nucleus of the amygdala (CeA) in anxiety-related behavior and alcohol's anxiolytic effects in mice. C57BL/6J (B6) mice underwent surgery for bilateral infusion of GFP-tagged lentivirus targeting *Gabra2* or a scramble control lentivirus into the CeA. Following 12-13 days of recovery, mice were assessed for anxiety-like behavior in the elevated plus maze (EPM) naïve or following IP injection of 0, 0.75, or 1.5 g/kg ethanol. After assessment, brains were extracted and sectioned through the CeA. Finally, GFP was quantified, the CeA was collected via laser microdissection, and $\alpha 2$ protein was quantified via ELISA.

In mice expressing GFP in the CeA, $\alpha 2$ protein concentrations were lower for Virus mice relative to Control mice. The EPM was anxiogenic, and alcohol was found to be anxiolytic. In naïve mice, while there was no difference between Control mice and Virus mice on any behavioral measure, there were significant correlations between CeA $\alpha 2$ protein concentration and time spent in closed arms as well as both total and average time spent in open arms. In mice receiving injection of 0, 0.75, or 1.5 g/kg ethanol, there was a main effect of dose on several behavioral measures, but no interaction between viral condition and dose, and only a main effect of viral condition on average time spent in closed arms. There were no significant correlations between CeA $\alpha 2$ protein concentration and behavioral measures within any injected dose. These results are consistent with *GABRA2*-anxiety associations and effects of *Gabra2* manipulation on anxiety-like behavior. Furthermore, they suggest that CeA $\alpha 2$ protein concentration is

positively related to basal anxiety, which could affect alcohol use through various routes. However, these results also suggest that CeA $\alpha 2$ protein concentration is not related to alcohol's anxiolytic capacity, at least when acutely administered in alcohol-naïve animals.

CHAPTER 1 INTRODUCTION

1.1 *GABRA2* & Alcohol-Related Behavior

Alcohol is a widely consumed drug, and its abuse can lead to negative individual and social consequences. Many of alcohol's neurobehavioral effects result from its interaction with GABA signaling within the central nervous system, and in particular with GABA_A receptors. These receptors are ligand-gated and composed of five protein subunits, and their functional properties can vary depending on subunit combinations. Among these possible protein subunits is $\alpha 2$, which is encoded by the *GABRA2* gene, one of the genes most frequently associated with alcohol-related behavior in human studies (Demers et al., 2014).

Some of the earliest evidence for a *GABRA2*-alcohol association was with respect to alcohol dependent individuals. In a sample of individuals with a family history of alcohol dependence, Edenberg et al. (2004) found 31 single nucleotide polymorphisms (SNPs) within the *GABRA2* gene to be associated with dependence. Furthermore, Soyka et al. (2008) found an association between a single, 8-SNP haplotype in *GABRA2* and dependence between treatment-seeking dependent and non-dependent individuals. These results support a relationship between *GABRA2* polymorphisms and chronic alcohol consumption.

GABRA2 has also been associated with alcohol-related behavior outside of dependence. In a longitudinal study examining drinking frequency, allelic variation at 6 SNPs within *GABRA2* was associated with a greater increase in drinking frequency during the transition from adolescence to adulthood (Dick et al., 2014). Additionally, several studies have found associations between single or multiple *GABRA2* SNPs and the subjective effects of acutely administered alcohol. These include increased stimulation and euphoria (Arias et al., 2014), increased happiness and vigor (Haughey et al., 2008), and decreased negative effects (Uhart et al., 2013) following oral administration, as well as increased stimulation and liking following IV administration (Haughey et al., 2008). Taken as a whole, the evidence points to a significant relationship between *GABRA2* and alcohol-related behavior, including both chronic use and acute subjective effects.

As manipulation of *GABRA2* in humans is not currently feasible, non-human animals can be extremely useful, and studies with rodents also support the relationship between *GABRA2* and alcohol. Alterations in $\alpha 2$ expression have been found following selection for alcohol consumption and preference, with greater $\alpha 2$ mRNA expression in the nucleus accumbens (NAc) of HAP vs LAP mice (Boehm, unpublished) and greater $\alpha 2$ protein expression in the CeA of P vs NP rats (Liu et al., 2011). That these rodents were alcohol-naïve suggests that $\alpha 2$ alterations might play a role in alcohol consumption or preference.

Experimental manipulation of $\alpha 2$ expression also supports its role in alcohol-related behavior. Boehm II et al. (2004) found that $\alpha 2$ knock-out mice had a reduction in the duration of alcohol-induced loss-of-righting reflex (LORR), a measure of alcohol's

hypnotic effects. Furthermore, $\alpha 2$ knock-out mice showed a reduction in alcohol consumption and preference, but this was only the case in females. However, their (non-significant) reduced preference for bitter taste (quinine) may also have contributed to this effect. Although not part of this study, $\alpha 2$ knock-out mice have also been found to show a reduction in conditioned taste aversion (CTA) to alcohol (Blednov, unpublished). In a more refined approach to assessing the role of $\alpha 2$ subunits in alcohol-induced behavioral responses, Blednov et al. (2011) used knock-in mice containing point mutations at specific amino acids which yielded functional, but alcohol-insensitive $\alpha 2$ -containing GABA_A receptors. These knock-in mice showed both decreases and increases in alcohol consumption and preference depending on the paradigm (2-bottle choice, 1-bottle DID, or 2-bottle DID), access schedule (continuous or limited) and sex. Overall, there was no alcohol-drinking assessment that yielded a lack of effect in knock-in mice. In addition, these mice showed an increase in the duration of alcohol-induced LORR and a decrease in alcohol-induced CTA and locomotion. While the results of these studies suggest a role of $\alpha 2$ -containing receptors in multiple alcohol-related behaviors, the implications are not completely clear as there were differences in effects based on study, paradigm, and sex. Furthermore, an effect of compensatory structural or functional changes within these congenital knock-out/in mice cannot be ruled out.

1.2 Anxiety's Mediation of GABRA2 & Alcohol-Related Behavior

Both human and rodent data suggest, relatively consistently, a relationship between *GABRA2* and alcohol-related behavior. However, this relationship may not always be direct, as human studies indicate that there may be important mediating factors, including anxiety and impulsiveness (Enoch et al., 2006; Villafuerte et al., 2013),

with anxiety being of particular interest for this study. Enoch et al. (2006) used two distinct populations, Finnish and Plains American Indian, to investigate a potential *GABRA2*-alcohol-anxiety relationship. *GABRA2* SNPs were associated with alcoholism in both populations, replicating previous findings. Furthermore, risk (of alcohol dependence)-conferring haplotype configurations within *GABRA2* were associated with harm avoidance (a measure of trait anxiety) but not associated with alcoholism. Harm avoidance was higher in alcoholics, and haplotype frequencies varied with level of harm avoidance, but only for alcoholics. These results indicate that some of the relationship between *GABRA2* haplotype and alcoholism can be revealed when anxiety is taken into consideration. Additionally, it has been suggested that anxiety may partly explain the discrepancy between the ability of high- and low-risk *GABRA2* alleles to differentiate the effects of various psychosocial treatments in alcoholics (Bauer et al., 2007).

1.3 *GABRA2* & Anxiety-Related Behavior

Given the prevalence and clinical relevance of comorbidity of anxiety and alcohol use disorders (J. P. Smith & Randall, 2012), as well as the significant role that GABA_A receptors play in these disorders, it would not be surprising if *GABRA2* played a role in both, and possibly even their overlap. Studies in rodents with global $\alpha 2$ expression manipulation tend to support its role in anxiety, but results aren't unequivocal. $\alpha 2$ knock-out mice have been found to display more risk assessment and longer latencies to explore a novel environment during free-choice exploration, as well as less time in light and more time in the tunnel during a light/dark (L/D) choice situation (Koester et al., 2013). In addition, mice with alcohol-insensitive $\alpha 2$ -containing receptors have displayed a reduced

percentage of open arm entries in the elevated plus maze (EPM) (Blednov et al., 2011). These results indicate a role for $\alpha 2$ expression in basal anxiety.

With respect to drug-induced anxiolysis, the results are mixed. Boehm II et al. (2004) found no significant difference in alcohol's anxiolytic effects in the EPM in $\alpha 2$ knock-out mice, and the same was true of mice with alcohol-insensitive, $\alpha 2$ -containing receptors (Blednov et al., 2011). However, when looking at studies with mutant mice containing point mutations rendering GABA_A receptors containing specific subunits ($\alpha 1$, $\alpha 2$, or $\alpha 3$) insensitive to modulation by benzodiazepines, the evidence for an $\alpha 2$ -anxiolysis relationship is strong. Low et al. (2000) found that only $\alpha 2$ mutant mice showed no effect of diazepam on time in light in the L/D test and open arm time and entries in the EPM. Furthermore, only $\alpha 2$ mutant mice have shown a consistent lack of effect of diazepam and chlordiazepoxide in EPM and fear-potentiated startle assessments without affecting locomotor activity (K. S. Smith, Engin, Meloni, & Rudolph, 2012). Benzodiazepines are effective anxiolytics in non-mutated mice, and while multiple GABA_A α -subunits are implicated in a number of benzodiazepines' behavioral effects, it appears that $\alpha 2$ -containing GABA_A receptors play a principal role in mediating their anxiolytic effects. That global $\alpha 2$ expression manipulations in rodents tend to alter the anxiolytic properties of benzodiazepines but not alcohol might suggest a drug-specific relationship between $\alpha 2$ expression and anxiolysis. However, evidence from region-specific studies (discussed below) suggests that targeted investigation of the role of $\alpha 2$ expression in alcohol's anxiolytic effects is warranted.

1.4 GABA_A Involvement in CeA-Mediated Behavior

The amygdala is widely implicated in both anxiety disorders and addiction, and its circuitry is well conserved across species, allowing for meaningful translational research (Janak & Tye, 2015). Based on location and connectivity, the CeA is an important mediator of responses (physiological and behavioral) to alcohol- and stress-related stimuli (Roberto et al., 2012), leading to the possibility of behaviorally-relevant, alcohol-anxiety interactions in this region. With respect to alcohol, acute administration in rats or mice increases c-Fos expression in the CeA (McBride, 2002), and CeA lesions have been shown to reduce alcohol intake in a 2-bottle choice procedure (Moller, Wiklund, Sommer, Thorsell, & Heilig, 1997). In rats, infusion of a GABA_A antagonist (SR5531) into the CeA has been shown to decrease responding for alcohol in an operant choice task at lower concentrations than infusion into other brain regions (Hyytia & Koob, 1995); infusion of a selective, $\alpha 1$ GABA_A benzodiazepine mixed agonist/antagonist (β CCt) into the CeA has been shown to decrease responding for alcohol, with and without sucrose (Foster et al., 2004); and infusion of a GABA_A agonist (muscimol) into the CeA has been shown to substitute for alcohol in a discrimination task (Hodge & Cox, 1998). With respect to anxiety, lesions of the CeA in rats have been shown to abolish the anxiogenic effects of restraint stress in the EPM (Moller et al., 1997). Furthermore, rats infused with muscimol or midazolam in the CeA have shown a reduction in a number of anxiety-related behaviors in the EPM, and these effects were not present after infusions in the basolateral amygdala (Carvalho, Moreira, Zanoveli, & Brandao, 2012; Moreira, Masson, Carvalho, & Brandao, 2007). These results indicate a role for the CeA broadly, and GABA_A receptors within the CeA specifically, in

subjective and behavioral responses to alcohol as well as in the exhibition of anxiety-related behaviors.

Alcohol- and anxiety-related responses/behaviors also appear to interact within the CeA based on the following studies assessing CeA-mediated anxiety-related behaviors in the context of prior alcohol exposure. Sharko, Kaigler, Fadel, and Wilson (2013) split rats into high- and low-drinking groups after multiple, limited-access alcohol sessions and found an increase in the percent time in open arms in the EPM in the high-drinking group as well as higher c-Fos activation in the CeA, which was correlated with both alcohol intake and percent time in open arms. Rassnick, Heinrichs, Britton, and Koob (1993) infused a corticotropin-releasing factor (CRF) antagonist into the CeA (and intracerebroventricularly) and assessed anxiety in the EPM after withdrawal from either an alcohol- or sucrose-based liquid diet. CRF antagonism was found to be anxiolytic, but only after infusion into the CeA, and only in alcohol-dependent rats. The fact that the aforementioned effects were only found in animals with prior (high) alcohol experience suggests the possibility that alcohol-related alterations in the CeA are relevant to expression of anxiety-related behavior.

1.5 GABRA2 & the CeA

GABRA2 is expressed at relevant levels in the CeA of multiple species. There is moderate to strong expression of $\alpha 2$ protein in the rat CeA (Pirker, Schwarzer, Wieselthaler, Sieghart, & Sperk, 2000) and $\alpha 2$ mRNA and protein expression in the mouse CeA (Hortnagl et al., 2013), and concerning GABA_A subunit distribution in the (extended) amygdala, $\alpha 2$ protein expression is most prominent in the CeA (and NAc, BNST) while $\alpha 1$ protein expression is most prominent in the lateral amygdala (and VP).

With respect to alcohol, post-mortem analysis of CeA tissue from humans has revealed decreases in mRNA expression of a number of glutamate receptor subunits, but only one GABA receptor subunit, $\alpha 2$, in alcoholics compared to controls (Jin et al., 2014).

Additionally, both alcohol-naïve and alcohol-exposed P rats have been shown to have increased expression of $\alpha 2$ (and $\alpha 1$) protein compared to NP rats in the CeA (Liu et al., 2011). Although these human and rat results differ in direction and expression measure, they both indicate a relationship between $\alpha 2$ expression in the CeA and a chronic alcohol consumption phenotype. With respect to anxiety, diazepam-modulated inhibitory post-synaptic currents have been shown to be mediated by $\alpha 2$ -containing receptors in the CeA and $\alpha 1$ - and $\alpha 2$ -containing receptors in the lateral/basolateral amygdala in mutant mice with diazepam-insensitive $\alpha 1$ -, $\alpha 2$ -, or $\alpha 3$ -containing GABA_A receptors (Marowsky, Fritschy, & Vogt, 2004). This goes beyond the effects of the lesioning of, and GABA_A agonism within, the CeA on anxiety to provide evidence for the importance of $\alpha 2$ subunits in the CeA in regulating benzodiazepine-induced inhibition and related anxiolysis.

In what appears to be the only study with direct manipulation of $\alpha 2$ expression specifically in the CeA including an assessment of behavior, Liu et al. (2011) infused P rats with virally-mediated, small interfering RNA (siRNA) targeting either $\alpha 1$ or $\alpha 2$ subunit expression in the CeA, NAc, or ventral pallidum (VP) and assessed them on an operant, multiple-scheduled-access procedure. siRNA infusions resulted in a drastic reduction from baseline responding for alcohol (but not sucrose) when targeting $\alpha 2$ in the CeA and when targeting $\alpha 1$ in the VP, providing strong evidence for subunit-specific, subregion-specific effects within the extended amygdala. Interestingly, alcohol and

sucrose responding were both lower on the first day of assessment post-surgery; however, while sucrose responding returned immediately thereafter, alcohol responding slowly returned to baseline over the course of several days, which could suggest a stress-alcohol- $\alpha 2$ interaction within the CeA.

In summary, *GABRA2* appears to play a role in both alcohol- and anxiety-related behavior. The CeA is an important substrate for the mediation of these behaviors, and $\alpha 2$ -containing GABA_A receptors within this region appear to play an important role in each. The prospect that this subunit's expression within the CeA might play a role in their interaction (alcohol-induced anxiolysis) is intriguing. However, there appear to have been no studies manipulating $\alpha 2$ expression exclusively in the CeA and assessing its effects on basal anxiety and alcohol-induced anxiolysis. The current study sought to fill this gap.

1.6 Specific Aims

1. *Determine the appropriate parameters and dosage for assessing anxiety and alcohol's anxiolytic effects in the elevated plus maze in C57BL/6J mice.*
2. *Assess the effects of virally-mediated, *Gabra2* knock-down in the central nucleus of the amygdala on basal anxiety and alcohol's anxiolytic effects in the elevated plus maze in C57BL/6J mice.*

CHAPTER 2 EFFECT OF CENTRAL AMYGDALA *GABRA2* EXPRESSION ON ANXIETY AND ALCOHOL'S ANXIOLYTIC CAPACITY

2.1 Materials and Methods

2.1.1 General Design

In brief, male B6 mice in Aim 1 were assessed in the EPM naïve or following IP injection of 0, 0.75, 1.0, or 1.5 g/kg ethanol to determine the appropriate parameters for subsequent assessment with viral manipulation. Male B6 mice in Aim 2 underwent surgery for bilateral infusion of GFP-tagged lentivirus targeting *Gabra2* or a scramble control lentivirus into the CeA. Following 12-13 days of recovery, mice were assessed in the EPM naïve or following IP injection of 0, 0.75, or 1.5 g/kg ethanol. After assessment, brains were extracted and sectioned through the CeA. Finally, GFP was quantified, the CeA was collected via laser microdissection, and $\alpha 2$ protein was quantified via ELISA.

2.1.2 Subjects

For Aim 1, a total of 60 male B6 mice were obtained from Jackson Labs (Bar Harbor, ME) at 8 weeks of age. For Aim 2, a total of 104 male B6 mice were obtained from Jackson Labs at 8 weeks of age, and of these, 6 mice died during or after surgery and 2 mice were not used due to cagemate death, leaving a total of 96 mice used for experimentation. Mice were 9-10 weeks of age at the time of surgery and 11-12 weeks of age at the time of behavioral assessment. All mice had ad libitum access to food and water for the duration of the experiment and were pair-housed (except for 5-6 days during

recovery from surgery) under a 12-hour, reverse light/dark schedule, with lights off at 8:00 am. Procedures were approved by the IUPUI School of Science Institutional Animal Care and Use Committee and conformed to the Guide for the Care and Use of Laboratory Animals (The National Academic Press, 2003).

2.1.3. Surgery

Mice were anesthetized with a 0.01 ml/g IP injection of a ketamine/xylazine cocktail (10:1:89 ratio of ketamine (100 mg/ml), xylazine (100 mg/ml), and 0.9% sterile saline, respectively). Once anesthetized, mice had their heads shaved and were placed in a Kopf stereotaxic alignment system (Tujunga, CA). Mice's eyes were kept moist with commercially available tear ointment. A small incision was made from approximately bregma to lambda, and the skull was sterilized with 70% alcohol and Nolvasan disinfectant. For accuracy, the distance between bregma and lambda was measured and divided by 4.21 mm, the published average distance between lambda and bregma for B6 mice (Franklin and Paxinos, 1997), to create a scaling factor for each mouse. The targeted coordinates for the CeA (-1.20 mm A/P, ± 2.80 mm M/L, and -4.85 mm D/V) were adjusted by the scaling factor, and one hole was drilled in the skull in each hemisphere above the CeA. A microinjection syringe (Hamilton Co., Reno, NV) was twice lowered into the CeA to bilaterally infuse a GFP-tagged lentivirus targeting *Gabra2* (Virus) via short hairpin RNA (shRNA) or a GFP-tagged scramble control lentivirus (Control) (Sigma-Aldrich, Inc., St. Louis, MO). An illustration and description of RNA interference via shRNA are given in **Figure 1**. Viruses were diluted 1:1 with 0.9% sterile saline, as exploratory surgery indicated this reduces extra-regional incorporation. This produced final titers of 2.4×10^9 and 2.2×10^9 TU/ml for Virus and Control lentiviruses,

respectively. 0.15 μ l of virus was infused at a rate of 0.1 μ l/min and allowed to diffuse for an additional 10 minutes. The microinjector was raised 0.05 mm between infusion and diffusion, as exploratory surgery indicated this reduces residual virus on the microinjector tip. Between infusions, the syringe pump (World Precision Instruments, Sarasota, FL) was run until virus was visible in order to confirm patency. Following infusions, incisions were sealed with VetBond skin glue (3M Animal Care, St. Paul, MN), and mice received 0.01 ml/g SC injections of buprenorphine (0.03 mg/ml) and rimadyl (5 mg/ml) for analgesia and anti-inflammation, respectively. Following surgery, mice were single-housed in a biohazard-approved, filter-top cage for 5-6 days of recovery and subsequently pair-housed with their original cagemate, with 4 exceptions. 2 mice were housed with an age-matched, surgery-naïve mouse, and 2 mice were housed with each other due to the death of their original cagemate during surgery.

2.1.4. EPM Assessment

The testing room was separated by a canvas curtain into two sections, one under red light (the same conditions as the vivarium) and the other, containing the EPM, under red light plus dim ambient lighting. The EPM (Med Associates, Inc., St. Albans, VT) was black in color, was 74.5 cm above the ground, and had arms measuring 76.0 cm in length. The closed arms had walls measuring 20.5 cm, and the open arms had no walls, but a slight lip on the sides but not ends. The luminance across testing was 26 ± 2 , 5 ± 1 , and 33 ± 2 lux for Aim 1 and 43 ± 1 , 6 ± 1 , and 54 ± 4 lux for Aim 2, for the center, closed arms, and open arms, respectively. Luminance was increased slightly in Aim 2 in order to enhance the EPM's anxiogenic nature.

Previous work in our lab has shown viral incorporation and significant effects on alcohol consumption at 2 weeks post-infusion (Boehm, unpublished). Therefore, mice were assessed in the EPM following 12-13 days of recovery from surgery. Mice were transported to the testing room 5 minutes prior to lights off (8:00 am) and allowed to habituate in the non-maze portion of the testing room for at least 1 hour prior to assessment. Prior to placement on the EPM, each mouse was weighed and placed in an empty holding cage for 10 minutes. During this time, injected mice were taken just outside the testing room for 30 seconds to receive an IP injection of either 0, 0.75, or 1.5 g/kg of 20% ethanol in 0.9% sterile saline (Aim 1 also included a 1.0 g/kg dose). All injections took place 10 minutes prior to placement in the EPM. Mice were placed in the center of the EPM, facing an open arm, and allowed to explore for 5 minutes while being video-recorded by a camera mounted directly overhead. All condition-dose combinations were represented on each testing day except for 2 instances.

Following assessment, the number of fecal boli were counted, and the EPM was cleaned with Clidox-S. Retro-orbital sinus blood samples were collected in Aim 1 from a subset of mice in each injection condition (n 's = 8) to determine BEC resulting from IP administration of ethanol. Videos were later scored for entries and time spent in center, closed arms, and open arms as well as the number of head dips made. Mice were considered to have entered an arm when all 4 paws transitioned from outside to inside an arm and considered to be in an arm as long as all 4 paws remained inside. Mice were considered to have made a head dip when the shoulders moved forward and the head was facing downward over the edge of the EPM. Head dips were differentiated as protected when initiated from outside of an open arm.

2.1.5. Histology

Approximately 2-4 hours after assessment in the EPM, mice's brain were extracted, flash frozen in 2-methylbutane at $< -40^{\circ}$ C, and stored at -80° C until processed. Brains were sliced in a cryostat at -16° C. Serial sections through the CeA at $35\ \mu\text{m}$ thickness were mounted to PEN-Membrane slides, dehydrated with ascending concentrations of ethanol (70% - 5 sec, 95% - 5 sec, 100% - 20 sec), and stored at -80° C until microdissection. For sections containing the CeA, tissue was laser microdissected using the following technique. The area expressing GFP was encircled under 5x magnification with fluorescence using laser microdissection software (Leica Biosystems Inc., Buffalo Grove, IL), which calculates area in nm^2 . Bilateral CeA was then laser microdissected under 5x magnification in bright field while referencing a mouse brain atlas (Paxinos & Franklin, 1997), and the tissue was captured in 1x RIPA buffer (1x RIPA buffer with $100\ \mu\text{l/ml}$ of 10x PI and $10\ \mu\text{l/ml}$ of 20x PMSF). The remaining area expressing GFP was again encircled and calculated under 5x magnification with fluorescence. Bilateral CeA tissue was collected from all sections whether or not they expressed GFP fluorescence. The pre- and post-microdissection GFP area calculations were used to estimate the amount of GFP expressing tissue within and without the CeA. Tissue samples were homogenized manually, and total protein concentration was calculated for each. Samples were assessed via ELISA (MyBioSource, Inc., San Diego, CA), in triplicate and with $60\ \mu\text{g}$ of total protein loaded per sample, for quantification of $\alpha 2$ protein concentration according to the manufacturer's suggested procedures. An illustration and description of the sandwich ELISA assay are given in **Figure 2**. A total of 4 ELISA assays were needed to assess all samples, with all 3 samples from each brain

run within one assay, and the mean value of samples from each brain was used as the $\alpha 2$ protein concentration.

2.1.6. Statistical Analysis

Statistical significance for all analyses was set at $p < .05$. For Aim 1, confirmation of the anxiogenic nature of the EPM was assessed using only naïve mice. Both a single-sample t-test comparing % of entries in open arms to 50% and a paired-samples t-test comparing time spent in the open vs. closed arms were run. To determine appropriate doses for future assessment in Aim 2, the following measures were submitted to a one-way ANOVA with dose (0, 0.75, 1.0, 1.5 g/kg) as the lone factor, using only mice receiving injection: % of entries in open arms, time spent in open arms, and total arm entries. Each ANOVA was followed by Dunnett's test using the 0 g/kg dose as the reference.

For Aim 2 histological analyses, only mice expressing GFP in the CeA were included, and mice with total protein and/or $\alpha 2$ protein concentration values ≥ 3 SD from the overall mean were excluded (see **Table 1** for n's). Consistency of ELISA output was assessed using a one-way ANOVA on $\alpha 2$ protein concentrations, with ELISA pass as the lone factor, for Control mice only. Because there was a significant effect of ELISA pass on $\alpha 2$ protein concentrations, $F(3,27) = 3.84$, $p = .021$, mean $\alpha 2$ protein concentrations for all mice were normalized via z-score transformations based on distributions within their respective ELISA pass (**Fig. 5A**). Normalized (z-score) values were used to represent $\alpha 2$ protein concentrations for all subsequent analyses. A one-tailed, independent samples t-test was used to test the prediction that $\alpha 2$ protein concentrations would be lower in the Virus relative to the Control condition.

For Aim 2 behavioral analyses, only mice expressing GFP in the CeA were included, and mice with ≤ 1 open arm entry or an environmental confound (noise during assessment) were excluded (see **Table 1** for n's). To limit the influence of general activity levels on measures of arm entries, the % of entries in open arms $((\text{open}/\text{total}) \times 100)$ was calculated for each mouse. In addition, % of head dips protected $((\text{protected}/\text{total}) \times 100)$ was calculated for each mouse. Lastly, latency to first arm entry and fecal boli were largely unrelated to other EPM measures and were not analyzed further (**Table 2**). The following behavioral measures were used as dependent variables for analyses: total arm entries; % of entries in open arms; total and average time spent in the center, closed arms, and open arms; head dips; and % of head dips protected. To assess the effect of viral condition on basal anxiety, these measures were each submitted to an independent-samples t-test comparing Control to Virus using only naïve mice. To assess both alcohol's anxiolytic capacity and potential augmentation by viral condition, these measures were each submitted to a two-way ANOVA with viral condition and dose as factors using only mice receiving 0, 0.75, or 1.5 g/kg ethanol injections. Main effects of dose were followed by Dunnett's test using the 0 g/kg dose as the reference, and main effects of viral condition were followed by Bonferroni-corrected t-tests comparing Control to Virus within each dose.

For Aim 2 brain-behavior correlations, all mice were included, with the following exceptions: mice with total protein and/or $\alpha 2$ protein concentration values ≥ 3 SD from the overall mean and mice with ≤ 1 open arm entry or an environmental confound (see **Table 1** for n's). The relationships between $\alpha 2$ protein concentrations and behavioral

measures in the EPM, as well as the relationships between the various EPM behavioral measures, were assessed using Pearson correlations.

2.2 Results

2.2.1. Aim 1

The anxiogenic nature of the EPM was confirmed for this experiment by mice making significantly fewer open arm entries than would be expected by chance (50%), $t(11) = -4.15, p = .002$ and by mice spending significantly less time in open vs. closed arms, $t(11) = 6.35, p < .001$ (data not shown). The anxiolytic capacity of alcohol was confirmed by a significant increase in the % of entries in open arms, $F(3, 38) = 4.57, p = .008$, with Dunnett's test indicating that all doses differed significantly from 0 g/kg (p 's $< .048$) (**Fig. 3A**); however, alcohol was not found to affect either time spent in open arms, $F(3, 38) = 1.63, p = .199$ or total arm entries, $F(3, 38) = 2.54, p = .071$ (**Figs. 3B** and **3C**, respectively). Based on these results, the 0.75 and 1.5 g/kg doses were chosen for use in Aim 2. These doses produced mean BEC values of 74.58 and 151.62 mg%, respectively.

2.2.2. Histology

Representative images showing both successful and unsuccessful viral incorporation with reference to a mouse brain atlas (Paxinos & Franklin, 1997) are shown in **Figure 4**. As shown in **Figure 5B**, a one-tailed, independent-samples t-test revealed a decrease in normalized $\alpha 2$ protein concentrations for Virus mice relative to Control mice, $t(62) = 1.81, p = .038$, thus confirming the effectiveness of the virus in reducing expression of the target protein. An assessment of the potential impact of dose on these

normalized protein concentrations via two-way ANOVA yielded no significant effect of dose or dose x condition interaction (p 's > .470, data not shown).

2.2.3. EPM Assessment

The anxiogenic nature of the EPM was confirmed for this experiment by mice making significantly fewer open arm entries than would be expected by chance (50%), $t(10) = -4.97$, $p = .001$ and by mice spending significantly less time in open vs. closed arms, $t(10) = 4.61$, $p = .001$ (**Figs. 6B** and **7B/C**, respectively). Regarding basal anxiety in naïve mice, there were no significant differences between Control mice and Virus mice on any behavioral measure (p 's > .376) (**Figs. 6**, **7**, and **8**). Regarding alcohol's anxiolytic capacity in mice receiving 0, 0.75, or 1.5 g/kg ethanol injections, two-way ANOVA's yielded a significant main effect of viral condition for average time spent in closed arms, $F(1, 44) = 5.23$, $p = .027$, with average time reduced in Virus mice relative to Control mice (**Fig. 7E**). Post hoc independent-samples t-tests yielded no significant differences between Virus mice and Control mice at any given dose (p 's > .104). In contrast to viral condition, a significant main effect of dose was found for several behavioral measures including % of entries in open arms, $F(2, 44) = 9.73$, $p < .001$, time spent in center, $F(2, 44) = 7.10$, $p = .002$, time spent in open arms, $F(2, 44) = 3.60$, $p = .036$, head dips, $F(2, 44) = 7.20$, $p = .002$, and % of head dips protected, $F(2, 44) = 4.98$, $p = .011$ (**Figs. 6B**, **7A**, **7C**, **8A**, and **8B**, respectively). Follow-up Dunnett's tests using the 0 g/kg dose as the reference revealed that both the 0.75 and 1.5 g/kg doses increased % of entries in open arms ($p = .03$ and $p < .001$, respectively). However, it was only the 1.5 g/kg dose that decreased time spent in center ($p = .001$), increased time spent in open arms ($p = .022$), increased head dips ($p = .002$), and decreased % of head dips

protected ($p = .011$). There were no significant interactions between viral condition and dose for any behavioral measure (p 's $> .186$) (**Figs.6, 7, and 8**).

2.2.4. Brain-Behavior Correlations

Pearson correlations were run within each dose (and naïve) to further explore the potential relationship between CeA $\alpha 2$ protein concentration and behavior. For naïve mice, normalized $\alpha 2$ protein concentration was significantly positively correlated with time spent in closed arms, $r(22) = .449$, $p = .036$ and significantly negatively correlated with both time spent in open arms overall $r(22) = -.519$, $p = .013$ and on average $r(22) = -.463$, $p = .03$ (**Figs. 9A, 9B, and 9C**, respectively). There were no significant correlations between these measures in mice receiving 0, 0.75, or 1.5 g/kg ethanol (p 's $> .068$) (**Table 2**).

CHAPTER 3 DISCUSSION

3.1. EPM & Alcohol's Anxiolytic Capacity

The current EPM parameters were confirmed to be anxiogenic in completely naïve mice (Aim 1) as well as mice having undergone surgery (Aim 2). As expected, alcohol was found to be an effective anxiolytic in both Aims; however, this effect was more pronounced in Aim 2. In addition to differential histories, naive (Aim 1) vs. surgery (Aim 2), EPM luminance was increased for Aim 2, and each aim was run as a separate experiment, so a direct comparison of alcohol's anxiolytic capacity between them may not be appropriate. However, these parameters did provide an appropriate context for the assessment of the effect of CeA *Gabra2* expression on basal anxiety and alcohol's anxiolytic capacity.

3.2. Virally-Mediated *Gabra2* Knock-Down

Several mice in this experiment exhibited no GFP expression, yet all mice had visible needle track marks in or near the CeA. Lack of expression could have been due to clogged/blocked microinjector tips or ineffective viral incorporation. Although, considering the majority of mice exhibited GFP expression, ineffective incorporation seems much less likely. Quantification of the area of GFP expression within the CeA was intended to provide an indirect measure of the amount/degree of viral incorporation. While there should have been no relationship between CeA GFP expression and $\alpha 2$

protein concentration in Control mice, a negative relationship was expected for Virus mice, as more incorporation should have produced a greater knock-down. On the contrary, there was a positive correlation between CeA GFP expression and $\alpha 2$ protein concentration in Virus mice. Both the inability to measure GFP expression in three dimensions and the inability to account for the intensity of fluorescence could have been responsible for this unexpected result. Therefore, GFP expression was useful as an indicator of viral incorporation but not as a measure of amount/degree of incorporation. Overall, lentivirus targeting *Gabra2* was effective in reducing $\alpha 2$ protein concentrations relative to a scramble control lentivirus.

3.3. CeA *Gabra2* Expression & Basal Anxiety

The current study found no significant difference on any behavioral measure between naïve Control and Virus mice. However, when looking at naïve mice collectively, there was a significant relationship between CeA $\alpha 2$ protein concentration and multiple behavioral measures. This discrepancy could be accounted for by the fact that the statistically significant reduction in $\alpha 2$ protein concentrations seen for Virus mice relative to Control mice may not have been behaviorally significant. Taken together, the results from naïve mice support a role for CeA *Gabra2* expression in basal anxiety, specifically a positive relationship between CeA $\alpha 2$ protein concentration and anxiety-like behavior. In addition to the significant correlations, the relationships between $\alpha 2$ protein concentration and behavior were all consistently in this direction (except average time spent in closed arms), and although not statistically significant, the difference in means between Control mice and Virus mice for these behaviors were also consistently in the same direction (except head dips).

The current demonstration of a *Gabra2*-anxiety relationship is in line with several previous studies. In humans, haplotype configurations within *GABRA2* have been found to be associated with harm avoidance, the tendency to be anxious and fearful (Enoch et al., 2006). Of course these haplotypes don't indicate a specific change in $\alpha 2$ protein concentration, and these individuals likely had varied environmental histories. Similarly, compared to wild-type mice, $\alpha 2$ knock-out mice show alterations in anxiety-like behavior, specifically increased risk-assessment in and latency to explore a novel environment as well as decreased time in light in a L/D test (Koester et al., 2013). That this negative relationship between *Gabra2* expression and anxiety-like behavior is opposite in direction to the current results could be due to a number of factors including the congenital and global nature of the *Gabra2* manipulation, the type of assessment used, and the different strain of mouse used (129X1/SvJ). Nevertheless, it supports a *Gabra2*-anxiety relationship in animals with controlled environmental histories. The current results also implicate $\alpha 2$ protein concentration specifically in the CeA in affecting anxiety-like behavior. Additional support for this relationship comes from the observation that naïve P rats have elevated CeA $\alpha 2$ protein concentrations compared to NP rats (Liu et al., 2011) and also display greater anxiety-like behavior in the EPM (R. B. Stewart, Gatto, Lumeng, Li, & Murphy, 1993). Although there are likely several structural/functional differences between P and NP rats, the difference in CeA $\alpha 2$ protein concentrations is significant (~100% increase) and the direction of its relationship to behavior is the same as in the current study. Finally, while the current results can't rule out other brain-related factors as contributing to the observed behavioral differences, the fact that B6 mice are inbred, resulting in relatively homogeneous brain structure/function,

and that viral manipulation of the CeA produced significant alterations in $\alpha 2$ protein concentrations specifically in this region strengthens the interpretation that CeA *Gabra2* expression is positively related to basal anxiety.

3.4. CeA *Gabra2* Expression & Alcohol's Anxiolytic Capacity

Although alcohol was clearly found to be anxiolytic in this study, affecting a number of EPM measures in a predictable fashion, there was no interaction between viral condition and dose on any measure and only a main effect of viral condition across doses on one measure, average time spent in closed arms. That this behavior alone, and not in conjunction with behaviors typically reported in EPM studies (e.g. time spent in open arms and % of entries in open arms), was affected by viral condition decreases confidence in the interpretation that a reduction in $\alpha 2$ protein concentration is anxiogenic across these doses. In fact, mean group differences for this behavior are most pronounced at the 0.75 and 1.5 g/kg doses, yet correlations indicate a null relationship and a weak relationship in the opposite direction for these doses, respectively. Furthermore, increases in average time spent in closed arms could be explained by behaviors not necessarily related to anxiety, like increases in grooming. Finally, there were no significant relationships between $\alpha 2$ protein concentration and behavioral measures within any dose. Taken together, these results suggest that alterations in CeA *Gabra2* expression do not affect alcohol's anxiolytic capacity.

A lack of effect of *Gabra2* expression on alcohol's anxiolytic capacity is consistent with a lack of effect seen in both $\alpha 2$ knock-out mice (Boehm II et al., 2004) and knock-in mice containing alcohol-insensitive $\alpha 2$ -containing GABA_A receptors (Blednov et al., 2011). However, this is in contrast to the role of *Gabra2* expression in

mediating the anxiolytic actions of benzodiazepines, as mice with benzodiazepine-insensitive $\alpha 2$ -containing GABA_A receptors fail to show an anxiolytic response to these drugs in a number of behavioral assessments (Low et al., 2000; K. S. Smith et al., 2012). In addition, while modulation of inhibitory post-synaptic currents by diazepam in the CeA appears to be mediated by $\alpha 2$ -containing GABA_A receptors (Marowsky et al., 2004), the results of the current study do not suggest a role for CeA $\alpha 2$ subunits in alcohol's pharmacologic actions. This difference could be related to, among other things, alcohol's enhancement of inhibitory GABAergic transmission in the CeA via both pre- and post-synaptic mechanisms (Roberto, Madamba, Moore, Tallent, & Siggins, 2003) or alcohol's interactions with anxiogenic CRF and anxiolytic neuropeptide Y (NPY) signaling in the CeA (Gilpin, Herman, & Roberto, 2015). Taken together, these studies suggest a drug- or drug class- specific relationship between $\alpha 2$ subunits and anxiolysis.

3.5. Conclusions, Limitations, & Future Directions

The current study supports the idea that CeA $\alpha 2$ protein expression is positively related to anxiety. Enoch et al. (2006) found that risk (of alcohol dependence)-conferring *GABRA2* haplotypes were associated with trait anxiety, which was increased in alcoholics, and one could speculate that these individuals had increased CeA *GABRA2* expression. However, post-mortem analysis of CeA tissue from alcoholics has revealed decreased *GABRA2* mRNA expression (Jin et al., 2014). These findings can be reconciled when considering that a substantial change in CeA $\alpha 2$ protein expression and corresponding change in basal anxiety in either direction could lead to increases in alcohol use via different routes. Increased anxiety, associated with an increase in CeA $\alpha 2$ protein expression, would lead to an increase in the opportunity for alcohol to provide

relief and acquire negatively-reinforcing properties. This negative-reinforcement route is direct and fairly intuitive, and it is consistent with several models of human alcoholism (J. P. Smith & Randall, 2012). On the other hand, decreased anxiety, associated with a decrease in CeA $\alpha 2$ protein expression, might lead to alcohol use through a more indirect route via a relationship with externalizing behaviors, which are commonly found to be comorbid with substance use disorders (SUDs) (Hofmann, Richey, Kashdan, & McKnight, 2009). In a direct assessment of the relationship between anxiety, externalizing problems, and SUDs, Hofmann, Richey, Kashdan, and McKnight (2009) found that while having either an anxiety disorder or externalizing problems increased the probability of having a SUD, the probability of having a SUD for individuals with externalizing problems was decreased for those with a comorbid anxiety disorder compared to those without. Thus anxiety can moderate the relationship between externalizing problems and substance use, with decreased anxiety being associated with greater substance use in those with externalizing problems.

Interestingly, both *GABRA2* and the amygdala have been associated with externalizing behaviors. With respect to *GABRA2*, a risk (of alcohol dependence)-conferring *GABRA2* genotype has been found to be associated with a greater likelihood of maintaining elevated expression of externalizing behaviors across the transition from adolescence to adulthood (Dick et al., 2009). With respect to the amygdala, preschool externalizing behavior has been shown to predict decreased amygdala volume in adolescence in males (Caldwell et al., 2015), and adolescents with conduct disorder have also been shown to have decreased amygdala volume (Wallace et al., 2014). While there is growing evidence that *GABRA2*, anxiety, externalizing behavior, and alcohol use all

share some degree of interrelation, the exact nature of these relationships is not clear. Future studies could further investigate the *GABRA2*-anxiety relationship by assessing (central) amygdala structure/connectivity or activity in response to anxiogenic cues/situations as a function of *GABRA2* genotype in individuals prior to the initiation of alcohol use. In addition, a longitudinal study assessing *GABRA2* genotype in relation to measures of anxiety and externalizing behaviors before and after the initiation of alcohol use might further elucidate the impact and potential interaction of these *GABRA2*-associated phenotypes in relation to alcohol use.

In contrast to basal anxiety, the current study does not support the idea that CeA $\alpha 2$ protein expression is related to alcohol's anxiolytic capacity. However, one limitation of the current study was that it assessed this capacity with only a single acute exposure. Several studies indicate that chronic exposure to alcohol can alter basal GABAergic transmission and response to GABAergic drugs in the CeA (Koob, 2004), and it appears that these alterations might affect anxiety-like behavior. Sharko et al. (2013) assessed anxiety-like behavior in the EPM using rats with water or low- or high-drinking alcohol histories. High-drinking rats had greater CeA c-Fos expression, and CeA c-Fos expression was associated with anxiety-like behavior in alcohol- but not water-drinking animals, suggesting alcohol-induced alterations in the relationship between CeA activity and anxiety. Withdrawal from chronic alcohol exposure can also augment CeA functioning with respect to anxiety, as infusion of a CRF antagonist in the CeA has been shown to be anxiolytic in alcohol-dependent but not non-dependent animals 8 hours post-exposure (Rassnick et al., 1993). Lastly, Liu et al. (2011) found a decrease in alcohol consumption in rats with virally-mediated CeA *Gabra2* knock-down which occurred just

3 days after invasive brain surgery, which could be interpreted as a change in the negatively-reinforcing capacity of alcohol. Given that these rats had 21 days of drinking history prior to surgery, alcohol-induced changes in CeA functioning could play a role in this finding, and it remains to be seen how CeA *Gabra2* manipulation would affect drinking in alcohol-naïve animals.

Taken together, the aforementioned studies provide evidence for alcohol history as being an important factor in the CeA-alcohol-anxiety relationship and provide rationale for future investigations. Surprisingly, it appears that an assessment of the effect of alcohol infusion directly into the CeA on anxiety-like behavior has not been conducted. Performing this assessment in both alcohol-naïve and chronically alcohol-exposed animals would help to clarify the role of the CeA in alcohol's anxiolytic capacity and address potential augmentation of this role based on prior exposure. In addition, replicating the current study in chronically exposed mice would indicate whether or not the relationship between CeA *Gabra2* expression and anxiety-like behavior is still present after a history of alcohol exposure and whether or not a history of alcohol exposure alters the lack of relationship between CeA *Gabra2* expression and alcohol's anxiolytic capacity.

One major limitation of the current study that has yet to be mentioned is that only male mice were used, yet gender differences do exist for many of the behaviors discussed. Females have a higher prevalence of anxiety disorders than males even though age of onset and duration tend not to differ (McLean, Asnaani, Litz, & Hofmann, 2011). In the context of alcohol, females and males with high anxiety sensitivity differ in the degree to which the nature of a stressor impacts the acute effects of consumed alcohol

(Zack, Poulos, Aramakis, Khamba, & MacLeod, 2007). With respect to *GABRA2*, in individuals with a risk (of alcohol dependence)-conferring *GABRA2* genotype, increased levels of positive life events serves as a protective factor for males but not females (Perry et al., 2013). Finally, $\alpha 2$ knock-out and knock-in mice have shown sex-dependent differences in alcohol consumption (Blednov et al., 2011; Boehm II et al., 2004). Given these, and other, gender differences, especially regarding anxiety and alcohol use, an important future study would be to replicate the current experiment in female mice.

In summary, CeA $\alpha 2$ protein expression does not appear to be related to alcohol's anxiolytic capacity but does appear to be positively related to basal anxiety. The CeA may be a region where changes in $\alpha 2$ protein expression resulting from *GABRA2* polymorphisms could affect basal anxiety, which in turn could affect alcohol use through various routes. Investigating the role that these subunits play in GABAergic transmission within the CeA and its effect on efferent signaling at downstream targets in response to anxiogenic or stressful conditions could lead to a better understanding of mechanisms underlying the relationship between anxiety and alcohol use.

REFERENCES

REFERENCES

- Arias, A. J., Covault, J., Feinn, R., Pond, T., Yang, B.-Z., Ge, W., . . . Kranzler, H. R. (2014). A GABRA2 Variant Is Associated with Increased Stimulation and ‘High’ Following Alcohol Administration. *Alcohol and Alcoholism (Oxford, Oxfordshire)*, *49*(1), 1-9. doi:10.1093/alcalc/agt163
- Bauer, L. O., Covault, J., Harel, O., Das, S., Gelernter, J., Anton, R., & Kranzler, H. R. (2007). Variation in GABRA2 Predicts Drinking Behavior in Project MATCH Subjects. *Alcoholism, clinical and experimental research*, *31*(11), 1780-1787. doi:10.1111/j.1530-0277.2007.00517.x
- Blednov, Y. A., Borghese, C. M., McCracken, M. L., Benavidez, J. M., Geil, C. R., Osterdorff-Kahanek, E., . . . Harris, R. A. (2011). Loss of Ethanol Conditioned Taste Aversion and Motor Stimulation in Knockin Mice with Ethanol-Insensitive α 2-Containing GABA(A) Receptors. *The Journal of Pharmacology and Experimental Therapeutics*, *336*(1), 145-154. doi:10.1124/jpet.110.171645
- Boehm II, S. L., Ponomarev, I., Jennings, A. W., Whiting, P. J., Rosahl, T. W., Garrett, E. M., . . . Harris, R. A. (2004). γ -Aminobutyric acid A receptor subunit mutant mice: new perspectives on alcohol actions. *Biochemical Pharmacology*, *68*(8), 1581-1602. doi:http://dx.doi.org/10.1016/j.bcp.2004.07.023
- Caldwell, J. Z., Armstrong, J. M., Hanson, J. L., Sutterer, M. J., Stodola, D. E., Koenigs, M., . . . Davidson, R. J. (2015). Preschool externalizing behavior predicts gender-specific variation in adolescent neural structure. *PLoS One*, *10*(2), e0117453. doi:10.1371/journal.pone.0117453
- Carvalho, M. C., Moreira, C. M., Zanoveli, J. M., & Brandao, M. L. (2012). Central, but not basolateral, amygdala involvement in the anxiolytic-like effects of midazolam in rats in the elevated plus maze. *J Psychopharmacol*, *26*(4), 543-554. doi:10.1177/0269881110389209
- Demers, C. H., Bogdan, R., & Agrawal, A. (2014). The Genetics, Neurogenetics and Pharmacogenetics of Addiction. *Current behavioral neuroscience reports*, *1*(1), 33-44. doi:10.1007/s40473-013-0004-8
- Dick, D. M., Cho, S. B., Latendresse, S. J., Aliev, F., Nurnberger, J. I., Jr., Edenberg, H. J., . . . Kuperman, S. (2014). Genetic influences on alcohol use across stages of development: GABRA2 and longitudinal trajectories of drunkenness from adolescence to young adulthood. *Addict Biol*, *19*(6), 1055-1064. doi:10.1111/adb.12066
- Dick, D. M., Latendresse, S. J., Lansford, J. E., Budde, J. P., Goate, A., Dodge, K. A., . . . Bates, J. E. (2009). Role of GABRA2 in trajectories of externalizing behavior across development and evidence of moderation by parental monitoring. *Arch Gen Psychiatry*, *66*(6), 649-657. doi:10.1001/archgenpsychiatry.2009.48

- Edenberg, H. J., Dick, D. M., Xuei, X., Tian, H., Almasy, L., Bauer, L. O., . . . Begleiter, H. (2004). Variations in GABRA2, encoding the alpha 2 subunit of the GABA(A) receptor, are associated with alcohol dependence and with brain oscillations. *Am J Hum Genet*, 74(4), 705-714. doi:10.1086/383283
- Enoch, M. A., Schwartz, L., Albaugh, B., Virkkunen, M., & Goldman, D. (2006). Dimensional anxiety mediates linkage of GABRA2 haplotypes with alcoholism. *Am J Med Genet B Neuropsychiatr Genet*, 141b(6), 599-607. doi:10.1002/ajmg.b.30336
- Foster, K. L., McKay, P. F., Seyoum, R., Milbourne, D., Yin, W., Sarma, P. V., . . . June, H. L. (2004). GABA(A) and opioid receptors of the central nucleus of the amygdala selectively regulate ethanol-maintained behaviors. *Neuropsychopharmacology*, 29(2), 269-284. doi:10.1038/sj.npp.1300306
- Gilpin, N. W., Herman, M. A., & Roberto, M. (2015). The central amygdala as an integrative hub for anxiety and alcohol use disorders. *Biol Psychiatry*, 77(10), 859-869. doi:10.1016/j.biopsych.2014.09.008
- Haughey, H. M., Ray, L. A., Finan, P., Villanueva, R., Niculescu, M., & Hutchison, K. E. (2008). Human gamma-aminobutyric acid A receptor alpha2 gene moderates the acute effects of alcohol and brain mRNA expression. *Genes Brain Behav*, 7(4), 447-454. doi:10.1111/j.1601-183X.2007.00369.x
- Hodge, C. W., & Cox, A. A. (1998). The discriminative stimulus effects of ethanol are mediated by NMDA and GABA(A) receptors in specific limbic brain regions. *Psychopharmacology (Berl)*, 139(1-2), 95-107.
- Hofmann, S. G., Richey, J. A., Kashdan, T. B., & McKnight, P. E. (2009). Anxiety disorders moderate the association between externalizing problems and substance use disorders: data from the National Comorbidity Survey-Revised. *J Anxiety Disord*, 23(4), 529-534. doi:10.1016/j.janxdis.2008.10.011
- Hortnagl, H., Tasan, R. O., Wieselthaler, A., Kirchmair, E., Sieghart, W., & Sperk, G. (2013). Patterns of mRNA and protein expression for 12 GABAA receptor subunits in the mouse brain. *Neuroscience*, 236, 345-372. doi:10.1016/j.neuroscience.2013.01.008
- Hutvagner, G., & Zamore, P. D. (2002). RNAi: nature abhors a double-strand. *Curr Opin Genet Dev*, 12(2), 225-232.
- Hyytia, P., & Koob, G. F. (1995). GABAA receptor antagonism in the extended amygdala decreases ethanol self-administration in rats. *Eur J Pharmacol*, 283(1-3), 151-159.
- Janak, P. H., & Tye, K. M. (2015). From circuits to behaviour in the amygdala. *Nature*, 517(7534), 284-292. doi:10.1038/nature14188
- Jin, Z., Bhandage, A. K., Bazov, I., Kononenko, O., Bakalkin, G., Korpi, E. R., & Birnir, B. (2014). Expression of specific ionotropic glutamate and GABA-A receptor subunits is decreased in central amygdala of alcoholics. *Front Cell Neurosci*, 8, 288. doi:10.3389/fncel.2014.00288
- Jolliff, K. (2007). Lentiviral shRNA constructs for gene silencing studies. *Drug Discovery & Development*. Retrieved from <http://www.dddmag.com/articles/2007/09/lentiviral-shrna-constructs-gene-silencing-studies> on April 2016

- Koester, C., Rudolph, U., Haenggi, T., Papilloud, A., Fritschy, J. M., & Crestani, F. (2013). Dissecting the role of diazepam-sensitive gamma-aminobutyric acid type A receptors in defensive behavioral reactivity to mild threat. *Pharmacol Biochem Behav*, *103*(3), 541-549. doi:10.1016/j.pbb.2012.10.004
- Koob, G. F. (2004). A role for GABA mechanisms in the motivational effects of alcohol. *Biochem Pharmacol*, *68*(8), 1515-1525. doi:10.1016/j.bcp.2004.07.031
- Liu, J., Yang, A. R., Kelly, T., Puche, A., Esoga, C., June, H. L., Jr., . . . Aurelian, L. (2011). Binge alcohol drinking is associated with GABAA alpha2-regulated Toll-like receptor 4 (TLR4) expression in the central amygdala. *Proc Natl Acad Sci U S A*, *108*(11), 4465-4470. doi:10.1073/pnas.1019020108
- Low, K., Crestani, F., Keist, R., Benke, D., Brunig, I., Benson, J. A., . . . Rudolph, U. (2000). Molecular and neuronal substrate for the selective attenuation of anxiety. *Science*, *290*(5489), 131-134.
- Marowsky, A., Fritschy, J. M., & Vogt, K. E. (2004). Functional mapping of GABA A receptor subtypes in the amygdala. *Eur J Neurosci*, *20*(5), 1281-1289. doi:10.1111/j.1460-9568.2004.03574.x
- McBride, W. J. (2002). Central nucleus of the amygdala and the effects of alcohol and alcohol-drinking behavior in rodents. *Pharmacol Biochem Behav*, *71*(3), 509-515.
- McLean, C. P., Asnaani, A., Litz, B. T., & Hofmann, S. G. (2011). Gender differences in anxiety disorders: prevalence, course of illness, comorbidity and burden of illness. *J Psychiatr Res*, *45*(8), 1027-1035. doi:10.1016/j.jpsychires.2011.03.006
- Moller, C., Wiklund, L., Sommer, W., Thorsell, A., & Heilig, M. (1997). Decreased experimental anxiety and voluntary ethanol consumption in rats following central but not basolateral amygdala lesions. *Brain Res*, *760*(1-2), 94-101.
- Moore, C. B., Guthrie, E. H., Huang, M. T., & Taxman, D. J. (2010). Short hairpin RNA (shRNA): design, delivery, and assessment of gene knockdown. *Methods Mol Biol*, *629*, 141-158. doi:10.1007/978-1-60761-657-3_10
- Moreira, C. M., Masson, S., Carvalho, M. C., & Brandao, M. L. (2007). Exploratory behaviour of rats in the elevated plus-maze is differentially sensitive to inactivation of the basolateral and central amygdaloid nuclei. *Brain Res Bull*, *71*(5), 466-474. doi:10.1016/j.brainresbull.2006.10.004
- Paxinos, G., & Franklin, K. B. J. (1997). *The mouse brain in stereotaxic coordinates*. San Diego: Academic Press.
- Perry, B. L., Pescosolido, B. A., Bucholz, K., Edenberg, H., Kramer, J., Kuperman, S., . . . Nurnberger, J. I., Jr. (2013). Gender-specific gene-environment interaction in alcohol dependence: the impact of daily life events and GABRA2. *Behavior Genetics*, *43*(5), 402-414. doi:10.1007/s10519-013-9607-9
- Pirker, S., Schwarzer, C., Wieselthaler, A., Sieghart, W., & Sperk, G. (2000). GABA(A) receptors: immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience*, *101*(4), 815-850.
- Rassnick, S., Heinrichs, S. C., Britton, K. T., & Koob, G. F. (1993). Microinjection of a corticotropin-releasing factor antagonist into the central nucleus of the amygdala reverses anxiogenic-like effects of ethanol withdrawal. *Brain Res*, *605*(1), 25-32.

- Roberto, M., Gilpin, N. W., & Siggins, G. R. (2012). The central amygdala and alcohol: role of gamma-aminobutyric acid, glutamate, and neuropeptides. *Cold Spring Harb Perspect Med*, 2(12), a012195. doi:10.1101/cshperspect.a012195
- Roberto, M., Madamba, S. G., Moore, S. D., Tallent, M. K., & Siggins, G. R. (2003). Ethanol increases GABAergic transmission at both pre- and postsynaptic sites in rat central amygdala neurons. *Proc Natl Acad Sci U S A*, 100(4), 2053-2058. doi:10.1073/pnas.0437926100
- Sharko, A. C., Kaigler, K. F., Fadel, J. R., & Wilson, M. A. (2013). Individual differences in voluntary ethanol consumption lead to differential activation of the central amygdala in rats: relationship to the anxiolytic and stimulant effects of low dose ethanol. *Alcohol Clin Exp Res*, 37 Suppl 1, E172-180. doi:10.1111/j.1530-0277.2012.01907.x
- Smith, J. P., & Randall, C. L. (2012). Anxiety and alcohol use disorders: comorbidity and treatment considerations. *Alcohol Res*, 34(4), 414-431.
- Smith, K. S., Engin, E., Meloni, E. G., & Rudolph, U. (2012). Benzodiazepine-induced anxiolysis and reduction of conditioned fear are mediated by distinct GABAA receptor subtypes in mice. *Neuropharmacology*, 63(2), 250-258. doi:10.1016/j.neuropharm.2012.03.001
- Soyka, M., Preuss, U. W., Hesselbrock, V., Zill, P., Koller, G., & Bondy, B. (2008). GABA-A2 receptor subunit gene (GABRA2) polymorphisms and risk for alcohol dependence. *J Psychiatr Res*, 42(3), 184-191. doi:10.1016/j.jpsychires.2006.11.006
- Stewart, R. B., Gatto, G. J., Lumeng, L., Li, T. K., & Murphy, J. M. (1993). Comparison of alcohol-preferring (P) and nonpreferring (NP) rats on tests of anxiety and for the anxiolytic effects of ethanol. *Alcohol*, 10(1), 1-10.
- Stewart, S. A., Dykxhoorn, D. M., Palliser, D., Mizuno, H., Yu, E. Y., An, D. S., . . . Novina, C. D. (2003). Lentivirus-delivered stable gene silencing by RNAi in primary cells. *Rna*, 9(4), 493-501.
- Uhart, M., Weerts, E. M., McCaul, M. E., Guo, X., Yan, X., Kranzler, H. R., . . . Wand, G. S. (2013). GABRA2 markers moderate the subjective effects of alcohol. *Addict Biol*, 18(2), 357-369. doi:10.1111/j.1369-1600.2012.00457.x
- Villafuerte, S., Strumba, V., Stoltenberg, S. F., Zucker, R. A., & Burmeister, M. (2013). Impulsiveness mediates the association between GABRA2 SNPs and lifetime alcohol problems. *Genes Brain Behav*, 12(5), 525-531. doi:10.1111/gbb.12039
- Wallace, G. L., White, S. F., Robustelli, B., Sinclair, S., Hwang, S., Martin, A., & Blair, R. J. (2014). Cortical and subcortical abnormalities in youths with conduct disorder and elevated callous-unemotional traits. *J Am Acad Child Adolesc Psychiatry*, 53(4), 456-465 e451. doi:10.1016/j.jaac.2013.12.008
- Zack, M., Poulos, C. X., Aramakis, V. B., Khamba, B. K., & MacLeod, C. M. (2007). Effects of drink-stress sequence and gender on alcohol stress response dampening in high and low anxiety sensitive drinkers. *Alcohol Clin Exp Res*, 31(3), 411-422. doi:10.1111/j.1530-0277.2006.00322.x

TABLES

TABLES

Table 1. Group sizes overall and for specific analyses.

		<u>All</u>	<u>Exclusions</u>			<u>Used for Analyses</u>		
			GFP	Protein	Behavior	Brain	Behavior	Correlations
Control	NI	11	2	0	-	9	-	-
	0	12	6	0	-	6	-	-
	0.75	12	4	0	-	8	-	-
	1.5	12	3	1	-	8	-	-
Virus	NI	12	3	1	-	8	-	-
	0	12	2	1	-	9	-	-
	0.75	13	4	1	-	8	-	-
	1.5	12	3	1	-	8	-	-
Total		96	27	5	-	64	-	-

		<u>All</u>	<u>Exclusions</u>			<u>Used for Analyses</u>		
			GFP	Protein	Behavior	Brain	Behavior	Correlations
Control	NI	11	2	-	0	-	9	-
	0	12	6	-	0	-	6	-
	0.75	12	4	-	0	-	8	-
	1.5	12	3	-	0	-	9	-
Virus	NI	12	3	-	0	-	9	-
	0	12	2	-	1	-	9	-
	0.75	13	4	-	0	-	9	-
	1.5	12	3	-	0	-	9	-
Total		96	27	-	1	-	68	-

		<u>All</u>	<u>Exclusions</u>			<u>Used for Analyses</u>		
			GFP	Protein	Behavior	Brain	Behavior	Correlations
Control	NI	11	-	0	0	-	-	11
	0	12	-	0	0	-	-	12
	0.75	12	-	0	1	-	-	11
	1.5	12	-	2	0	-	-	10
Virus	NI	12	-	1	0	-	-	11
	0	12	-	1	1	-	-	10
	0.75	13	-	0	0	-	-	13
	1.5	12	-	1	1	-	-	10
Total		96	-	5	3	-	-	88

Note: Mice with multiple exclusions are listed in their leftmost occurring exclusion column.

Table 2. Pearson correlations between behavioral measures overall (Top) and between normalized (Z-Score) CeA $\alpha 2$ expression and behavioral measures in naïve mice and within each ethanol dose (Bottom). Behavioral measures in parentheses were excluded from additional analyses. $^{\wedge}p < .05$, $*p < .01$, $**p < .001$ uncorrected for multiple comparisons. Bold values are significant at the more stringent criteria of $p < .01$.

	Entries Total	%Entries OA	Time Center	Time CA	Time OA	Avg. Time Center	Avg. Time CA	Avg. Time OA	(1st Entry Latency)	Head Dips	% Protected HD	(Fecal Boli)
Entries Total	1											
%Entries OA	-0.01	1										
Time Center	0.31 *	-0.52 **	1									
Time CA	-0.13	-0.61 **	0.12	1								
Time OA	-0.05	0.75 **	-0.59 **	-0.87 **	1							
Avg. Time Center	-0.49 **	-0.38 **	0.59 **	0.06	-0.34 *	1						
Avg. Time CA	-0.67 **	0.20	-0.39 **	0.43 **	-0.15	0.17	1					
Avg. Time OA	-0.51 **	0.24 $^{\wedge}$	-0.39 **	-0.56 **	0.65 **	0.26 $^{\wedge}$	0.02	1				
(1st Entry Latency)	-0.11	0.05	0.24 $^{\wedge}$	-0.10	-0.03	0.43 **	0.04	0.19	1			
Head Dips	0.21	0.57 **	-0.22 $^{\wedge}$	-0.49 **	0.51 **	-0.27	-0.09	0.14	0.03	1		
% Protected HD	0.03	-0.64 **	0.60 **	0.45 **	-0.66 **	0.43 **	-0.11	-0.36 *	0.06	-0.47 **	1	
(Fecal Boli)	-0.21 $^{\wedge}$	-0.05	-0.19	0.05	0.05	0.10	0.09	0.35 *	0.12	-0.16	0.02	1
CeA $\alpha 2$ - NI	0.27	-0.37	0.19	0.45 $^{\wedge}$	-0.52 $^{\wedge}$	-0.01	-0.17	-0.46 $^{\wedge}$	-0.22	-0.32	0.22	0.03
CeA $\alpha 2$ - 0 g/kg	0.09	0.19	0.15	-0.23	0.09	0.11	-0.07	0.09	0.34	-0.01	-0.12	0.25
CeA $\alpha 2$ - 0.75 g/kg	0.01	-0.19	-0.08	0.25	-0.16	-0.12	0.00	-0.11	0.36	-0.26	0.13	-0.14
CeA $\alpha 2$ - 1.5 g/kg	-0.42	-0.19	-0.15	0.19	-0.11	0.15	0.28	0.05	0.07	-0.12	0.22	-0.01

FIGURES

FIGURES




Figure 1. RNA interference via shRNA. Illustration taken from (Jolliff, 2007) and description synthesized from (Hutvagner & Zamore, 2002; Moore, Guthrie, Huang, & Taxman, 2010; S. A. Stewart et al., 2003). Lentiviral transduction delivers a *Gabra2*-targeting genetic sequence which is integrated with a cell's DNA. The cell is induced to synthesize a double-stranded RNA (dsRNA) comprised of sense and anti-sense sequences linked by a short hairpin loop (shRNA), with the anti-sense sequence being complementary to endogenous *Gabra2* mRNA. The enzyme Dicer cleaves shRNA into small interfering RNA (siRNA) which is incorporated into a protein complex to form an RNA-induced silencing complex (RISC). RISC uses siRNA as a guide to cleave mRNA that is complementary to the siRNA, in this case *Gabra2*.




Figure 2. Sandwich ELISA procedure. Illustration adapted from LifeSpan Biosciences, Inc. (lsbio.com). A) The wells on each ELISA plate come pre-coated with a *Gabra2*-specific capture antibody (monoclonal, developed using rats). B) Samples and standards are loaded into wells and incubated, and the excess liquid is removed. $\alpha 2$ protein should remain bound to the capture antibody. C) A second *Gabra2*-specific detection antibody (polyclonal, developed using rabbits), conjugated with horseradish peroxidase (HRP), is added to each well and incubated, and the wells are subsequently washed of unbound antibody. D) HRP substrate solution, containing tetramethylbenzidine, is added to each well and incubated, resulting in blue color within each well proportional to $\alpha 2$ protein concentration. A stop solution is then added, ceasing color development. Finally, the optical density of each sample is determined via spectrophotometry at the 450nm wavelength using a microplate reader.




Figure 3. EPM behaviors in mice receiving 0, 0.75, 1.0, or 1.5 g/kg ethanol in Aim 1 (n 's = 10, 12, 9, 11 (left to right)). A) All doses of ethanol increased % of entries in open arms. However, there was no significant effect of dose on either (B) time spent in open arms or (C) total arm entries. * $p < .05$, ** $p < .01$ vs. 0 g/kg.




Figure 4. Lentiviral incorporation in the CeA. A) CeA and surrounding areas under 5x magnification in bright field. C) The same section under 5x magnification with fluorescence showing GFP expression in the CeA. B) A section under 5x magnification showing accurate microinjector placement in the CeA; however, this brain showed no GFP expression. D) Mouse brain atlas with the CeA highlighted for reference.




Figure 5. CeA $\alpha 2$ protein concentrations. A) CeA $\alpha 2$ protein concentrations differed across ELISA passes (n 's = 3, 8, 8, 12 (left to right)). B) Normalized (Z-Score) $\alpha 2$ concentrations were lower for Virus mice than Control mice (n 's = 31, 33 (left to right)).




Figure 6. Arm entries made by naïve mice and mice receiving 0, 0.75, or 1.5 g/kg ethanol (n 's = 9, 9, 6, 9, 8, 9, 9, 9 (left to right)). A) There was no effect of viral condition or dose on total arm entries for naïve mice or mice receiving injection. B) In naïve mice or mice receiving injection, there was no effect of viral condition on % of entries in open arms. However, % of entries in open arms was increased for mice receiving 0.75 or 1.5 g/kg ethanol. * $p < .05$, *** $p < .001$ vs. 0 g/kg.




Figure 7. Total and average time spent in center, closed arms, and open arms by naïve mice and mice receiving 0, 0.75, or 1.5 g/kg ethanol (n 's = 9, 9, 6, 9, 8, 9, 9, 9 (left to right)). In naïve mice, there was no effect of viral condition on any time-related measure (A-F). In mice receiving injection, 1.5 g/kg ethanol reduced time spent in center (A) and increased time spent in open arms (C). Also in mice receiving injection, Virus mice showed a trend ($p = 0.059$) towards increased time spent in closed arms (B) and showed an increase in average time spent in closed arms (E) compared to Control mice. There was no effect of viral condition or dose on average time spent in center (D) or open arms (F). * $p < .05$, ** $p < .01$ vs. 0 g/kg, ^ $p < .05$ vs. Control.




Figure 8. Head dips and % protected head dips made by naïve mice and mice receiving 0, 0.75, or 1.5 g/kg ethanol (n 's = 9, 9, 6, 9, 8, 9, 9, 9 (left to right)). In naïve mice, there was no effect of viral condition on head dips or % protected head dips (A-B). However, in mice receiving injection, 1.5 g/kg ethanol increased head dips (A) and decreased % protected head dips (B). * $p < .05$, ** $p < .01$ vs. 0 g/kg.




Figure 9. Correlations between $\alpha 2$ protein concentration and behavioral measures in naïve mice. Normalized (Z-Score) $\alpha 2$ protein concentration was positively correlated with time spent in closed arms (A) and negatively correlated with both total (B) and average (C) time spent in open arms. Regression lines incorporate both Control and Virus data points ($n = 11$ for each viral condition)

VITA

VITA

Michael P. Smoker

Education

- 1999 – 2003 **B.S., Mathematics**
Marian University, Indianapolis, IN
- 2011 – 2013 **Non-Degree, Psychology**
IUPUI, Indianapolis, IN
- 2013 – Present **Graduate Student, Addiction Neuroscience**
IUPUI, Indianapolis, IN

Teaching Experience

- 2004 – 2005 Mathematics Teacher
North Vermillion Jr/Sr High School, Cayuga, IN
- 2008 – 2009 Adjunct Mathematics Instructor
Ivy Tech Community College, Indianapolis, IN
- 2005 – 2010 Teacher/Tutor
Sylvan Learning Center, Indianapolis, IN & Phoenix, AZ
- Spring 2012 Teaching Assistant – Behavioral Neuroscience
- Summer 2012 Teaching Assistant – Statistics
- Fall 2013 Teaching Assistant – Behavioral Neuroscience, Statistics
- Spring 2014 Instructor – Statistics Lab
Teaching Assistant – Drugs and Behavior
- Summer 2014 Teaching Assistant – Alcoholism and Drug Abuse
- Fall 2014 Teaching Assistant – Foundations of Neuroscience, Learning
- Spring 2015 Instructor – Statistics Lab
- Summer 2015 Instructor – Statistics Lab
Teaching Assistant – Behavioral Neuroscience, Foundations of Neuroscience

Research Experience

- 2011 – 2012 **Research Assistant – Simon Katner, Ph.D.**
 Institute of Psychiatric Research, IUPUI, Indianapolis, IN
- Handling/weighing rats
 - Solution/equipment preparation – ethanol, nicotine, aCSF, HPLC mobile phase, pH, sipper tubes, microdialysis probes
 - Taking/running samples – microdialysate, tail blood, nicotine extraction, HPLC
 - Data recording/analysis – HPLC, Excel
 - Stereotaxic surgery and cannula implantation
 - Histology – cryostat slicing, probe placement verification
- 2012 – 2013 **Research Assistant – Christopher Lapish, Ph.D.**
 Department of Psychology, IUPUI, Indianapolis, IN
- Handling/weighing rats, IP injections, euthanization
 - Solution preparation – ethanol, sucrose
 - Behavioral tasks – catalepsy, social interaction, 2-way drinking, delay discounting
 - Delay discounting – development, experimental design, programming (Med PC)
 - Data recording/analysis – Excel, R
- 2013 – Present **Graduate Student – Stephen Boehm II, Ph.D.**
 Department of Psychology, IUPUI, Indianapolis, IN
- Handling/weighing mice, IP injections, retro-orbital blood sampling, vaginal-cell sampling, brain extraction, tissue punching, euthanization
 - Microscope cell visualization, BEC analysis, histology, laser microdissection
 - Drug preparation/administration
 - Stereotaxic surgery, cannula implantation, site-specific microinjection
- Graduate Student – Christopher Lapish, Ph.D.**
 Department of Psychology, IUPUI, Indianapolis, IN
- Delay discounting in P and Wistar rats

Awards

Research Society on Alcoholism Student Merit Award 2015 (\$230)
 School of Science Graduate Student Council Travel Award 2015 (\$500)

Published Manuscripts

Katner SN, Toalston JE, **Smoker MP**, Rodd ZA, McBride WJ, & Engleman EA (2015). Time-course of Extracellular Nicotine and Cotinine Levels in Rat Brain Following Administration of Nicotine: Effects of Route and Ethanol Co-administration. *Psychopharmacology* 232(3), 551-560.

Linsenbardt DN, **Smoker MP**, Janetsian SJ, & Lapish CC (In Revision). Impulsivity in Rodents with a Genetic Predisposition for Excessive Alcohol Consumption is Associated with a Lack of a Prospective Strategy.

Fritz BM, Kasten CR, Quoilin C, **Smoker MP**, Boehm SL (Submitted). Binge Co-Consumption of Caffeine and Ethanol Increases Ethanol Intake in Adolescent and Adult Mice and Produces Age-Specific Motor Effects.

Published Abstracts – Poster Presentations

Smoker MP, Linsenbardt DN, Oberlin BG, & Lapish CC (2013). P rats show greater discounting than Wistars in a sucrose pellet-reinforced delay discounting task. Poster presented at the Indianapolis Chapter of the Society for Neuroscience conference, Indianapolis, IN.

Smoker MP, Linsenbardt DN, Lukic NA, Oberlin BG, & Lapish CC (2014). P rats show greater discounting than Wistars in a sucrose pellet-reinforced delay discounting task. Poster presented at the Research Society on Alcoholism Conference, Bellevue, WA.

Smoker MP, Linsenbardt DN, & Lapish CC (2014). Differences in impulsivity and prospective-like memory between P rats and Wistars in a sucrose pellet-reinforced delay discounting task. Poster presented at the Indianapolis Chapter of the Society for Neuroscience conference, Indianapolis, IN.

Smoker MP, Zhang Y, Liang T, & Boehm SL (2015). Effect of Gabra2 knockdown in the nucleus accumbens shell on ethanol intake and preference in C57BL/6J mice. Poster presented at the Research Society on Alcoholism Conference, San Antonio, TX.